

# A Bijective Proof of an Identity Concerning Nodes of Fixed Degree in Planted Trees

John W. Moon

Department of Mathematical Sciences  
University of Alberta  
Edmonton, AB Canada T6G 2G1  
email: jwmoon@vega.math.ualberta.ca

Helmut Prodinger

Mathematics Department  
University of the Witwatersrand  
P.O. Wits  
2050 Johannesburg  
South Africa  
email: helmut@gauss.cam.wits.ac.za

## 1 Introduction

We consider *planted plane trees*, which are *rooted*, embedded into the *plane* and enumerated by Catalan numbers  $y_n = \frac{1}{n} \binom{2n-2}{n-1}$ ; see e.g. [2].

In [1], among other things, the following was considered. Denote by  $y_{nd}$  the number of (planted plane) trees of size  $n$  and root degree  $d$ . (It is easy to see that  $y_{nd} = \frac{d}{n-1} \binom{2n-3-d}{n-2}$ , a *ballot number*.) Also, denote by  $D_{nd}$  the total number of nodes of degree  $d$  in trees of size  $n$ . It was shown by generating function arguments that  $D_{nd} = 2 \binom{2n-3-d}{n-2}$  whence

$$d D_{nd} = 2(n-1)y_{nd}. \tag{1}$$

It is this formula (1) that we will “explain” combinatorially in the present note.

**Example.**



**Figure 1.**

$$n = 4, y_4 = 5, y_{41} = 2, y_{42} = 2, y_{43} = 1, D_{41} = 12, D_{42} = 6, D_{43} = 2.$$

**Remark.** In the case of labelled trees, the equivalent of formula (1) reads simply  $D_{nd} = n y_{nd}$ .

## 2 The Bijection

Consider the  $D_{nd}$  trees of size  $n$  with root node  $r$  and a designated node  $u$  of degree  $d$ .

Let  $\mathcal{L}$  denote the set of  $dD_{nd}$  (partially coloured) trees obtained by assigning (a) the colour red to the node  $r$  and to the left-most edge incident with  $r$  and (b) the colour blue to the designated node  $u$  and to one of the  $d$  edges incident with  $u$ . (Note that it is possible for a node or an edge to receive both colours.)

Now consider the  $y_{nd}$  trees of size  $n$  in which the root node has degree  $d$ .

Let  $\mathcal{R}$  denote the set of  $2(n-1)y_{nd}$  (partially coloured) trees obtained by assigning (a) the colour blue to the root node and to the left-most edge incident with the root node and (b) the colour red to one of the  $n-1$  edges of the tree and to one of the two nodes incident with this red edge.

To see that the sets  $\mathcal{L}$  and  $\mathcal{R}$  are equivalent, it suffices to regard each tree in  $\mathcal{L}$  as being rooted at its blue node with the blue edge as the left-most edge incident with the root. (Or, conversely, we may regard each tree in  $\mathcal{R}$  as being rooted at its red node with the red edge as the left-most edge incident with the root.)

**Remark.** Let  $c_0 (= 1)$ ,  $c_1, c_2, \dots$  denote a sequence of constants and suppose each rooted plane tree  $T$  is assigned the weight

$$w(T) = \prod_{v \in V(T)} c_{d(v)}$$

where  $V(T)$  denotes the set of nodes of  $T$  and  $d(v)$  denotes the number of edges incident with node  $v$  that lead away from the root of the tree. Let  $N_\beta(T)$  denote the number of nodes of degree  $\beta$  in a tree  $T$  and let

$$N_n(\alpha, \beta) = \sum N_\beta(T) w(T)$$

where the sum is over all (weighted rooted plane) trees  $T$  of size  $n$  and root degree  $\alpha$ . Then the relation

$$\beta c_\beta c_{\alpha-1} N_n(\alpha, \beta) = \alpha c_\alpha c_{\beta-1} N_n(\beta, \alpha) \quad (2)$$

follows by a slight extension of the bijective argument used above.

## References

- [1] A. Meir and J.W. Moon, Survival under random coverings of trees, *Graphs and Combinatorics* 4 (1988), 49–65.
- [2] R. Sedgewick and P. Flajolet, *An Introduction to the Analysis of Algorithms*, Addison-Wesley, 1996.