

A note on graphs without k -connected subgraphs

Raphael Yuster
Department of Mathematics
University of Haifa at Oranim
Tivon 36006, Israel.
e-mail: raphy@research.haifa.ac.il

Abstract

Given integers $k \geq 2$ and $n \geq k$, let $c(n, k)$ denote the maximum possible number of edges in an n -vertex graph which has no k -connected subgraph. It is immediate that $c(n, 2) = n - 1$. Mader [2] conjectured that for every $k \geq 2$, if n is sufficiently large then $c(n, k) \leq (1.5k - 2)(n - k + 1)$, where equality holds whenever $k - 1$ divides n . In this note we prove that when n is sufficiently large then $c(n, k) \leq \frac{193}{120}(k - 1)(n - k + 1) < 1.61(k - 1)(n - k + 1)$, thereby coming rather close to the conjectured bound.

1 Introduction

All graphs considered here are finite, undirected and have no loops or multiple edges. For the standard terminology used the reader is referred to [1]. This paper is about a classical extremal problem in graph connectivity, raised by Mader in [3]. Let $k \geq 2$ be an integer. Recall that a graph with $n \geq k + 1$ vertices is k -connected if the removal of any set of $k - 1$ vertices from the graph results in a connected subgraph (graphs with $n \leq k$ vertices are considered non k -connected). For $n \geq k$, let $c(n, k)$ denote the maximum possible number of edges in an n -vertex graph which has no k -connected subgraph. It is easy to see that $c(n, 2) = n - 1$ since any tree does not have a 2-connected subgraph, and any n -vertex graph with n edges contains a cycle, which is a 2-connected subgraph. For the rest of this paper we shall assume $k \geq 3$, whenever necessary. Trivially, $c(k, k) = \binom{k}{2}$. Since the complete graph K_{k+1} is the only k -connected graph with $k + 1$ vertices, one has $c(k + 1, k) = \binom{k+1}{2} - 1$ where the unique extremal graph is K_{k+1}^- (the complete graph missing one edge).

In [2], Mader gave a construction of an n -vertex graph with no k -connected subgraph, and with a rather large number of edges. Let $G_{n,k}$ be defined as follows. Assume $n = (k - 1)q + r$ where $1 \leq r \leq k - 1$. The vertices of $G_{n,k}$ are arranged in $q + 1$ classes V_0, \dots, V_q , where each class contains exactly $k - 1$ vertices, except for the final class V_q which contains r vertices. V_0 is an independent set, and V_i is a complete graph for $i = 1, \dots, q$. Furthermore, there is an edge between each vertex of V_0 and each vertex of V_i for $i \geq 1$. Note that V_0 is a disconnecting set of size $k - 1$. It is thus easy to check that $G_{n,k}$ has no k -connected subgraph. Let $e(n, k)$ denote the number of edges of $G_{n,k}$. We have:

$$e(n, k) = (q-1) \binom{k-1}{2} + \binom{r}{2} + (k-1)(n-k+1) \leq \left(\frac{3}{2}k-2\right)(n-k+1), \quad (1)$$

and equality is obtained whenever n is a multiple of $k - 1$. It follows that $c(n, k) \geq e(n, k)$. Mader [2] has conjectured the following:

Conjecture 1.1 (Mader [2]) *For n sufficiently large, $c(n, k) \leq \left(\frac{3}{2}k - 2\right)(n - k + 1)$. Consequently, if n is a multiple of $k - 1$ then $c(n, k) = \left(\frac{3}{2}k - 2\right)(n - k + 1)$, and $G_{n,k}$ is an extremal graph.*

Mader [3] has proved Conjecture 1.1 for all $k \leq 7$. The reason that n needs to be sufficiently large in Conjecture 1.1 follows from the fact that there exist n -vertex graphs with more than $\left(\frac{3}{2}k - 2\right)(n - k + 1)$ edges, and with no k -connected subgraph, for $n = \Theta(k^2)$.

A simple upper bound showing that $c(n, k) < (2k-3)(n-k+1)$ whenever $n \geq 2k - 1$ is presented in [1], p. 45. Mader showed that for n sufficiently large, $c(n, k) < (1 + \sqrt{2}/2)(k - 1)(n - k + 1)$. In this note we present a further improvement which is about halfway between Mader's bound and the bound in Conjecture 1.1:

Theorem 1.2 *For $k \geq 3$ and for $n \geq \frac{9}{4}(k - 1)$, $c(n, k) \leq \frac{193}{120}(k - 1)(n - k + 1)$.*

2 Proof of Theorem 1.2

An (S, A, B) -partition of a non k -connected graph G is a partition of the vertex set of G into three parts S , A and B , where $|S| = k - 1$, $|A| \leq |B|$ and there is no edge connecting a vertex of A and a vertex of B . Clearly, every non k -connected graph with at least $k + 1$ vertices has an (S, A, B) -partition. Given an (S, A, B) -partition, let G_A and G_B denote the subgraphs of G induced by $S \cup A$ and $S \cup B$ respectively.

Proof of Theorem 1.2: Matula has proved [4] that

$$c(n, k) \leq \binom{n}{2} - \frac{(n - k + 1)^2 - 1}{3}. \quad (2)$$

We shall use this fact. For completeness, we reprove (2). This is done by induction on n . For $n = k$, (2) is obvious. For $n = k + 1$ we have $c(k + 1, k) = \binom{k+1}{2} - 1$, so (2) holds. Assume it holds for all $k \leq a < n$. Let G be an n -vertex graph without a k -connected subgraph. Consider an (S, A, B) -partition of G . Clearly, G misses at least the $|A||B|$ possible edges between A and B , and by the induction hypothesis, G_B , as a subgraph of G with $|B| + k - 1 < n$ vertices, misses at least $(|B|^2 - 1)/3$ additional edges. Hence, since $|A| \leq |B|$:

$$e(G) \leq \binom{n}{2} - |A||B| - (|B|^2 - 1)/3 \leq \binom{n}{2} - \frac{(|A| + |B|)^2 - 1}{3} = \binom{n}{2} - \frac{(n - k + 1) - 1}{3}.$$

This proves (2) for all $n \geq k$.

Now let $n \geq \frac{9}{4}(k-1)$, and let G be an n -vertex graph without a k -connected subgraph. Put $n = \gamma(k - 1)$ and assume first that $\gamma \leq \frac{17}{5}$. According to (1) we have:

$$e(G) \leq \frac{\gamma^2(k-1)^2 - \gamma(k-1)}{2} - \frac{(\gamma-1)^2(k-1)^2 - 1}{3} \leq$$

$$(k-1)^2 \left(\frac{\gamma^2}{2} - \frac{(\gamma-1)^2}{3} \right) \leq \frac{193}{120}(\gamma-1)(k-1)^2 = \frac{193}{120}(k-1)(n-k+1).$$

Now assume that $\gamma > \frac{17}{5}$. We use induction once again, and assume the theorem holds for each value smaller than n . Consider an (S, A, B) -partition of G , put $a = |A|$ and $b = |B|$, and recall that $a \leq b$. Let α and β be defined by $a = \alpha(k - 1)$ and $b = \beta(k - 1)$. Notice that $a + b + k - 1 = n$ and so $\alpha + \beta = \gamma - 1$. Consider first the case $\alpha \leq 1$. In this case, $\beta + 1 \geq \frac{12}{5}$, so the induction hypothesis holds for G_B . Hence, the number of edges of G is at most

$$\frac{a(a-1)}{2} + a(k-1) + \frac{193}{120}(k-1)b < 1.5(k-1)a + \frac{193}{120}(k-1)b <$$

$$\frac{193}{120}(k-1)(a+b) = \frac{193}{120}(k-1)(n-k+1).$$

Now consider the case where $\alpha \geq \frac{5}{4}$. Since $\beta \geq \alpha$ we also have $\beta \geq \frac{5}{4}$. In this case, both G_A and G_B have at least $\frac{9}{4}(k - 1)$ edges, and since $e(G) \leq e(G_A) + e(G_B)$ we have by the induction hypothesis that:

$$e(G) \leq \frac{193}{120}(k-1)a + \frac{193}{120}(k-1)b = \frac{193}{120}(k-1)(n-k+1).$$

We remain with the case where $1 < \alpha < \frac{5}{4}$. A useful observation is the following: For every $1 < \alpha < \frac{5}{4}$:

$$\frac{\alpha^2}{2} - \frac{(\alpha - 1)^2}{3} + \alpha - \frac{193}{120}\alpha \leq 0. \quad (3)$$

Furthermore, the l.h.s. of (3) is monotone increasing in the range $[1, \frac{5}{4}]$. Since there are at most $a(k - 1)$ edges between S and A we have that $e(G) \leq e(A) + a(k - 1) + e(G_B)$. If $\beta \geq \frac{5}{4}$ then, according to (2) applied to $e(A)$ and the induction hypothesis applied to $e(G_B)$, and using (3) we have:

$$\begin{aligned} e(G) &\leq \binom{a}{2} - \frac{(a - k + 1)^2 - 1}{3} + a(k - 1) + \frac{193}{120}b(k - 1) = \\ &\frac{\alpha(k - 1)(\alpha(k - 1) - 1)}{2} - \frac{(\alpha - 1)^2(k - 1)^2 - 1}{3} + \alpha(k - 1)^2 + \frac{193}{120}\beta(k - 1)^2 \leq \\ &(k - 1)^2 \left(\frac{\alpha^2}{2} - \frac{(\alpha - 1)^2}{3} + \alpha + \frac{193}{120}\beta \right) \leq (k - 1)^2 \frac{193}{120}(\alpha + \beta) = \\ &\frac{193}{120}(k - 1)(n - k + 1). \end{aligned}$$

Finally, if $\beta < \frac{5}{4}$ then we can use (2) also for $e(G_B)$ and obtain:

$$\begin{aligned} e(G) &\leq (k - 1)^2 \left(\frac{\alpha^2}{2} - \frac{(\alpha - 1)^2}{3} + \alpha \right) + e(G_B) \leq \\ &(k - 1)^2 \left(\frac{\alpha^2}{2} - \frac{(\alpha - 1)^2}{3} + \alpha + \frac{(\beta + 1)^2}{2} - \frac{\beta^2}{3} \right). \end{aligned}$$

We therefore need to show that:

$$\frac{\alpha^2}{2} - \frac{(\alpha - 1)^2}{3} + \alpha + \frac{(\beta + 1)^2}{2} - \frac{\beta^2}{3} \leq \frac{193}{120}(\alpha + \beta).$$

Since the l.h.s. of (3) is monotone increasing in the selected range, and since $\alpha \leq \beta$, the worst case in the last inequality occurs when $\alpha = \beta$. It therefore suffices to show that:

$$\frac{\alpha^2}{3} + \frac{8}{3}\alpha + \frac{1}{6} \leq \frac{193}{60}\alpha$$

which, in turn, is true for $1 < \alpha < \frac{5}{4}$. □

References

- [1] B. Bollobás, *Extremal Graph Theory*, Academic Press, London, 1978.
- [2] W. Mader, *Connectivity and edge-connectivity in finite graphs*, In: *Surveys in Combinatorics*, B. Bollobás, Ed., Cambridge University Press, London, 1979.
- [3] W. Mader, *Existenz n -fach zusammenhängender teilgraphen in graphen genügend grossen kantendichte*, *Abh. Math. Sem. Hamburg Univ.* 37 (1972), 86-97.
- [4] D. W. Matula, *Ramsey Theory for graph connectivity*, *J. Graph Theory* 7 (1983), 95-105.