

Parity Results for p -Regular Partitions with Distinct Parts

James A. Sellers

Department of Mathematics
Penn State University
107 Whitmore Lab
University Park, PA 16802

sellersj@math.psu.edu

August 6, 2001

Abstract

We consider the partition function $b'_p(n)$, which counts the number of partitions of the integer n into distinct parts with no part divisible by the prime p . We prove the following: Let p be a prime greater than 3 and let r be an integer between 1 and $p-1$, inclusively, such that $24r+1$ is a quadratic nonresidue modulo p . Then, for all nonnegative integers n , $b'_p(pn+r) \equiv 0 \pmod{2}$.

1 Introduction

A **partition** λ of the nonnegative integer n is a nonincreasing sequence of nonnegative integers $\lambda_1, \lambda_2, \dots, \lambda_r$ with $\lambda_1 + \lambda_2 + \dots + \lambda_r = n$. Each value λ_i , $1 \leq i \leq r$, is called a **part** of the partition. The number of partitions of n is counted by the **partition function** $p(n)$.

A partition $\lambda_1, \lambda_2, \dots, \lambda_r$ of n is **p -regular** if no part λ_i , $1 \leq i \leq r$, is divisible by p . The function which enumerates the p -regular partitions of n is often denoted $b_p(n)$. These functions have been the focus of much study in recent years [1], [4], [5]. The function $b_p(n)$ is of particular interest for

prime p , as it yields the number of irreducible p -modular representations of the symmetric group S_n [7].

The function which counts those p -regular partitions of n which consist of distinct parts will be denoted $b'_p(n)$ in this note. Such functions have appeared in a variety of recent works. For example, parity results for $b'_2(n)$, the number of partitions of n into distinct odd parts, were found by Hirschhorn [6]. Moreover, the function $b'_5(n)$ was studied by Andrews, Bessenrodt, and Olsson [3] as it relates to representation theory.

2 Main Results

Our main goal here is to prove the following parity result for b'_p by elementary means:

Theorem 2.1. *Let p be a prime greater than 3 and let r be an integer between 1 and $p-1$, inclusively, such that $24r+1$ is a quadratic nonresidue modulo p . Then, for all nonnegative integers n , $b'_p(pn+r) \equiv 0 \pmod{2}$.*

Before proving Theorem 2.1, we mention two propositions. The proofs of these can be found in [2, Chapter 1].

Proposition 2.2. *The generating function for $p(n)$ is given by*

$$\sum_{n \geq 0} p(n)q^n = \frac{1}{(q; q)_\infty}$$

where $(a; b)_\infty = (1-a)(1-ab)(1-ab^2)(1-ab^3)\dots$

Proposition 2.3 (Euler's Pentagonal Number Theorem).

$$(q; q)_\infty = \sum_{m \in \mathbb{Z}} (-1)^m q^{\frac{3}{2}m^2 - \frac{1}{2}m}$$

With these two tools in hand, we turn to the proof of Theorem 2.1.

Proof. Note that the generating function for $b'_p(n)$ is given by

$$\sum_{n \geq 0} b'_p(n)q^n = \frac{(-q; q)_\infty}{(-q^p; q^p)_\infty}.$$

Then we see that

$$\begin{aligned}
 \sum_{n \geq 0} b'_p(n)q^n &= (-q; q)_\infty \frac{(q^p; q^p)_\infty}{(q^{2p}; q^{2p})_\infty} \\
 &\equiv (q; q)_\infty \frac{(q^p; q^p)_\infty}{(q^p; q^p)_\infty^2} \pmod{2} \quad \text{since } 1 - q \equiv 1 + q \pmod{2} \\
 &= (q; q)_\infty \frac{1}{(q^p; q^p)_\infty} \\
 &= (q; q)_\infty \sum_{k \geq 0} p(k)q^{pk}
 \end{aligned}$$

thanks to Proposition 2.2. But this implies

$$\sum_{n \geq 0} b'_p(n)q^n \equiv \sum_{m \in \mathbb{Z}} q^{\frac{3}{2}m^2 - \frac{1}{2}m} \sum_{k \geq 0} p(k)q^{pk} \pmod{2} \quad (1)$$

by Proposition 2.3.

Now we assume

$$pn + r = \frac{3}{2}m^2 - \frac{1}{2}m + pk$$

for some integers m and k . Then we know

$$r \equiv \frac{3}{2}m^2 - \frac{1}{2}m \pmod{p}.$$

Hence,

$$\begin{aligned}
 24r + 1 &\equiv 36m^2 - 12m + 1 \pmod{p} \\
 &\equiv (6m - 1)^2 \pmod{p}.
 \end{aligned}$$

But this contradicts the assumption that $24r + 1$ is a quadratic nonresidue modulo p . Therefore, $pn + r$ can never be represented as $\frac{3}{2}m^2 - \frac{1}{2}m + pk$ for integers m and k . Thus, by (1), we know

$$b'_p(pn + r) \equiv 0 \pmod{2}.$$

□

We note that, for each prime $p > 3$, Theorem 2.1 guarantees $\frac{p-1}{2}$ different congruences modulo 2 for the function b'_p , which is a very satisfying result.

References

- [1] K. Alladi, Partition identities involving gaps and weights, *Trans. Amer. Math. Soc.* 349, no. 12 (1997), 5001-5019.
- [2] G.E. Andrews, *The Theory of Partitions*, Encyclopedia of Mathematics and its Applications, Volume 2, Addison-Wesley Publishing, Reading, MA, 1976.
- [3] G.E. Andrews, C. Bessenrodt, and J.B. Olsson, A refinement of a partition identity and blocks of some modular characters, *Arch. Math. (Basel)* 66, no. 2 (1996), 101-113.
- [4] B. Gordon and K. Ono, Divisibility of certain partition functions by powers of primes, *The Ramanujan Journal* 1 (1997), 25-34.
- [5] A. Granville and K. Ono, Defect zero p -blocks for finite simple groups, *Trans. Amer. Math. Soc.* 348, no. 1 (1996), 331-347.
- [6] M.D. Hirschhorn, On the residue mod 2 and mod 4 of $p(n)$, *Acta Arithmetica* XXXVIII (1980), 105-109.
- [7] G. James and A. Kerber, *The Representation Theory of the Symmetric Group*, Addison-Wesley, Reading, MA, 1979.

AMS Subject Classification Number: 05A17, 11P83