

# On $(t, k)$ -shredders in $k$ -connected graphs

Zeev Nutov

Department of Computer Science

The Open University of Israel

`nutov@openu.ac.il`

Masao Tsugaki

Department of Mathematical Information Science

Science University of Tokyo

## Abstract

Let  $G = (V, E)$  be a  $k$ -connected graph. For  $t \geq 3$  a subset  $T \subset V$  is a  $(t, k)$ -shredder if  $|T| = k$  and  $G - T$  has at least  $t$  connected components. It is known that the number of  $(t, k)$ -shredders in a  $k$ -connected graph on  $n$  nodes is less than  $2n/(2t - 3)$ . We show a slightly better bound for the case  $k \leq 2t - 3$ .

## 1 Introduction

Let  $G = (V, E)$  be a  $k$ -(node) connected graph, that is,  $G$  is simple and there are  $k$  pairwise internally disjoint paths between every pair of its nodes. For  $T \subseteq V$  the  $T$ -components are the connected components of  $G - T$  and let  $b(T)$  denote the number of  $T$ -components.  $T$  with  $|T| = k$  is: a  $k$ -separator if  $b(T) \geq 2$ , a  $k$ -shredder if  $b(T) \geq 3$ , and a  $(t, k)$ -shredder if  $b(T) \geq t \geq 3$ . Let  $B(t, k, G)$  denote number of  $(t, k)$ -shredders in  $G$ ; note that  $B(3, k, G)$  is just the number of  $k$ -shredders in  $G$ . Let  $B(t, k, n) = \max B(t, k, G)$  where the maximum is taken over all  $k$ -connected graphs  $G$  on  $n$  nodes.

A motivation for studying shredders comes from the node-connectivity augmentation problem, see [3, 1, 5]. Cheriyan and Thurimella [1] showed that

in a  $k$ -connected graph computing the number of  $k$ -separators (which may be roughly  $2^k n^2 / k^2$ ) is  $\#$ -complete, while the number of  $k$ -shredders separating two given nodes  $r, s$  is  $O(n)$  and that they all can be found using one max-flow computation. They also proved that  $B(3, k, n) = O(n^2)$  and conjectured that  $B(3, k, n) \leq n$ . Jordán [4] proved this conjecture, and established a tight bound for  $k \leq 3$ : if  $k \leq 3$  and if  $G$  is  $k$ -connected then  $B(3, k, G) \leq (n - k - 1)/2$  unless  $k = 3$  and  $G = K_{3,3}$ . For arbitrary  $k$ , Egawa [2] proved that  $B(3, k, n) < 2n/3$  and that this bound is (asymptotically) the best possible. Liberman and Nutov [5], and independently the second author of this paper, considered  $(t, k)$ -shredders and proved that  $B(t, k, n) < 2n/(2t - 3)$ .

**Remark:** The following simple example shows that the bound  $B(t, k, n) < 2n/(2t - 3)$  is asymptotically tight for  $k \geq 2(t - 1)$ . Let  $t, q$  be integers. Let  $G$  be  $(t - 1)$ -blow-up of a  $q$ -cycle, that is  $G$  is obtained from a cycle of length  $q$  by replacing every node  $a$  by a set  $V_a$  of  $t - 1$  nodes, and every edge  $ab$  by  $(t - 1)^2$  edges, so that  $V_a \cup V_b$  induces a complete bipartite graph  $K_{t-1, t-1}$ . For  $k = 2(t - 1)$ ,  $G$  is  $k$ -connected and  $n = qk/2 = q(t - 1)$ . Thus  $2n/(2t - 3) = 2q(t - 1)/(2t - 3) = q + q/(2t - 3)$ . On the other hand,  $B(t, k, G) = q$ . For  $2t - 3 = k - 1 > q$ , the above bound is tight. This example easily extends for the case  $k > 2(t - 1)$ , by adding  $k - 2(t - 1)$  nodes to  $G$  and connecting by an edge every added node to all the other nodes.

We show a slightly better bound for the case  $k \leq 2t - 3$ , and prove the following theorem:

**Theorem** Let  $k \leq 2t - 3$ . Then  $B(t, k, n) \leq (n - k - 1)/(t - 1)$  for  $n \geq 2k + 1$  and  $B(t, k, n) < n/(t - 1)$  for  $n \leq 2k$ .

**Remark:** Our bound generalizes the bound of Jordán [4] which states: For  $k \leq 3$  and  $t = 3$ ,  $B(t, k, G) \leq (n - k - 1)/(t - 1)$  unless  $k = 3$  and  $G = K_{3,3}$ .

Indeed, let  $t = 3$  and let  $k \leq 3$ . Then  $k \leq 2t - 3$  since  $t = 3$ . Our bound implies that  $B(t, k, G) \leq (n - k - 1)/(t - 1)$  for  $n \geq 2k + 1$ . For  $n \leq 2k \leq 6$ , an easy case analysis shows that this bound also holds, unless  $k = 3$  and

$$G = K_{3,3}.$$

The bound in the Theorem is sharp for  $n \geq 2k + 1$ , in the sense that there are infinitely many graphs that attain this bound. Let  $p$  be an integer, and  $k, t$  be as in the Theorem. Define a graph  $G = (V, E)$  with  $n = |V| = k + t \sum_{1 \leq i \leq p} (t - 1)^{i-1}$  by:

$$\begin{aligned} V &= \{a\} \\ &\cup \{b_{i,j,h} : 1 \leq i \leq t, 1 \leq j \leq p, 1 \leq h \leq (t - 1)^{j-1}\} \\ &\cup \{c_\ell : 1 \leq \ell \leq k - 1\} \\ E &= \{ab_{i,1,1}, b_{i,j,h}b_{i,j+1,\ell} | 1 \leq i \leq t, 1 \leq j \leq p - 1, \\ &\quad 1 \leq h \leq (t - 1)^{j-1}, (h - 1)(t - 1) + 1 \leq \ell \leq h(t - 1)\} \\ &\cup \{c_i c_j | 1 \leq i < j \leq k - 1\} \\ &\cup \{c_\ell a, c_\ell b_{i,j,h} | 1 \leq \ell \leq k - 1, 1 \leq i \leq t, 1 \leq j \leq p, \\ &\quad 1 \leq h \leq (t - 1)^{j-1}\}. \end{aligned}$$

Then  $G$  is  $k$ -connected and has  $1 + t \sum_{1 \leq i \leq p-1} (t - 1)^{i-1}$   $(t, k)$ -shredders which are:

$$\begin{aligned} &\{a, c_1, \dots, c_{k-1}\} \\ &\{b_{i,j,h}, c_1, \dots, c_{k-1}\} \quad 1 \leq i \leq t, 1 \leq j \leq p - 1, 1 \leq h \leq (t - 1)^{j-1}. \end{aligned}$$

Thus

$$\begin{aligned} \frac{n - k - 1}{t - 1} &= \frac{1}{t - 1} (k + t \sum_{1 \leq i \leq p} (t - 1)^{i-1} - k - 1) \\ &= \frac{1}{t - 1} (t(t - 1) \sum_{1 \leq i \leq p} (t - 1)^{i-2} - 1) \\ &= \frac{1}{t - 1} (t(t - 1) \sum_{2 \leq i \leq p} (t - 1)^{i-2} + t(t - 1)(t - 1)^{-1} - 1) \\ &= \frac{1}{t - 1} (t(t - 1) \sum_{1 \leq i \leq p-1} (t - 1)^{i-1} + t - 1) \\ &= 1 + t \sum_{1 \leq i \leq p-1} (t - 1)^{i-1} = B(t, k, G) \end{aligned}$$

## 2 Properties of separators and shredders

Let  $G = (V, E)$  be a  $k$ -connected graph. For  $Y \subseteq V$  let  $\Gamma(Y)$  denote the set of neighbors of  $Y$  in  $G$ , and let  $Y^* = V - Y - \Gamma(Y)$ .  $Y$  is *tight* if  $|\Gamma(Y)| = k$  and  $Y^* \neq \emptyset$ . A separator  $S$  *meshes* a separator  $T$  if  $S$  intersects at least two  $T$ -components. As was mentioned in [1], if  $S$  meshes  $T$ , then each one of  $S, T$  intersects all the components of the other; thus “meshing” is a symmetric relation. The following statement is immediate.

**Proposition 2.1** *Let  $S, T$  be distinct nonmeshing  $k$ -separators in a  $k$ -connected graph. Then there is an  $S$ -component  $X$  and a  $T$ -component  $Y$  so that  $T \subset X \cup S$  and  $S \subset Y \cup T$  holds; thus  $Y^* \subset X$  and  $X^* \subset Y$ .*

**Corollary 2.2** *Let  $\mathcal{T}$  be a family of pairwise nonmeshing  $k$ -separators in a  $k$ -connected graph  $G$ . Then  $G$  has a node  $r$  not belonging to any member of  $\mathcal{T}$ .*

**Proof:** Let  $\mathcal{C}$  be the family of tight sets obtained by picking the  $T$ -components for each  $T \in \mathcal{T}$ . Let  $X$  be an inclusion minimal set in  $\mathcal{C}$ , and let  $S = \Gamma(X)$ . We claim that no member of  $\mathcal{T}$  intersects  $X$ . Suppose this is not so, that is, there is  $T \in \mathcal{T}$  intersecting  $X$ . Then  $T \subset X \cup S$ , since  $S, T$  are nonmeshing. By Proposition 2.1, there is a  $T$ -component strictly contained in  $X$ , contradicting the minimality of  $X$ .  $\square$

**Lemma 2.3** *Let  $S, T$  be meshing  $k$ -separators in a  $k$ -connected graph  $G = (V, E)$  so that  $S \cup T \neq V$ . Then  $k \geq b(S) + b(T) - 2$ .*

**Proof:** Let  $t = b(T)$  and  $s = b(S)$ . Let  $Y$  be the union of  $T$ -components not containing  $r$ , and let  $Z$  be the union of  $S$ -components not containing  $r$ . Since  $S, T$  mesh,  $|\Gamma(Z) \cap Y| \geq t - 1$ ,  $|\Gamma(Y) \cap Z| \geq s - 1$ . Let  $W = Y^* \cap Z^*$ . Then  $r \in W^* \neq \emptyset$ . Thus  $|\Gamma(W)| \geq k$ , since  $G$  is  $k$ -connected. Furthermore,

$$\begin{aligned} |\Gamma(W)| &= |\Gamma(Y^* \cap Z^*)| \\ &\leq |\Gamma(Y^*)| + |\Gamma(Z^*)| - [|\Gamma(Y^*) \cap Z| + |\Gamma(Z^*) \cap Y|] \\ &\leq 2k - [(s - 1) + (t - 1)]. \end{aligned}$$

Thus we have  $k \leq 2k - [(s - 1) + (t - 1)]$ , that is  $k \geq s + t - 2$ .  $\square$

For  $r \in V$  let  $B_r(t, k, G)$  be the number of  $(t, k)$ -shredders in  $G$  not containing  $r$ . The following statement follows from a simple averaging argument, e.g., see [5, Lemma 2.4].

**Lemma 2.4**  $B(t, k, G) \leq \frac{n}{n-k} \max_{r \in V} B_r(t, k, G)$ . If  $r$  is a node of  $G$  not contained in any  $(t, k)$ -shredder then  $B(t, k, G) = B_r(t, k, G)$ .

Two intersecting sets  $X, Y$  are *crossing* (or  $Y$  *crosses*  $X$ ) if none of them contains the other. We will use the following key statement (see [6, Lemma 3.14] and [5, Lemma 2.3]).

**Lemma 2.5** ([6, 5]) *Let  $G$  be a  $k$ -connected graph, let  $T$  be a  $k$ -shredder in  $G$ , and let  $Y$  be a tight set in  $G$  so that  $Y^*$  intersects some  $T$ -component  $C$ . Then  $Y$  does not cross  $V - T - C$  nor a  $T$ -component distinct from  $C$ .*

### 3 Proof of the Theorem

Let  $r \in V$ . Consider the family  $\mathcal{L}$  obtained by picking for every  $(t, k)$ -shredders  $T$  the  $T$ -components that do not contain  $r$  and their union; color the former blue and the later red. Let  $U$  be the union of the sets in  $\mathcal{L}$ ; note that  $|U| \leq n - |\Gamma(r)| - 1 \leq n - k - 1$ . By Lemma 2.5,  $\mathcal{L}$  is laminar (that is, if two sets in  $\mathcal{L}$  intersect then one of them contains the other). Thus  $\mathcal{L}$  can be represented by a forest  $\mathcal{F}$  of rooted trees, if we order the sets in  $\mathcal{L}$  by inclusion:  $X$  is a child of  $Y$  if  $X$  is the largest set in  $\mathcal{L}$  properly contained in  $Y$ . Note that every red set is the union of its children. The forest  $\mathcal{F}$  has the following properties:

- (i) every member of  $\mathcal{L}$  is either blue or red, but not both;
- (ii) the children of every red set are blue, and there are at least  $t - 1$  of them;
- (iii) every child of a blue set is red.

**Claim 3.1** *If a blue set  $Z$  is the union of its children, then for every child  $Q$  of  $Z$  there exists a child  $R$  of  $Z$  so that  $S = \Gamma(Q)$  and  $T = \Gamma(R)$  are meshing. In particular, if  $Z$  has one child, then  $Z$  contains a node not contained in its children.*

**Proof:** Let  $Q$  be a child of  $Z$ . Since  $S \neq \Gamma(Z)$  and  $Q \subseteq Z$ , and since  $Z$  is the union of its children,  $Q$  has a neighbor in some child  $R$  of  $Z$ . Consequently,  $Q$  has a child  $X$  and  $R$  has a child  $Y$ , so that there is an edge in  $G$  with one end in  $X$  and the other end in  $Y$ . This implies that  $S$  and  $T$  mesh. Otherwise, by Proposition 2.1,  $Y^* \subset X$ ; this is a contradiction, since  $r \in Y^* - X$ .  $\square$

**Claim 3.2** *If every blue set has a node not contained in any of its children then  $B_r(t, k, G) \leq (n - k - 1)/(t - 1)$ .*

**Proof:** Let  $\ell$  be the number of blue sets. Then  $\ell \leq |U| \leq n - k - 1$ , since every blue set has a node not contained in any of its children. We will show that the number of red sets (which equals  $B_r(t, k, G)$ ) is at most  $\ell/(t - 1)$ . We claim that in any tree  $\mathcal{T}$  (and thus in any forest) that satisfies properties (i),(ii),(iii), the number of red nodes is at most  $\ell/(t - 1)$ . If  $\mathcal{T}$  has one red node, the statement is obvious. Otherwise,  $\mathcal{T}$  has a blue node  $X$  so that every red descendant of  $X$  is a child of  $X$ . Let  $q$  be the number of children of  $X$ . By deleting the children of  $X$  and their descendants (which are all blue leaves) we get a tree with the same properties, and  $\ell$  decreases by at least  $q(t - 1)$ . The claim follows.  $\square$

Combining Corollary 2.2 and Lemma 2.4 with the two claims above, we get:

**Corollary 3.3** *If no two  $(t, k)$ -shredders mesh, then  $B(t, k, G) \leq (n - k - 1)/(t - 1)$ .*

**Proof of the Theorem** By Lemma 2.3, if  $S, T$  are meshing  $(t, k)$ -shredders, then  $S \cup T = V$  and thus  $n \leq 2k$ . Thus for  $n \geq 2k + 1$  no two  $(t, k)$ -shredders mesh, and Corollary 3.3 implies the bound  $B(t, k, G) \leq (n - k - 1)/(t - 1)$ .

Assume  $n \leq 2k$ . Let  $r \in V$  and consider the corresponding forest  $\mathcal{F}$ . We claim that every blue set  $X$  has a node not contained in any of its children; thus by Claim 3.2  $B_r(t, k, G) \leq (n - k - 1)/(t - 1)$ , implying (via Lemma 2.4)  $B(t, k, G) < n/(t - 1)$ . Otherwise, by Claim 3.1,  $X$  has two (red) children  $Y, Z$  corresponding to meshing shredders. But then by Lemma 2.3  $k \geq 2t - 2$ , contradicting the assumption of the theorem.

## References

- [1] J. Cheriyan and R. Thurimella, Fast algorithms for  $k$ -shredders and  $k$ -node connectivity augmentation, *J. Algorithms* **33** (1999), No. 1, 15–50.
- [2] Y. Egawa,  $k$ -shredders in  $k$ -connected graphs, manuscript.
- [3] T. Jordán, On the optimal vertex-connectivity augmentation, *J. Combinatorial Theory Series B* **63** (1995), 8–20.
- [4] T. Jordán, On the number of shredders, *J. Graph Theory* **31** (1999), 195–200.
- [5] G. Liberman and Z. Nutov, On shredders and vertex-connectivity augmentation, manuscript.
- [6] Z. Nutov, Approximating rooted connectivity augmentation problems, manuscript.