

# A Note on Graceful Graphs with Large Chromatic Numbers

Ahmad Mahmoody

*Department of Mathematical Sciences, Sharif University of Technology,*

*Azadi Street, P. O. Box 11365-9415, Tehran, Iran*

*Email address: ahmadmahmoody@yahoo.com*

## Abstract

A *graceful labeling* of a graph  $G$  with  $m$  edges is a function  $f : V(G) \rightarrow \{0, \dots, m\}$  such that distinct vertices receive distinct numbers and  $\{|f(u) - f(v)| : uv \in E(G)\} = \{1, \dots, m\}$ . A graph is *graceful* if it has a graceful labeling. In [1] this question was posed: "Is there an  $n$ -chromatic graceful graph for  $n \geq 6$ ?" In this paper it is shown that for any natural number  $n$ , there exists a graceful graph  $G$  with  $\chi(G) = n$ .

For a graph  $G$  we denote the set of vertices and the set of edges of  $G$  with  $V(G)$  and  $E(G)$ , respectively. The chromatic number of a graph  $G$ , denoted by  $\chi(G)$  is the minimum number of independent subsets into which  $V(G)$  can be partitioned. A *graceful labeling* of a graph  $G$  with  $m$  edges is a function  $f : V(G) \rightarrow \{0, \dots, m\}$  such that distinct vertices receive distinct numbers and  $\{|f(u) - f(v)| : uv \in E(G)\} = \{1, \dots, m\}$ . A graph is *graceful* if it has a graceful labeling. The label of an edge is the difference between the labels of its ends. In [2, p. 266] it has been conjectured that


**Conjecture.** Graceful graphs with arbitrary large chromatic number do not exist.

In the following theorem we give a negative answer to the above conjecture.

**Theorem.** For any natural number  $n$ , there exists a graceful graph  $G$  such that  $\chi(G) = n$ .

**Proof.** For  $n = 1$  the assertion is trivial, thus suppose that  $n \geq 2$ . Let  $v_1, \dots, v_n$  be the vertices of the complete graph  $K_n$ . For each  $i, 1 \leq i \leq n$ , consider  $2^i$  as the label of  $v_i$ . First, we show that all edges of  $K_n$  in this labeling have different labels. Suppose  $v_i v_j$  and  $v_k v_l$  are two edges with the same labels. First assume that  $i > j$  and  $k > l$ . Thus the labels of  $v_i v_j$  and  $v_k v_l$  are  $2^i - 2^j$  and  $2^k - 2^l$ , respectively, and clearly they are distinct unless  $i = k$  and  $j = l$ . This implies that two edges  $v_i v_j$  and  $v_k v_l$  are the same.

Now the greatest label of the vertices is  $2^n$  and it is obvious that for each natural number  $n$ , we have  $2^n > \frac{n(n-1)}{2}$ . Add  $2^n - \frac{n(n-1)}{2}$  new vertices to the complete graph  $K_n$  and join all of them to  $v_n$ . Let us call this graph by  $G$ . We claim that  $G$  has a graceful labeling. For each  $x, x \in \{1, \dots, 2^n\}$



which does not occur as a label of an edge in  $K_n$  label one of the new vertices with  $2^n - x$ . It is obvious that all edges have different labels. Moreover the labels of vertices are contained in  $\{1, \dots, 2^n\}$  and they are distinct. Indeed if the labels of two vertices are the same, then we conclude that the labels of two edges (of which one end is  $v_n$ ) are the same, a contradiction. Note that  $n$  is an arbitrary number, and  $\chi(G) = n$  ( $n \geq 2$ ).  $\square$

**Acknowledgement.** The author wishes to thank Dr. Saieed Akbari for introducing the conjecture and for his fruitful comments.

## References

- [1] G. Chartrand, H. Hevia, O. R. Oellermann, The Chromatic Number of a Factorization of a Graph, Bull. Inst. Combin. Appl. 20(1997), 33-56.
- [2] G. Chartrand and L. Lesniak, Graphs & Digraphs, CHAPMAN & HALL/CRC, Fourth Edition, 2005.