

# Minimum Distance between Bent and 1-resilient Boolean Functions \*

Soumen Maity<sup>†</sup> and Subhamoy Maitra<sup>‡</sup>

## Abstract

In this paper we study the minimum distance between the set of bent functions and the set of 1-resilient Boolean functions and present lower bounds on that. The first bound is proved to be tight for functions up to 10 input variables and a revised bound is proved to be tight for functions up to 14 variables. As a consequence, we present a strategy to modify the bent functions, by toggling some of its outputs, in getting a large class of 1-resilient functions with very good nonlinearity and autocorrelation. In particular, the technique is applied up to 14-variable functions and we show that the construction provides a large class of 1-resilient functions reaching currently best known nonlinearity and achieving very low autocorrelation values which were not known earlier. The technique is sound enough to theoretically solve some of the mysteries of 8-variable, 1-resilient functions with maximum possible nonlinearity. However, the situation becomes complicated from 10 variables and above, where we need to go for complicated combinatorial analysis with trial and error using computational facility.

**Keywords:** Autocorrelation, Bent Function, Boolean Function, Non-linearity, Resiliency.

## 1 Introduction

Boolean functions have many applications in computer security practices including the construction of key stream generators based on a set of shift

---

\*This is a revised version of the paper presented in the Fast Software Encryption - FSE 2004, Lecture Notes in Computer Science, volume 3017, pages 143–160, Springer Verlag, 2004.

<sup>†</sup>Department of Mathematics, Indian Institute of Technology Guwahati, Guwahati 781 039, Assam, INDIA, Email: soumen@iitg.ernet.in

<sup>‡</sup>Applied Statistics Unit, Indian Statistical Institute, 203, B T Road, Kolkata 700 108, INDIA, Email: subho@isical.ac.in

registers. A Boolean function is said to be correlation immune if its output leaks no information about its input values. Balanced correlation immune functions are called resilient functions. Construction of resilient Boolean functions with very good parameters in terms of nonlinearity, algebraic degree and other cryptographic parameters has received lot of attention in literature [17, 18, 20, 21, 8, 23, 2, 3]. In [19, 7], it had been shown how bent functions can be modified to construct highly nonlinear balanced Boolean functions. A recent construction method [13] presents modification of some output points of a bent function to construct highly nonlinear 1-resilient function. A natural question that arises in this context is “at least how many bits in the output column of the truth table of a bent function need to be changed to get the output column of an 1-resilient Boolean function”. The answer of this question gives the minimum distance between the set of bent functions and the set of 1-resilient functions (we define this distance more formally in Subsection 1.1). We here try to answer this question and show that the minimum distance for  $n$ -variable functions is

$$dBR_n(1) \geq 2^{\frac{n}{2}-1} + 2 \left\lceil \frac{(r+1)(2^{\frac{n}{2}-1} - \sum_{i=0}^r \binom{n}{i}) + \sum_{i=1}^r i \binom{n}{i}}{n-r-1} \right\rceil,$$

where  $r$  is the integer such that  $\sum_{i=0}^r \binom{n}{i} \leq 2^{\frac{n}{2}-1} + 1 < \sum_{i=0}^{r+1} \binom{n}{i}$  is satisfied. We also show that this result is tight for  $n \leq 10$ . We observe that this bound is not tight for  $n \geq 12$  and provide further modification (see Subsection 5.1) to the bound that makes the new bound tight for  $n = 12, 14$ .

The immediate corollary is the construction of 1-resilient Boolean functions with nonlinearity  $\geq 2^{n-1} - 2^{\frac{n}{2}-1} - dBR_n(1)$  and maximum absolute value of autocorrelation spectra  $\leq 4dBR_n(1)$ . Interestingly, it is possible to get 1-resilient functions with better nonlinearity and autocorrelation than these bounds. In particular, we concentrate on construction of 1-resilient Boolean functions up to 14-variables with best known nonlinearity and autocorrelation. *Throughout the paper we consider the number of input variables ( $n$ ) is even.*

The bent functions chosen in [13, Section 3] use the concept of perfect nonlinear functions and one example function each for 8, 10 and 12 variables were presented. However, it is not clear how a generalized construction of such bent functions can be achieved in that manner. We here identify a large subclass of Maiorana-McFarland type bent functions which can be modified to get 1-resilient functions with currently best known parameters. Further our construction is superior to [13] in terms of number of points that need to be toggled (we need less in case of 10, 12 variables), the nonlinearity (we get better nonlinearity for 12 variables) and autocorrelation (we get 1-resilient functions with autocorrelation values that were not known earlier for 10, 12 and 14 variables).

## 1.1 Preliminaries

A Boolean function on  $n$  variables may be viewed as a mapping from  $\{0, 1\}^n$  into  $\{0, 1\}$ . A Boolean function  $f(x_1, \dots, x_n)$  is also interpreted as the output column of its *truth table*  $f$ , i.e., a binary string of length  $2^n$ ,

$$f = [f(0, 0, \dots, 0), f(1, 0, \dots, 0), f(0, 1, \dots, 0), \dots, f(1, 1, \dots, 1)].$$

The *Hamming distance* between two binary strings  $S_1, S_2$  is denoted by  $d(S_1, S_2)$ , i.e.,  $d(S_1, S_2) = \#(S_1 \neq S_2)$ . Also the *Hamming weight* or simply the weight of a binary string  $S$  is the number of ones in  $S$ . This is denoted by  $wt(S)$ . An  $n$ -variable function  $f$  is said to be *balanced* if its output column in the truth table contains equal number of 0's and 1's (i.e.,  $wt(f) = 2^{n-1}$ ). By distance between two Boolean functions on  $n$  variables  $f, g$ , we mean the  $d(f, g)$ , where  $f, g$  are interpreted as their truth tables, i.e., binary strings of length  $2^n$ .

Denote addition operator over  $GF(2)$  by  $\oplus$ . An  $n$ -variable Boolean function  $f(x_1, \dots, x_n)$  can be considered to be a multivariate polynomial over  $GF(2)$ . This polynomial can be expressed as a sum of products representation of all distinct  $k$ -th order products ( $0 \leq k \leq n$ ) of the variables. More precisely,  $f(x_1, \dots, x_n)$  can be written as

$$a_0 \oplus \bigoplus_{1 \leq i \leq n} a_i x_i \oplus \bigoplus_{1 \leq i < j \leq n} a_{ij} x_i x_j \oplus \dots \oplus a_{12\dots n} x_1 x_2 \dots x_n,$$

where the coefficients  $a_0, a_i, a_{ij}, \dots, a_{12\dots n} \in \{0, 1\}$ . This representation of  $f$  is called the *algebraic normal form* (ANF) of  $f$ . The number of variables in the highest order product term with nonzero coefficient is called the *algebraic degree*, or simply the degree of  $f$  and denoted by  $deg(f)$ .

A Boolean function is called *affine* if it has an ANF of the form  $a_0 \oplus a_1 x_1 \oplus \dots \oplus a_n x_n$ . An affine function with constant term equal to zero is called a *linear* function. The set of all  $n$ -variable affine (respectively linear) functions is denoted by  $A(n)$  (respectively  $L(n)$ ). The nonlinearity of an  $n$ -variable function  $f$  is

$$nl(f) = \min_{g \in A(n)} d(f, g),$$

i.e., the distance from the set of all  $n$ -variable affine functions.

Let  $x = (x_1, \dots, x_n)$  and  $\omega = (\omega_1, \dots, \omega_n)$  both belong to  $\{0, 1\}^n$  and

$$x \cdot \omega = x_1 \omega_1 \oplus \dots \oplus x_n \omega_n.$$

Let  $f(x)$  be a Boolean function on  $n$  variables. Then the *Walsh transform* of  $f(x)$  is a real valued function over  $\{0, 1\}^n$  which is defined as

$$W_f(\omega) = \sum_{x \in \{0, 1\}^n} (-1)^{f(x) \oplus x \cdot \omega}.$$

In terms of Walsh spectra, the nonlinearity of  $f$  is given by

$$nl(f) = 2^{n-1} - \frac{1}{2} \max_{\omega \in \{0,1\}^n} |W_f(\omega)|.$$

For  $n$ -even, the maximum nonlinearity of a Boolean function can be  $2^{n-1} - 2^{\frac{n}{2}-1}$  and the functions possessing this nonlinearity are called bent functions [16]. Further, for a bent function  $f$  on  $n$  variables,  $W_f(\omega) = \pm 2^{\frac{n}{2}}$  for all  $\omega$ . For  $n$  even, let us denote the set of  $n$ -variable bent functions by  $\mathcal{B}_n$ .

In [9], an important characterization of correlation immune and resilient functions has been presented, which we use as the definition here. A function  $f(x_1, \dots, x_n)$  is  $m$ -resilient (respectively  $m$ -th order correlation immune) iff its Walsh transform satisfies  $W_f(\omega) = 0$ , for  $0 \leq wt(\omega) \leq m$  (respectively  $W_f(\omega) = 0$ , for  $1 \leq wt(\omega) \leq m$ ).

As the notation used in [17, 18], by an  $(n, m, d, \sigma)$  function we denote an  $n$ -variable,  $m$ -resilient function with degree  $d$  and nonlinearity  $\sigma$ . Let us denote the set of  $n$ -variable  $m$ -resilient functions by  $\mathcal{R}_{n,m}$ . Thus we can formally define

$$dBR_n(m) = \min_{f \in \mathcal{B}_n, g \in \mathcal{R}_{n,m}} d(f, g).$$

We will now define *restricted Walsh transform* which will be frequently used in this text. The *restricted Walsh transform* of  $f(x)$  on a subset  $S$  of  $\{0, 1\}^n$  is a real valued function over  $\{0, 1\}^n$  which is defined as

$$W_f(\omega)|_S = \sum_{x \in S} (-1)^{f(x) \oplus x \cdot \omega}.$$

Now we present the following technical result.

**Proposition 1** *Let  $S \subset \{0, 1\}^n$  and  $b(x), f(x)$  be two  $n$ -variable Boolean functions such that  $f(x) = 1 \oplus b(x)$  when  $x \in S$  and  $f(x) = b(x)$  otherwise. Then  $W_f(\omega) = W_b(\omega) - 2W_b(\omega)|_S$ .*

**Proof:** Take  $\omega \in \{0, 1\}^n$ . Now  $W_f(\omega)$

$$\begin{aligned} &= \sum_{x \in \{0,1\}^n} (-1)^{f(x) \oplus \omega \cdot x} \\ &= \sum_{x \in \{0,1\}^n - S} (-1)^{f(x) \oplus \omega \cdot x} + \sum_{x \in S} (-1)^{f(x) \oplus \omega \cdot x} \\ &= \sum_{x \in \{0,1\}^n - S} (-1)^{b(x) \oplus \omega \cdot x} - \sum_{x \in S} (-1)^{b(x) \oplus \omega \cdot x} \\ &\quad \text{(since } f, b \text{ are same for the inputs } \notin S \\ &\quad \text{and complement when the inputs } \in S) \\ &= \sum_{x \in \{0,1\}^n - S} (-1)^{b(x) \oplus \omega \cdot x} + \sum_{x \in S} (-1)^{b(x) \oplus \omega \cdot x} \\ &\quad - 2 \sum_{x \in S} (-1)^{b(x) \oplus \omega \cdot x} \\ &= \sum_{x \in \{0,1\}^n} (-1)^{b(x) \oplus \omega \cdot x} - 2 \sum_{x \in S} (-1)^{b(x) \oplus \omega \cdot x} \\ &= W_b(\omega) - 2W_b(\omega)|_S. \quad \blacksquare \end{aligned}$$

Propagation Characteristics (PC) and Strict Avalanche Criteria (SAC) [15] are important properties of Boolean functions to be used in S-boxes. Further, Zhang and Zheng [24] identified related cryptographic measures called Global Avalanche Characteristics (GAC).

Let  $\alpha \in \{0, 1\}^n$  and  $f$  be an  $n$ -variable Boolean function. Define the autocorrelation value of  $f$  with respect to the vector  $\alpha$  as

$$\Delta_f(\alpha) = \sum_{x \in \{0,1\}^n} (-1)^{f(x) \oplus f(x \oplus \alpha)},$$

and the absolute indicator

$$\Delta_f = \max_{\alpha \in \{0,1\}^n, \alpha \neq 0} |\Delta_f(\alpha)|.$$

A function is said to satisfy PC( $k$ ), if  $\Delta_f(\alpha) = 0$  for  $1 \leq wt(\alpha) \leq k$ . Note that, for a bent function  $f$  on  $n$  variables,  $\Delta_f(\alpha) = 0$  for all nonzero  $\alpha$ , i.e.,  $\Delta_f = 0$ .

Analysis of autocorrelation properties of correlation immune and resilient Boolean functions has gained substantial interest recently as evident from [22, 25, 11, 4]. In [11, 4], it has been identified that some well known construction of resilient Boolean functions are not good in terms of autocorrelation properties. Since the present construction is modification of bent functions which possess the best possible autocorrelation properties, we get very good autocorrelation properties of the 1-resilient functions. We present a bound on the  $\Delta_f$  value of the 1-resilient functions and further achieve best known autocorrelation values for the cases  $n = 8, 10, 12$ .

## 2 The Distance

Initially we start with a simple technical result.

**Proposition 2**  $dBR_n(1) \geq 2^{\frac{n}{2}-1}$ .

**Proof :** For a bent function  $b$  on  $n$  variables,  $W_b(\omega) = \pm 2^{\frac{n}{2}}$ . Hence the minimum distance from a bent function to balanced functions equals  $2^{\frac{n}{2}-1}$ . The 1-resilient functions are balanced by definition and hence the result. ■

Now we present a restricted result. Let  $b(x)$  be an  $n$ -variable bent function with  $W_b(\omega) = +2^{\frac{n}{2}}$  for  $wt(\omega) \leq 1$ . We denote by  $M_b(n, 1)$  the minimum number of bits to be modified in the output column of  $b(x)$  to construct an  $n$  variable 1-resilient function from  $b(x)$ .

**Theorem 1** Let  $b(x)$  be an  $n$ -variable bent function with  $W_b(\omega) = 2^{\frac{n}{2}}$  for  $0 \leq wt(\omega) \leq 1$ . Then

$$M_b(n, 1) \geq 2^{\frac{n}{2}-1} + 2 \left\lceil \frac{(r+1)(2^{\frac{n}{2}-1} - \sum_{i=0}^r \binom{n}{i}) + \sum_{i=1}^r i \binom{n}{i}}{n-r-1} \right\rceil,$$

where  $r$  is the integer such that  $\sum_{i=0}^r \binom{n}{i} \leq 2^{\frac{n}{2}-1} + 1 < \sum_{i=0}^{r+1} \binom{n}{i}$  is satisfied.

**Proof :** Let  $S \subset \{0, 1\}^n$  and  $f(x)$  be an  $n$ -variable Boolean function obtained by modifying the  $b(x)$  values for  $x \in S$  and keeping the other bits unchanged. Then from Proposition 1,  $W_f(\omega) = W_b(\omega) - 2W_b(\omega)|_S \forall \omega$ , and in particular,  $W_f(\omega) = 2^{\frac{n}{2}} - 2W_b(\omega)|_S$  for  $0 \leq wt(\omega) \leq 1$ .

It is known that,  $f$  is 1-resilient iff  $W_f(\omega) = 0$  for  $0 \leq wt(\omega) \leq 1$ , i.e., iff  $W_b(\omega)|_S = 2^{\frac{n}{2}-1}$  for  $0 \leq wt(\omega) \leq 1$ . Thus, our problem is to find a lower bound on  $|S| = k$  with the constraint  $W_b(\omega)|_S = 2^{\frac{n}{2}-1}$  for  $0 \leq wt(\omega) \leq 1$ .

Given  $S = \{x^{i_1}, x^{i_2}, \dots, x^{i_k}\} \subset \{0, 1\}^n$ , consider the matrices

$$\mathbf{S}^{k \times n} = (x^{i_1}, x^{i_2}, \dots, x^{i_k})^T, \quad b(\mathbf{S})^{k \times 1} = (b(x^{i_1}), b(x^{i_2}), \dots, b(x^{i_k}))^T,$$

$$\text{and } (\mathbf{S} \oplus b(\mathbf{S}))^{k \times n} = (x^{i_1} \oplus b(x^{i_1}), x^{i_2} \oplus b(x^{i_2}), \dots, x^{i_k} \oplus b(x^{i_k}))^T.$$

By  $A^T$  we mean transpose of a matrix  $A$ . Also by abuse of notation,  $x^{i_j} \oplus b(x^{i_j})$  means the GF(2) addition (XOR) of the bit  $b(x^{i_j})$  with each of the bits of  $x^{i_j}$ .

Now  $W_b(\omega)|_S = 2^{\frac{n}{2}-1}$  for  $0 \leq wt(\omega) \leq 1$  implies that there are exactly  $\frac{k}{2} - 2^{\frac{n}{2}-2}$  many 1's in  $b(\mathbf{S})$  and in each column of  $\mathbf{S} \oplus b(\mathbf{S})$ . Since all the rows of  $\mathbf{S}$  are distinct and further  $b(\mathbf{S})$  contains  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  many 0's,  $\mathbf{S} \oplus b(\mathbf{S})$  should contain at least  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  distinct rows.

Consider that one such matrix  $\mathbf{S} \oplus b(\mathbf{S})$  is formed. The number of 1's in the matrix is exactly  $n \times (\frac{k}{2} - 2^{\frac{n}{2}-2})$  as each column contains exactly  $\frac{k}{2} - 2^{\frac{n}{2}-2}$  many 1's and there are  $n$  columns. We know that there must be at least  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  many distinct rows. Thus the total number of 1's in these distinct rows must be  $\leq n \times (\frac{k}{2} - 2^{\frac{n}{2}-2})$ . Note that the minimum number of 1's in  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  many distinct rows is at least

$$\sum_{i=1}^r i \binom{n}{i} + (r+1) \left( \frac{k}{2} + 2^{\frac{n}{2}-2} - \sum_{i=0}^r \binom{n}{i} \right)$$

(all the rows upto weight  $r$  and some of the rows with weight  $r+1$ ). Hence,

$$\sum_{i=1}^r i \binom{n}{i} + (r+1) \left( \frac{k}{2} + 2^{\frac{n}{2}-2} - \sum_{i=0}^r \binom{n}{i} \right) \leq n \times \left( \frac{k}{2} - 2^{\frac{n}{2}-2} \right).$$

This gives,

$$k \geq 2 \left[ \frac{(n+r+1)2^{\frac{n}{2}-2} + \sum_{i=1}^r i \binom{n}{i} - (r+1) \sum_{i=0}^r \binom{n}{i}}{n-r-1} \right].$$

Now we discuss how to choose this  $r$ . For this we need a easier lower bound on  $k$  which does not depend on  $r$  itself.

From Proposition 2,  $k \geq 2^{\frac{n}{2}-1}$ . We now show that  $k \geq 2^{\frac{n}{2}-1} + 2$ . This is because, to construct an 1-resilient function from bent function, the number of 1's in each column must be  $\geq 1$  (it cannot be 0 since then we will not be able to get distinct rows). As number of 1's in each column is  $\frac{k}{2} - 2^{\frac{n}{2}-2}$ , we get  $\frac{k}{2} - 2^{\frac{n}{2}-2} \geq 1$ , and hence  $k \geq 2^{\frac{n}{2}-1} + 2$ .

Since,  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  number of distinct rows has to be filled, we need to find the  $r$  such that  $\sum_{i=0}^r \binom{n}{i} \leq \frac{k}{2} + 2^{\frac{n}{2}-2} < \sum_{i=0}^{r+1} \binom{n}{i}$ . Putting the minimum value of  $k$ , i.e.,  $2^{\frac{n}{2}-1} + 2$ , we get  $r$  such that  $\sum_{i=0}^r \binom{n}{i} \leq 2^{\frac{n}{2}-1} + 1 < \sum_{i=0}^{r+1} \binom{n}{i}$ . ■

As example, for  $n = 8$ , take  $r = 1$  and  $9 = \sum_{i=0}^r \binom{n}{i} \leq$  (the  $=$  of  $\leq$  is satisfied here)  $2^{\frac{n}{2}-1} + 1 = 9 < \sum_{i=0}^{r+1} \binom{n}{i}$  is satisfied. For  $n = 10$ , take  $r = 1$  and  $11 = \sum_{i=0}^r \binom{n}{i} \leq$  (the  $<$  of  $\leq$  is satisfied here)  $2^{\frac{n}{2}-1} + 1 = 17 < \sum_{i=0}^{r+1} \binom{n}{i}$  is satisfied.

**Theorem 2** Let  $b(x)$  be any  $n$ -variable bent function. Then

$$dBR_n(1) \geq 2^{\frac{n}{2}-1} + 2 \left\lceil \frac{(r+1)(2^{\frac{n}{2}-1} - \sum_{i=0}^r \binom{n}{i}) + \sum_{i=1}^r i \binom{n}{i}}{n-r-1} \right\rceil,$$

where  $r$  is the integer such that  $\sum_{i=0}^r \binom{n}{i} \leq 2^{\frac{n}{2}-1} + 1 < \sum_{i=0}^{r+1} \binom{n}{i}$  is satisfied.

**Proof:** Without loss of generality, assume that  $W_b(\omega) = +2^{\frac{n}{2}}$  for  $wt(\omega) = 0$ . Let  $G_1 = \{\omega | wt(\omega) = 1, W_b(\omega) = +2^{\frac{n}{2}}\}$  and  $G_2 = \{\omega | wt(\omega) = 1, W_b(\omega) = -2^{\frac{n}{2}}\}$ . Let  $S \subset \{0, 1\}^n$  and  $f(x)$  be an  $n$ -variable Boolean function obtained by modifying the  $b(x)$  values for  $x \in S$  and keeping the other bits unchanged. Then from Proposition 1,  $W_f(\omega) = W_b(\omega) - 2W_b(\omega)|_S \forall \omega$ , and in particular,  $W_f(\omega) = 2^{\frac{n}{2}} - 2W_b(\omega)|_S$  for  $wt(\omega) = 0, \omega \in G_1$  and  $W_f(\omega) = -2^{\frac{n}{2}} - 2W_b(\omega)|_S$  for  $\omega \in G_2$ .

Given  $f$  is 1-resilient, we need to find a lower bound on  $|S| = k$  with the constraints  $W_b(\omega)|_S = 2^{\frac{n}{2}-1}$  for  $wt(\omega) = 0$  and  $\omega \in G_1$  and  $W_b(\omega)|_S = -2^{\frac{n}{2}-1}$  for  $\omega \in G_2$ .

Let  $|G_1| = \lambda$ . Using the same argument as in the proof of Theorem 1, our problem is to find a  $k \times n$  binary matrix  $S \oplus b(S)$  with minimum number of rows  $k$  such that there are  $\lambda$  columns with exactly  $\frac{k}{2} - 2^{\frac{n}{2}-2}$  many 1's in each column and exactly  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  many 1's in each of the remaining  $n - \lambda$  columns. Further, there are at least  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  distinct rows.

Let  $M_1^{k \times \lambda}$  (respectively  $M_2^{k \times (n-\lambda)}$ ) be a binary matrix with exactly  $\frac{k}{2} - 2^{\frac{n}{2}-2}$  (respectively  $\frac{k}{2} + 2^{\frac{n}{2}-2}$ ) many 1's in each column. Let  $J$  be the  $k \times (n - \lambda)$  matrix with all elements 1. Then the problem of "finding a binary matrix  $(M_1 : M_2)$  with minimum number of rows  $k$  such that there are at least  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  distinct rows" is equivalent to "finding a binary matrix

$(M_1 : J - M_2)$  with minimum number of rows  $k$  such that there are at least  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  distinct rows". Note that each column of  $(M_1 : J - M_2)$  contains exactly  $\frac{k}{2} - 2^{\frac{n}{2}-2}$  many 1's. Thus, if one can find a construction with

$$k < 2^{\frac{n}{2}-1} + 2 \left\lceil \frac{(r+1)(2^{\frac{n}{2}-1} - \sum_{i=0}^r \binom{n}{i}) + \sum_{i=1}^r i \binom{n}{i}}{n-r-1} \right\rceil$$

in this case that will contradict to the lower bound presented in Theorem 1. Hence the proof.  $\blacksquare$

For  $8 \leq n \leq 16$ , it can be checked that  $\sum_{i=0}^1 \binom{n}{i} \leq 2^{\frac{n}{2}-1} + 1 < \sum_{i=0}^{1+1} \binom{n}{i}$  is satisfied. In these cases, the lower bound on  $k$  is attained for  $r = 1$  itself. Thus we have the following result.

**Corollary 1** For even  $n$ ,  $8 \leq n \leq 16$ ,  $dBR_n(1) \geq 2^{\frac{n}{2}-1} + 2 \left\lceil \frac{2^{\frac{n}{2}-n-2}}{n-2} \right\rceil$ .

Assume that one can construct a bent function  $b$  on  $n$  variables such that  $dBR_n(1)$  bits at the output column of  $b$  are changed to get an  $n$ -variable 1-resilient function  $f$ . It is clear that toggling of a single bit can reduce the nonlinearity at most by 1 and increase the maximum absolute value of the autocorrelation spectra (absolute indicator) by at most 4. Thus we have the following result.

**Theorem 3**  $nl(f) \geq 2^{n-1} - 2^{\frac{n}{2}-1} - dBR_n(1)$  and  $\Delta_f \leq 4dBR_n(1)$ .

**Proof :** This follows from  $nl(f) \geq nl(b) - dBR_n(1)$  and  $\Delta_f \leq \Delta_b + 4dBR_n(1)$ , where  $b$  is a bent function.  $\blacksquare$

However, for the actual constructions of functions on 8, 10 and 12 variables, we will show that we get better nonlinearity and autocorrelation values than these bounds. Let us now present the case for  $n = 4, 6$ . We are not interested in the case  $n = 2$ , since there is no nonlinear 2-variable 1-resilient functions.

Note that  $r = 0$  for these two cases and then we arrive at  $dBR_4(1) \geq 4$  and  $dBR_6(1) \geq 6$ . These bounds are tight since we can construct 4-variable (respectively 6-variable) 1-resilient function by changing 4 (respectively 6) output points of 4-variable (respectively 6-variable) bent function.

For the 4-variable case, we have the following construction.

$$\mathbf{S} \oplus b(\mathbf{S}) = \begin{pmatrix} x_4 & x_3 & x_2 & x_1 \\ \hline 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ \hline 0 & 1 & 0 & 0 \end{pmatrix}$$

The constraint is that the number of 1's in each column has to be 1 and there are at least 3 distinct rows. Then we take a bent function with truth

table 0000011001010011 and toggle the function at the inputs

$$\{(0, 0, 0, 1), (0, 0, 1, 0), (1, 0, 0, 0), (1, 0, 1, 1)\}.$$

Note that the last row of the matrix is complemented when considered as the input point. The bent function has output zero at the first three points and output one at the last point. This gives a (4, 1, 2, 4) function with the truth table 0110011011000011.

For the 6-variable case, let us consider the following matrix.

$$\mathbf{S} \oplus b(\mathbf{S}) = \begin{pmatrix} x_6 & x_5 & x_4 & x_3 & x_2 & x_1 \\ \hline 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

We take a bent function with truth table

0000000001011010001111000110011001101001001100110101010100001111

(note that the bent function has output zero at the first five points and output one at the last point) and toggle the outputs at the input points

$\{(0, 0, 0, 0, 0, 1), (0, 0, 0, 0, 1, 0), (0, 0, 0, 1, 0, 0), (0, 0, 1, 0, 0, 0), (1, 0, 0, 0, 0, 0), (1, 0, 1, 1, 1, 1)\}$ . Then we get a (6, 1, 4, 24) function with the truth table

0110100011011010001111000110011011101001001100100101010100001111.

Before proceeding further let us note the following issues towards the construction of 1-resilient functions in general.

1. Given even  $n$ , one needs to design a  $k \times n$  binary matrix  $\mathbf{S} \oplus b(\mathbf{S})$  with minimum number of rows  $k$ . We have provided a lower bound in Theorem 2 on  $k$ . However, we could not provide any algorithm to construct such a matrix. For  $n \leq 8$ , the situation can be handled easily, but it becomes complicated for  $n \geq 10$ . We could construct such matrices for  $n = 10, 12$  and  $14$  by trial and error.
2. For the construction, we use the restricted case as described in Theorem 1, where we need a bent function  $b$  with  $W_b(\omega) = 2^{\frac{n}{2}}$  for  $0 \leq wt(\omega) \leq 1$ . It is possible to get Maiorana-McFarland type bent functions with such properties (see the next sections for exact details).

3. We need to partition the rows of the  $k \times n$  binary matrix  $\mathbf{S} \oplus b(\mathbf{S})$  in such a manner that one partition (call it  $P_1$ ) contains  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  many rows and the other partition (call it  $P_2$ ) contains  $\frac{k}{2} - 2^{\frac{n}{2}-2}$  many rows. The rows in each partition are distinct among themselves.
4. Let  $P_2^c$  denotes the set of rows which are bitwise complement to the rows in  $P_2$ . It is clear that the rows of  $P_1$  and  $P_2^c$  taken together are distinct among themselves. Each row can be seen as an input point of an  $n$ -variable Boolean function. We need a bent function such that  $b(x) = 0$  for  $x \in P_1$  and  $b(x) = 1$  for  $x \in P_2^c$ . We show that given the matrix  $\mathbf{S} \oplus b(\mathbf{S})$ , constructing such bent functions from Maiorana-McFarland class is possible for  $n \leq 14$ .
5. Note that this construction needs a closer look when we try to achieve very high nonlinearity. That needs a detailed analysis of the Walsh spectra of the bent functions. This will be discussed in full details in the coming sections.

Next we consider the cases for  $n = 8, 10, 12$  and  $14$  one by one.

### 3 The 8-variable 1-resilient Functions

In the previous section we have presented a lower bound of the minimum distance between the bent and 1-resilient functions and presented an outline how exactly a construction is possible. Further to achieve the currently best known parameters (or even better than that, if possible) we may need to consider some other issues. In this section we consider the construction of an  $(8, 1, 6, 116)$  function. Construction of this function was an important open question and the function has been first reported in [10] by inter linking combinatorial technique and computer search. Later this function has also been found by meta heuristic search (simulated annealing) in [5]. Further the function found in [5] has  $\Delta_f = 24$ , which is currently the best known value. We here follow the similar kind of technique used in [13]. In the course of discussion it will be clear that how our technique is an improvement over [13]. We present a generalized construction method of  $(8, 1, 6, 116)$  functions by modifying Maiorana-McFarland type bent functions and in specific cases, these functions have the  $\Delta_f$  value as low as 24, the best known one [5].

For the 8-variable case, the matrix looks as follows.

$$S \oplus b(S) = \begin{pmatrix} x_8 & x_7 & x_6 & x_5 & x_4 & x_3 & x_2 & x_1 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Note that apart from the conditions mentioned in Theorem 1, we also add one extra condition  $W_b(\omega) = -16$  for  $wt(\omega) = 8$  for the 8-variable bent function  $b$  from which we construct the 1-resilient function. This condition is not required for 1-resiliency of the function, which is handled by the conditions of Theorem 1 itself. However, this extra condition is required to maintain the nonlinearity at 116. If one takes  $W_b(\omega) = 16$  for  $wt(\omega) = 8$ , then the nonlinearity will fall to 112. This is explained in detail in the proof of Theorem 4.

**Construction 1** Take a bent function  $b(x)$  on 8 variables with the following properties : (1)  $b(x) = 0$  for  $wt(x) \leq 1$  and  $b(x) = 1$  for  $wt(x) = 8$ , (2)  $W_b(\omega) = 16$  for  $wt(\omega) \leq 1$  and  $W_b(\omega) = -16$  for  $wt(\omega) = 8$ . Define a set  $S = \{x \in \{0, 1\}^8 \mid wt(x) = 0, 1, 8\}$ . Construct a function  $f(x)$  as :

$$\begin{aligned} f(x) &= 1 \oplus b(x), & \text{if } x \in S \\ &= b(x), & \text{otherwise.} \end{aligned}$$

From Corollary 1, we get that  $dBR_8(1) \geq 10$  and we here choose exactly 10 positions and modify them. It is important to point out that we here start with bent functions with some specific properties. The reason for choosing such bent functions is to get an actual construction of 1-resilient function with very high nonlinearity.

**Theorem 4** The function  $f(x)$  as described in Construction 1 is an  $(8, 1, 6, 116)$  function.

**Proof :** Take  $\omega \in \{0, 1\}^8$  with  $wt(\omega) = i$ . Now

$$\begin{aligned} W_f(\omega) &= \sum_{x \in \{0, 1\}^8} (-1)^{b(x) \oplus \omega \cdot x} - 2 \sum_{x \in S} (-1)^{b(x) \oplus \omega \cdot x} \\ &= W_b(\omega) - 2(8 - 2wt(\omega) + 2(wt(\omega) \bmod 2)). \end{aligned}$$

Now we explain how the last step is deduced. Note that  $b(x) = 0$  when  $wt(x) = 0$  and  $b(x) = 1$ , when  $wt(x) = 8$ . Thus,

$$\begin{aligned} \sum_{x \in \{0,1\}^8 | wt(x)=0,8} (-1)^{b(x) \oplus \omega \cdot x} &= 0, \text{ when } wt(\omega) \text{ is even,} \\ &= 2, \text{ when } wt(\omega) \text{ is odd.} \end{aligned}$$

Moreover,  $\sum_{x \in \{0,1\}^8 | wt(x)=1} (-1)^{b(x) \oplus \omega \cdot x} = 8 - 2wt(\omega)$ , as

- (i)  $b(x) = 0$  when  $wt(x) = 1$  and
- (ii)  $\omega \cdot x = 1$  at  $wt(\omega)$  input points when  $wt(x) = 1$ .

Since  $\sum_{x \in S} (-1)^{b(x) \oplus \omega \cdot x} = \sum_{x \in \{0,1\}^8 | wt(x)=0,8} (-1)^{b(x) \oplus \omega \cdot x} + \sum_{x \in \{0,1\}^8 | wt(x)=1} (-1)^{b(x) \oplus \omega \cdot x}$ , we get,

$$W_f(\omega) = W_b(\omega) - 2 ( 8 - 2wt(\omega) + 2(wt(\omega) \bmod 2) ).$$

When  $wt(\omega) \leq 1$ ,  $W_f(\omega) = W_b(\omega) - 16 = 16 - 16 = 0$ . Thus the function is 1-resilient.

Further, if  $wt(\omega) = 8$ ,  $W_f(\omega) = W_b(\omega) + 16 = -16 + 16 = 0$ . For any other choice, i.e., for  $2 \leq wt(\omega) \leq 7$ , we have  $|8 - 2wt(\omega) + 2(wt(\omega) \bmod 2)| \leq 4$  and hence,  $|W_f(\omega)| \leq |W_b(\omega)| + 8 = 16 + 8 = 24$ . Hence,  $nl(f) = 2^{8-1} - \frac{24}{2} = 116$ .

Since the function attains the maximum possible nonlinearity, the algebraic degree [1, 3] of the function must be  $8 - 2 = 6$ . ■

$wt(\omega)$	0	1	2	3	4	5	6	7	8
$W_f(\omega) = W_b(\omega) +$	-16	-16	-8	-8	0	0	8	8	16

Table 1: Relationship between Walsh spectra of  $f, g$  as described in Construction 1.

Based on Table 1 and the previous discussion, we get related results with respect to (i) nonexistence of some 8-variable bent functions and (ii) some relationship between 8-variable bent functions and balanced Boolean functions with nonlinearity 118 (whose existence is not known till date).

Note that, in the Walsh spectra of a bent function on 8 variables, there are 120 values of +16 and 136 values of -16 or vice versa. It is known that even if that condition is satisfied for some Walsh spectra, the inverse Walsh transform may not produce a Boolean function. We here discuss that issue.

**Lemma 1** Consider a function  $b(x)$  on 8 variables with the properties :

1.  $b(x) = 0$  for  $wt(x) \leq 1$  and  $b(x) = 1$  for  $wt(x) = 8$ ,

2.  $W_b(\omega) = 16$  for  $wt(\omega) \leq 3$  and  $W_b(\omega) = -16$  for  $wt(\omega) \geq 6$ .

*This function can not be bent.*

**Proof :** If such a function  $b$  is bent, then Table 1, we will get an 1-resilient function with nonlinearity 120. This is a contradiction. ■

**Corollary 2** Consider a function  $b(x)$  on 8 variables with the properties :

1.  $b(x) = 0$  for  $wt(x) \leq 3$  and  $b(x) = 1$  for  $wt(x) \geq 6$ ,
2.  $W_b(\omega) = 16$  for  $wt(\omega) \leq 1$  and  $W_b(\omega) = -16$  for  $wt(\omega) = 8$ .

*This function can not be bent.*

**Proof :** The result follows from Lemma 1 and the duality property of bent functions. ■

Next we present an important result related to the existence of balanced 8-variable function with nonlinearity 118.

**Theorem 5** Take a bent function  $h(x)$  on 8 variables with the following properties :

1.  $h(x) = 0$  for  $wt(x) \leq 1$  and  $h(x) = 1$  for  $wt(x) = 8$ ,
2.  $W_h(\omega) = 16$  for  $wt(\omega) \leq 2$  and  $W_h(\omega) = -16$  for  $wt(\omega) \geq 6$ .

Define a set  $T = \{x \in \{0, 1\}^8 | wt(x) = 1\}$ . Construct a function  $g(x)$  as :

$$\begin{aligned} f(x) &= 1 \oplus h(x), & \text{if } x \in T \\ &= h(x), & \text{otherwise.} \end{aligned}$$

*Then  $g$  is a balanced 8-variable function with nonlinearity 118.*

**Proof :** The proof is similar to the proof of Theorem 4. ■

We have tried some heuristic search to find a bent function as mentioned in Theorem 5, but could not get any. Getting such a bent function or proving its nonexistence is an interesting open question.

### 3.1 A subclass of Maiorana-McFarland bent functions

The original Maiorana-McFarland class of bent function is as follows [6]. Consider  $n$ -variable Boolean functions on  $(X, Y)$ , where  $X, Y \in \{0, 1\}^{\frac{n}{2}}$  of the form  $f(X, Y) = X \cdot \pi(Y) + g(Y)$  where  $\pi$  is a permutation on  $\{0, 1\}^{\frac{n}{2}}$  and  $g$  is any Boolean function on  $\frac{n}{2}$  variables. The function  $f$  can be seen as concatenation of  $2^{\frac{n}{2}}$  distinct (upto complementation) affine function on  $\frac{n}{2}$  variables.

Once again we write what kind of bent function  $b(x)$  on 8 variables we require.

1.  $b(x) = 0$  for  $wt(x) \leq 1$  and  $b(x) = 1$  for  $wt(x) = 8$ ,
2.  $W_b(\omega) = 16$  for  $wt(\omega) \leq 1$  and  $W_b(\omega) = -16$  for  $wt(\omega) = 8$ .

In this case,  $n = 8$ , i.e.,  $\frac{n}{2} = 4$ . We have to decide what permutations  $\pi$  on  $\{0, 1\}^4$  and what kind of functions  $g$  on  $\{0, 1\}^4$  we can take such that the conditions on  $b$  are satisfied. We present a set of conditions below, which taken all together, provides sufficient condition for construction of such functions. Before going into the conditions, let us fix the notation and ordering of input variables as  $x = (x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8)$ ,  $X = (X_1, X_2, X_3, X_4)$ , and  $Y = (Y_1, Y_2, Y_3, Y_4)$ . Further we identify  $X_1 = x_1, X_2 = x_2, X_3 = x_3, X_4 = x_4, Y_1 = x_5, Y_2 = x_6, Y_3 = x_7, Y_4 = x_8$ .

1. First of all, the function  $b$  has the value zero at the points  $(0, 0, 0, 0, 0, 0, 0, 0)$ ,  $(1, 0, 0, 0, 0, 0, 0, 0)$ ,  $(0, 1, 0, 0, 0, 0, 0, 0)$ ,  $(0, 0, 1, 0, 0, 0, 0, 0)$ ,  $(0, 0, 0, 1, 0, 0, 0, 0)$  and this condition is satisfied if we choose  $\pi(0, 0, 0, 0) = (0, 0, 0, 0)$  and  $g(0, 0, 0, 0) = 0$ .
2. Next we need function  $b$  should have value 0 at points  $(0, 0, 0, 0, 1, 0, 0, 0)$ ,  $(0, 0, 0, 0, 0, 1, 0, 0)$ ,  $(0, 0, 0, 0, 0, 0, 1, 0)$ ,  $(0, 0, 0, 0, 0, 0, 0, 1)$ , and this condition is satisfied if we choose  $g(Y) = 0$  for  $wt(Y) = 1$ .
3. We need  $b$  to be 1 when the input is  $(1, 1, 1, 1, 1, 1, 1, 1)$ . Thus if  $\pi(1, 1, 1, 1)$  is a vector of odd weight then  $g(1, 1, 1, 1)$  need to be 0. Otherwise if  $\pi(1, 1, 1, 1)$  is a vector of even weight then  $g(1, 1, 1, 1)$  has to be 1.
4. Since we have already decided that  $\pi(0, 0, 0, 0) = (0, 0, 0, 0)$  and  $g(0, 0, 0, 0) = 0$ , the  $W_f(\omega)$  values for  $\omega \in \{(0, 0, 0, 0, 1, 0, 0, 0)$ ,  $(0, 0, 0, 0, 0, 1, 0, 0)$ ,  $(0, 0, 0, 0, 0, 0, 1, 0)$ ,  $(0, 0, 0, 0, 0, 0, 0, 1)\}$  becomes  $+2^{\frac{n}{2}} = 16$ .
5. Further if  $\pi(Y) \in \{(1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 1, 0), (0, 0, 0, 1)\}$ , then we take  $g(Y) = 0$ . This guarantees that  $W_f(\omega)$  values for  $\omega \in \{(1, 0, 0, 0, 0, 0, 0, 0)$ ,  $(0, 1, 0, 0, 0, 0, 0, 0)$ ,  $(0, 0, 1, 0, 0, 0, 0, 0)$ ,  $(0, 0, 0, 1, 0, 0, 0, 0)\}$  becomes  $+2^{\frac{n}{2}} = 16$ .
6. Lastly, if  $\pi(Y) = (1, 1, 1, 1)$ , we have to fix  $g(Y) = (wt(Y) + 1) \bmod 2$ . This guarantees that  $W_f(1, 1, 1, 1, 1, 1, 1, 1) = -2^{\frac{n}{2}} = -16$ .

Given a bent function from the Maiorana-McFarland class  $f(X, Y) = X \cdot \pi(Y) + g(Y)$ , the dual of such function  $f$  is  $Y \cdot \pi^{-1}(X) + g(\pi^{-1}(X))$ . It is interesting to check whether the above points can be replaced by more precise arguments using this idea.

**Theorem 6** Let  $n = 8$ ,  $x \in \{0, 1\}^n$  and  $X, Y \in \{0, 1\}^{\frac{n}{2}}$ . Let  $b(x)$  be a Maiorana-McFarland type bent function  $b(x) = b(X, Y) = X \cdot \pi(Y) + g(Y)$  where  $\pi$  is a permutation on  $\{0, 1\}^{\frac{n}{2}}$  and  $g$  is a Boolean function on  $\frac{n}{2}$  variables with the following conditions.

- (1) if  $Y = (0, 0, 0, 0)$ ,  $\pi(Y) = Y$ ;
- (2) if  $wt(\pi(Y)) \leq 1$ , or  $wt(Y) \leq 1$ , then  $g(Y) = 0$ ;
- (3) if  $Y = (1, 1, 1, 1)$ ,  $g(Y) = (wt(\pi(Y)) + 1) \bmod 2$ ;
- (4) if  $wt(\pi(Y)) = 4$ ,  $g(Y) = (wt(Y) + 1) \bmod 2$ .

Then (1)  $b(x) = 0$  for  $wt(x) \leq 1$  and  $b(x) = 1$  for  $wt(x) = 8$ , (2)  $W_b(\omega) = 16$  for  $wt(\omega) \leq 1$  and  $W_b(\omega) = -16$  for  $wt(\omega) = 8$ .

Further there are  $\geq 2^{46 \cdot 297}$  many distinct  $b$ 's (upto complementation) satisfying these conditions and in turn there are  $\geq 2^{46 \cdot 297}$  many distinct (upto complementation)  $(8, 1, 6, 116)$  functions.

**Proof :** The proof of the properties of  $b$  is discussed above in detail. The count of such functions is arrived as follows. Note that there are  $2^{\frac{n}{2}} = 16$  places for the permutation  $\pi$ .

Let there are  $i$  many  $Y$ 's,  $0 \leq i \leq 4$  such that  $wt(\pi(Y)) = 1$  for  $wt(Y) = 1$ . There are 4 elements of weight 1 and 10 elements of weight 2 or 3. Thus the  $\pi(Y)$ 's for  $wt(Y) = 1$  may be chosen in  $\binom{4}{i} \binom{10}{4-i}$  ways. Note that  $\pi(Y)$  can not be  $(1, 1, 1, 1)$  for  $wt(Y) = 1$ . Now there are two cases.

1. Consider that  $\pi(1, 1, 1, 1) = (1, 1, 1, 1)$ . Then the number of options is  $\binom{4}{i} \cdot \binom{10}{4-i} \cdot 4! \cdot 10! \cdot 2^{6+i}$ . This is because the 4 elements where  $wt(Y) = 1$  can be permuted in  $4!$  ways. The 4 elements where  $wt(Y) = 2, 3$  can be permuted in  $10!$  ways. The function  $g(Y)$  is fixed when  $Y$  is  $(0, 0, 0, 0)$  (1 place,  $g(Y) = 0$ ) or  $wt(Y) = 1$  (4 places,  $g(Y) = 0$ ) or  $wt(\pi(Y)) = 1$  ( $4 - i$  places,  $g(Y) = 0$ ) or  $wt(Y) = wt(\pi(Y)) = 4$  (1 place,  $g(Y) = 1$ ). Thus  $g(Y)$  is fixed in  $10 - i$  places and we can put any choice from  $\{0, 1\}$  for  $16 - (10 - i) = 6 + i$  places.
2. Consider that  $\pi(1, 1, 1, 1) \neq (1, 1, 1, 1)$ . Then the number of options is  $\binom{4}{i} \cdot \binom{10}{4-i} \cdot 10 \cdot 4! \cdot 10! \cdot 2^{5+i}$ . Choose one element of  $wt(Y) \neq 4$  as  $\pi(1, 1, 1, 1)$ . This can be done in 10 ways. The 4 elements where  $wt(Y) = 1$  can be permuted in  $4!$  ways. The 4 elements where  $wt(Y) = 2, 3$  can be permuted in  $10!$  ways. The function  $g(Y)$  is fixed when  $Y$  is  $(0, 0, 0, 0)$  (1 place,  $g(Y) = 0$ ) or  $wt(Y) = 1$  (4 places,  $g(Y) = 0$ ) or  $wt(\pi(Y)) = 1$  ( $4 - i$  places,  $g(Y) = 0$ ) or  $wt(Y) = 4$  (1 place,  $g(Y) = 1$  if  $wt(\pi(Y)) = 0$ , else  $g(Y) = 1$ ) or  $wt(\pi(Y)) = 4$  (1 place,  $g(Y) = (wt(Y) + 1) \bmod 2$ ). Thus  $g(Y)$  is fixed in  $11 - i$  places and we can put any choice from  $\{0, 1\}$  for  $16 - (11 - i) = 5 + i$  places.

So the total number of options is  $6 \sum_{i=0}^4 \binom{4}{i} \cdot \binom{10}{4-i} \cdot 4! \cdot 10! \cdot 2^{6+i} = 6 \cdot 4! \cdot 10! \cdot 2^6 \sum_{i=0}^4 \binom{4}{i} \cdot \binom{10}{4-i} \cdot 2^i \approx 2^{46.297492}$ . ■

**Remark 1** *Following Theorem 3, it is clear that for the function  $f$  as discussed in Theorem 4,  $\Delta_f \leq 40$ . Now we present the following specific case.*

*Consider  $\pi(Y) = Y$  for all  $Y \in \{0, 1\}^4$ ,  $g(Y) = 0$  for all  $Y \in \{0, 1\}^4 \setminus \{(1, 1, 1, 1)\}$  and  $g(Y) = 1$  for  $Y = (1, 1, 1, 1)$ . Let  $b(x) = b(X, Y) = X \cdot \pi(Y) + g(Y)$  and  $f(x)$  is as given in Construction 1. Then  $f$  is an  $(8, 1, 6, 116)$  function with  $\Delta_f = 24$ .*

*Note that we get an  $(8, 1, 6, 116)$  function  $f$  with  $\Delta_f = 24$  in this method which has earlier been found by simulated annealing and linear transformation in [5].*

## 4 The 10-variable 1-resilient Functions

We here start with 10-variable bent functions. Theorem 1 and Theorem 2 do not directly provide the idea how the exact construction of a 1-resilient function from a bent function is possible. Let us now describe a method where we will be able to identify a subclass of 10-variable Maiorana-McFarland type bent functions for this purpose.

As described in Section 2, we need to modify at least  $k = 22$  points (see Corollary 1). Now following Theorem 1 and Theorem 2, it is clear that we first need to select  $\frac{k}{2} + 2^{\frac{k}{2}-2} = 19$  distinct points. Note that we can have 1 point of weight 0 and 10 points of weight 1. Thus we need to find out 8 more points from weight 2. Once these 19 points are selected, further there are 3 more points to be chosen.

Now we refer to the  $S \oplus b(S)$  matrix given here. We present the first 19 points and after the horizontal line we show the next 3 points. Note that the choice of the all zero point and the points of weight 1 are clear from the discussion in Theorem 1. However, it is still to be sorted out how exactly the 8 points of weight 2 are chosen. We here do that by observation and choose the 8 points of weight 2 out of total  $\binom{10}{2} = 45$  weight 2 points. The rest 3 points (one of weight 0 and other two of weight 2) are chosen properly to satisfy that weight of each column should be  $\frac{k}{2} - 2^{\frac{k}{2}-2} = 3$ . Now we need a bent function  $b$  on 10 variables with the property that  $b(x) = 0$  when  $x$  is any of the first 19 points and  $b(x) = 1$  when  $x$  is complement of any of the last 3 points. This means that the last three rows need to be complemented when they will be considered as input points in the function.

$$\mathbf{S} \oplus b(\mathbf{S}) = \begin{pmatrix}
 x_{10} & x_9 & x_8 & x_7 & x_6 & x_5 & x_4 & x_3 & x_2 & x_1 \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\
 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\
 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
 \hline
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\
 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0
 \end{pmatrix}$$

Thus, we construct two sets  $S_1$  and  $S_2$  as follows and then denote  $S =$

$$S_1 \cup S_2. \quad S_1 = \{
 (0,0,0,0,0,0,0,0,0,0), (0,0,0,0,0,0,0,0,0,1), (0,0,0,0,0,0,0,0,1,0), \\
 (0,0,0,0,0,0,0,1,0,0), (0,0,0,0,0,0,1,0,0,0), (0,0,0,0,1,0,0,0,0,0), \\
 (0,0,0,0,1,0,0,0,0,0), (0,0,1,0,0,0,0,0,0,0), \\
 (0,1,0,0,0,0,0,0,0,0), (1,0,0,0,0,0,0,0,0,0), (0,0,0,0,0,0,0,1,1,0), \\
 (0,0,0,0,0,1,1,0,0,0), (0,0,0,0,0,1,0,0,0,1), (0,0,0,1,1,0,0,0,0,0), \\
 (0,0,1,1,0,0,0,0,0,0), (0,1,1,0,0,0,0,0,0,0), (1,1,0,0,0,0,0,0,0,0), \\
 (1,0,0,0,1,0,0,0,0,0)\} \text{ and } S_2 =$$

$$\{(1,1,1,1,1,1,1,1,1,1), (1,1,1,1,1,1,1,1,0,0), (1,1,1,1,1,1,0,0,1,1)\}.$$

$$\text{Also consider } S'_1 = \{
 (0,0,0,0,0,0,0,0,0,0), (0,0,0,0,0,0,0,0,0,1), (0,0,0,0,0,0,0,0,1,0), \\
 (0,0,0,0,0,0,0,1,0,0), (0,0,0,0,0,0,1,0,0,0), (0,0,0,0,1,0,0,0,0,0), \\
 (0,0,0,0,1,0,0,0,0,0), (0,0,0,1,0,0,0,0,0,0), (0,0,1,0,0,0,0,0,0,0), \\
 (0,1,0,0,0,0,0,0,0,0), (1,0,0,0,0,0,0,0,0,0), (0,0,0,0,0,0,0,1,1,0), \\
 (0,0,0,0,0,1,1,0,0,0), (0,0,0,0,0,1,0,0,0,1)\},$$

$$S_3 = \{(0,0,0,0,0,0,0,1,0,1), (0,0,0,0,0,0,0,1,1,1), (0,0,0,0,0,0,1,0,0,1)\}.$$

$(0, 0, 0, 0, 0, 0, 1, 0, 1, 0), (0, 0, 0, 0, 0, 0, 1, 1, 1, 0), (0, 0, 0, 0, 0, 1, 0, 0, 1, 1),$   
 $(0, 0, 0, 0, 1, 1, 1, 0, 0, 1), (0, 0, 0, 0, 0, 1, 1, 1, 0, 0), (0, 0, 0, 0, 0, 1, 1, 1, 1, 1)\}$   
 and  $S_4 =$   
 $\{(0, 0, 1, 1, 1, 0, 0, 0, 0, 0), (0, 1, 1, 1, 0, 0, 0, 0, 0, 0), (1, 0, 0, 1, 1, 0, 0, 0, 0, 0),$   
 $(0, 0, 0, 0, 0, 1, 1, 0, 0, 0), (1, 1, 0, 0, 1, 0, 0, 0, 0, 0), (1, 1, 1, 0, 0, 0, 0, 0, 0, 0),$   
 $(1, 1, 1, 1, 0, 0, 0, 0, 0, 0)\}$ . We will talk about these sets  $S'_1, S_3$  and  $S_4$  little later. Let us now write the exact construction.

**Construction 2** We need a 10-variable bent function  $b(x)$  with the following properties:

1.  $b(x) = 0$  when  $x \in S_1$  and  $b(x) = 1$  when  $x \in S_2$ ,
2.  $W_b(\omega) = +32$  when  $\omega \in S'_1 \cup S_3 \cup S_4$ .

The function  $f(x)$  is as follows.

$$\begin{aligned}
 f(x) &= 1 \oplus b(x), & \text{if } x \in S \\
 &= b(x), & \text{otherwise.}
 \end{aligned}$$

From Theorem 1, it is clear that the function  $f(x)$  is 1-resilient. Now we need to calculate the nonlinearity of  $f$ . In fact, we will prove that  $nl(f) = 488$ , the currently best known nonlinearity for 10-variable 1-resilient functions. By Proposition 1,  $W_f(\omega) = W_b(\omega) - 2W_b(\omega)|_S$ . Thus, it is important to analyse the values of  $W_b(\omega)|_S$  for all  $\omega \in \{0, 1\}^{10}$ . However, this can not be done in a nice way as it has been done in the 8-variable case in Theorem 4. So we use a computer program to calculate  $W_b(\omega)|_S$  for all  $\omega \in \{0, 1\}^{10}$ . Note that when  $|W_b(\omega)|_S| \leq 8$ , then at those points  $|W_f(\omega)| \leq 48$ . Thus, we have no restriction on the Walsh spectra of the bent function  $b$  at these points to get the nonlinearity 488 for  $f$ . However, we need to concentrate on the cases when  $|W_b(\omega)|_S| \geq 12$ . We have checked that this happens when  $\omega \in S'_1 \cup S_3 \cup S_4$  and all these values are either +12 or +16. Thus as given in Construction 2, the Walsh spectra of the function  $b$  should be +32 at these points. Hence Construction 2 provides 10-variable 1-resilient functions having nonlinearity 488. Using similar technique as in Theorem 6, it is possible to get the count of such functions.

Note that we have not yet discussed the algebraic degree and autocorrelation properties of the functions. We now consider a specific case and check the algebraic degree and autocorrelation property. Take  $x = (x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10})$ ,  $X = (X_1, X_2, X_3, X_4, X_5)$ , and  $Y = (Y_1, Y_2, Y_3, Y_4, Y_5)$ . Further we identify  $X_1 = x_1, X_2 = x_2, X_3 = x_3, X_4 = x_4, X_5 = x_5, Y_1 = x_6, Y_2 = x_7, Y_3 = x_8, Y_4 = x_9, Y_5 = x_{10}$ .

Consider a 10-variable Maiorana-McFarland type bent function

$$b(x) = b(X, Y) = X \cdot \pi(Y) + g(Y),$$

where  $\pi$  is a permutation on  $\{0, 1\}^5$  with  $\pi(Y) = Y$  and  $g$  is a Boolean function on 5 variables which is a constant 0 function. It can be checked that this bent function satisfies the conditions required in Construction 2. Then we prepare  $f$  as given in Construction 2. We checked that nonlinearity of  $f$  is 488, algebraic degree is 8 and  $\Delta_f = 48$ . Now it is important to note the following two points.

1. The construction in [13, Theorem 4] required 26 points to be modified to get 1-resilient function from a bent function. We here need only 22 points to modify. Further, we have checked that the  $\Delta_f$  value of the function constructed in [13] is 64. The function we construct here has  $\Delta_f = 48$  and this is the best known value which is achieved for the first time here.
2. The  $(10, 1, 8, 488)$  function was first constructed in [10] and we have checked that  $\Delta_f$  value is 320 for that function. Thus our construction provides better parameter.

## 5 Improving the bound: Functions on 12 and 14 variables

From Corollary 1, we find that  $dBR_{12}(1) \geq 42$ . However, it seems that it is not possible to construct an 1-resilient function by toggling 42 bits of a bent function. Instead we succeeded to construct a  $(12, 1, 10, 2000)$  function  $f$ , with  $\Delta_f = 120$  by toggling 44 points of a bent function. Later in this section we will present arguments why at least 44 points are needed and that eventually provides direction towards better lower bound.

Thus taking  $k = 44$ , we have to first find  $\frac{k}{2} + 2^{\frac{k}{2}-2} = 38$  distinct points. We select the all zero input point and the twelve input points each of weight one. Now there are  $\binom{12}{2} = 66$  input points of weight two. Out of them we choose  $38 - 13 = 25$  points by trial and error. These points are 2560, 2304, 2176, 2112, 1280, 1152, 1088, 640, 576, 320, 1536, 384, 40, 36, 34, 33, 20, 18, 17, 10, 9, 5, 24, 6, 2080 when written as decimal integers corresponding to 12-bit binary numbers. We need a bent function such that it will have output zero at these 38 input points. Next we take the six input points 4095, 3055, 3575, 3835, 3965, 4030. We need a bent function which provides output one at these six points. Now we present the bent function.

Take  $x = (x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}, x_{11}, x_{12})$ ,  $X = (X_1, X_2, X_3, X_4, X_5, X_6)$ , and  $Y = (Y_1, Y_2, Y_3, Y_4, Y_5, Y_6)$ . Further we identify  $X_1 = x_1, X_2 = x_2, X_3 = x_3, X_4 = x_4, X_5 = x_5, X_6 = x_6, Y_1 = x_7, Y_2 = x_8, Y_3 = x_9, Y_4 = x_{10}, Y_5 = x_{11}, Y_6 = x_{12}$ . Consider a 12-variable Maiorana-McFarland type bent function  $b(x) = b(X, Y) = X \cdot \pi(Y) + g(Y)$

where  $\pi$  is a permutation on  $\{0,1\}^6$  with  $\pi(Y) = Y$ , except the cases  $\pi(1,1,1,1,1,0) = (1,1,1,1,1,1)$ ,  $\pi(1,1,1,1,1,1) = (1,1,1,1,1,0)$ . Here  $g$  is a Boolean function on 6 variables which is a constant 0 function.

The construction presented in [13] requires 54 points to be toggled and they could achieve a nonlinearity 1996. Thus our construction is clearly better. Further we get  $\Delta_f = 120$  for the  $(12,1,10,2000)$  function that we construct here. This is the best known autocorrelation parameter which was not known earlier.

### 5.1 Improving the bound

We like to refer the proof of Theorem 1. We have found the inequality

$$\sum_{i=1}^r i \binom{n}{i} + (r+1) \left( \frac{k}{2} + 2^{\frac{n}{2}-2} - \sum_{i=0}^r \binom{n}{i} \right) \leq n \times \left( \frac{k}{2} - 2^{\frac{n}{2}-2} \right), \quad (1)$$

that has been instrumental in getting the lower bound

$$k \geq 2^{\frac{n}{2}-1} + 2 \left[ \frac{(r+1)(2^{\frac{n}{2}-1} - \sum_{i=0}^r \binom{n}{i}) + \sum_{i=1}^r i \binom{n}{i}}{n-r-1} \right], \quad (2)$$

where  $r$  is the integer such that  $\sum_{i=0}^r \binom{n}{i} \leq 2^{\frac{n}{2}-1} + 1 < \sum_{i=0}^{r+1} \binom{n}{i}$  is satisfied. Note that the inequality 1 can be updated as

$$\begin{aligned} & \sum_{i=1}^r i \binom{n}{i} + (r+1) \left( \frac{k}{2} + 2^{\frac{n}{2}-2} - \sum_{i=0}^r \binom{n}{i} \right) + \\ & \sum_{i=1}^{r_1} i \binom{n}{i} + (r_1+1) \left( \frac{k}{2} - 2^{\frac{n}{2}-2} - \sum_{i=0}^{r_1} \binom{n}{i} \right) \leq n \times \left( \frac{k}{2} - 2^{\frac{n}{2}-2} \right), \quad (3) \end{aligned}$$

where  $r_1$  is such that,  $\sum_{i=0}^{r_1} \binom{n}{i} \leq \frac{k}{2} - 2^{\frac{n}{2}-2} < \sum_{i=0}^{r_1+1} \binom{n}{i}$ .

The reason for this updation is as follows. In the proof of Theorem 1, we noted that since all the rows of  $\mathbf{S}$  are distinct and further  $b(\mathbf{S})$  contains  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  many 0's,  $\mathbf{S} \oplus b(\mathbf{S})$  should contain at least  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  distinct rows. Keeping these  $\frac{k}{2} + 2^{\frac{n}{2}-2}$  distinct rows apart, as  $b(\mathbf{S})$  contains  $\frac{k}{2} - 2^{\frac{n}{2}-2}$  many 1's, in  $\mathbf{S} \oplus b(\mathbf{S})$  the rest  $\frac{k}{2} - 2^{\frac{n}{2}-2}$  rows are also distinct among themselves. This gives the updated inequality 3.

Note that, once we calculate  $k$  using inequality 2, we can check whether this value of  $k$  satisfies inequality 3. If the inequality 3 is satisfied, then we may live with that value of  $k$ , but if it is not, then it is clearly not possible to get a matrix  $\mathbf{S} \oplus b(\mathbf{S})$  as required and one needs to increase  $k$  by at least 2 as  $k$  is always even.

We first revisit the cases  $n = 8, 10$ . For  $n = 8$ , from inequality 2 we get  $k = 10$ . Thus,  $r = 1, r_1 = 0$ . This satisfies  $(8 + 2(5 + 4 - 9)) +$

$(0 + 1(5 - 4 - 1)) \leq 8 \times (1)$ , i.e., this satisfies inequality 3. For  $n = 10$ , from inequality 2 we get  $k = 22$ . Thus,  $r = 1, r_1 = 0$ . This satisfies  $(10 + 2(11 + 8 - 11)) + (0 + 1(11 - 8 - 1)) \leq 10 \times (3)$ , i.e., this satisfies inequality 3.

Now we look into the case for  $n = 12$ . From inequality 2 we get  $k = 42$ . Thus,  $r = 1, r_1 = 0$ . This does not satisfy  $(12 + 2(21 + 16 - 13)) + (0 + 1(21 - 16 - 1)) \leq 12 \times (5)$ , i.e., inequality 3 is not satisfied. Hence we need to increase  $k$  by 2, which gives  $k = 44$ . This satisfies  $(12 + 2(22 + 16 - 13)) + (0 + 1(22 - 16 - 1)) \leq 12 \times (6)$ , i.e., this satisfies inequality 3.

## 5.2 The 14-variable case

From Corollary 1, we find that  $dBR_{14}(1) \geq 84$ , which is also given by inequality 2. Putting  $k = 84$ , we get  $r = 1, r_1 = 0$ . We find that this does not satisfy  $141 = (14 + 2(42 + 32 - 15)) + (0 + 1(42 - 32 - 1)) \leq 14 \times (10) = 140$ , i.e., inequality 3 is not satisfied. Hence we need to increase  $k$  from 84 to 86. This satisfies  $144 = (14 + 2(43 + 32 - 15)) + (0 + 1(43 - 32 - 1)) \leq 14 \times (11) = 154$ , i.e., this satisfies inequality 3.

Now we construct the matrix. Thus taking  $k = 86$ , we have to first find  $\frac{k}{2} + 2^{\frac{k}{2}-2} = 75$  distinct points. We select the all zero input point and the fourteen input points each of weight one. Now there are  $\binom{14}{2} = 91$  input points of weight two. Out of them we choose  $75 - 15 = 60$  points by trial and error. These points are 384, 640, 768, 1152, 1280, 1536, 2176, 2304, 2560, 3072, 4224, 4352, 4608, 5120, 6144, 8320, 8448, 8704, 9216, 10240, 12288, 3, 5, 6, 9, 10, 12, 17, 18, 20, 24, 33, 34, 36, 40, 48, 65, 66, 68, 72, 80, 96, 8256, 4128, 2064, 1032, 516, 258, 129, 4160, 2080, 1040, 520, 260, 130, 8193, 1088, 544, 272, 136, when written as decimal integers corresponding to 14-bit binary numbers. We need a bent function such that it will have output zero at these 75 input points. Next we take the 11 input points 16383, 14847, 15999, 16287, 16359, 4095, 6143, 10239, 16377, 16378, 16380. We need a bent function which provides output one at these eleven points. Now we present the bent function.

Take  $x = (x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}, x_{11}, x_{12}, x_{13}, x_{14})$ ,  $X = (X_1, X_2, X_3, X_4, X_5, X_6, X_7)$ , and  $Y = (Y_1, Y_2, Y_3, Y_4, Y_5, Y_6, Y_7)$ . Further we identify  $X_1 = x_1, X_2 = x_2, X_3 = x_3, X_4 = x_4, X_5 = x_5, X_6 = x_6, X_7 = x_7, Y_1 = x_8, Y_2 = x_9, Y_3 = x_{10}, Y_4 = x_{11}, Y_5 = x_{12}, Y_6 = x_{13}, Y_7 = x_{14}$ . Consider a 14-variable Maiorana-McFarland type bent function  $b(x) = b(X, Y) = X \cdot \pi(Y) + g(Y)$  where  $\pi$  is a permutation on  $\{0, 1\}^7$  with  $\pi(Y) = Y$ , except the cases

$$\begin{aligned} \pi(0, 0, 0, 0, 0, 0, 1) &= (0, 0, 0, 0, 1, 0, 0), & \pi(0, 0, 0, 0, 1, 0, 0) &= (0, 0, 0, 0, 0, 0, 1), \\ \pi(0, 0, 0, 0, 0, 1, 0) &= (0, 0, 0, 1, 0, 0, 0), & \pi(0, 0, 0, 1, 0, 0, 0) &= (0, 0, 0, 0, 0, 1, 0), \\ \pi(0, 0, 0, 1, 0, 0, 1) &= (0, 0, 1, 0, 0, 0, 0), & \pi(0, 0, 1, 0, 0, 0, 0) &= (0, 0, 0, 1, 0, 0, 1), \\ \pi(0, 0, 0, 0, 1, 0, 1) &= (0, 1, 0, 0, 0, 0, 0), & \pi(0, 1, 0, 0, 0, 0, 0) &= (0, 0, 0, 0, 1, 0, 1), \end{aligned}$$

$\pi(0, 0, 0, 0, 1, 1, 0) = (1, 0, 0, 0, 0, 0, 0)$ ,  $\pi(1, 0, 0, 0, 0, 0, 0) = (0, 0, 0, 0, 1, 1, 0)$ . Here  $g$  is a Boolean function on 7 variables which is a constant 0 function.

This function  $f$  is a (14, 1, 12, 8092) function with  $\Delta_f = 168$ . Note that in [12], a (14, 1, 12, 8104) function  $g$  has been reported with  $\Delta_g = 800$ . Thus the function constructed in this method is little worse in nonlinearity, but much better in terms of autocorrelation spectra.

## 6 Conclusion

In this paper we present lower bound on the minimum distance  $dBR_n(1)$  between bent and 1-resilient functions on  $n$  variables, where  $n$  is even. We have also shown that it is possible to get 1-resilient functions by modifying exactly  $dBR_n(1)$  many bits for  $n = 4, 6, 8, 10$  which shows that the minimum distance is tight in these cases. This bound is not tight for  $n \geq 12$  and we improve the basic bound to get a tighter bound. The case for  $n = 8$  could be nicely handled, but it starts to become complicated from  $n = 10$  and requires some computer simulation for exact construction of the 1-resilient functions.

A lot of open questions are still to be solved. First of all, a relatively hard question is to find out the minimum distance between bent and  $m$ -resilient functions on  $n$  variables, which we may denote as  $dBR_n(m)$ . It seems natural that  $dBR_n(n-2) > dBR_n(n-3) > \dots > dBR_n(1)$ , though it needs a proof. Note that  $(n-2)$ -resilient functions on  $n$  variables are basically the affine functions, which are known to be at maximum distance from the bent function [16].

The functions we provide here up to 12 variables possess currently best known parameters. The upper bound on nonlinearity of 1-resilient functions is  $2^{n-1} - 2^{\frac{n}{2}-1} - 4$  for  $n$  even as described in [18]. The tightness of this bound [18] has been shown upto  $n = 8$ . For  $n \geq 10$ , there is no evidence of an 1-resilient function attaining that bound [18]. Our construction modifies  $dBR_n(1) > 2^{\frac{n}{2}-1}$  many bits and it seems unlikely that modifying these many bits will result in a fall of nonlinearity only 4 for  $n \geq 10$ .

It seems that the improved bound presented in Subsection 5.1 can be a candidate to be the exact minimum distance. Proving that this improved bound is an exact one, the following steps need to be proved.

1. Given the value of  $k$  one can always construct a  $k \times n$  binary matrix  $S \oplus b(S)$  with the required properties.
2. Given the matrix, one can always construct a bent function on  $n$  variables which possesses proper outputs 0 or 1 at the corresponding input points.

Solving this problem will be an interesting research contribution.

**Acknowledgment :** The authors like to acknowledge Dr. Kishan Chand Gupta of Indian Statistical Institute, Kolkata, for detailed discussion on the proof of Theorem 1.

## References

- [1] C. Carlet. On the coset weight divisibility and nonlinearity of resilient and correlation immune functions. In *Sequences and their Applications - SETA 2001*, Discrete Mathematics and Theoretical Computer Science, pages 131-144, Bergen, Norvge, May 2001. Springer-Verlag.
- [2] C. Carlet. A larger Class of Cryptographic Boolean Functions via a Study of the Maiorana-McFarland Constructions. In *Advances in Cryptology - CRYPTO 2002*, number 2442 in Lecture Notes in Computer Science, pages 549–564. Springer Verlag, 2002.
- [3] C. Carlet and P. Sarkar. Spectral domain analysis of correlation immune and resilient Boolean functions. *Finite Fields and Its Applications*, 8(1):120–130, January 2002.
- [4] P. Charpin and E. Pasalic. On Propagation Characteristics of Resilient Functions. In *SAC 2002*, number 2595 in Lecture Notes in Computer Science, pages 175–195. Springer-Verlag, 2003.
- [5] J. Clark, J. Jacob, S. Stepney, S. Maitra and W. Millan. Evolving Boolean Functions Satisfying Multiple Criteria. In *INDOCRYPT 2002*, number 2551 in Lecture Notes in Computer Science, pages 246–259, Springer Verlag, 2002.
- [6] J. F. Dillon. Elementary Hadamard Difference sets. PhD Thesis, University of Maryland, 1974.
- [7] H. Dobbertin. Construction of bent functions and balanced Boolean functions with high nonlinearity. In *Fast Software Encryption - FSE 1994*, number 1008 in Lecture Notes in Computer Science, pages 61–74. Springer-Verlag, 1994.
- [8] M. Fedorova and Y. V. Tarannikov. On the constructing of highly nonlinear resilient Boolean functions by means of special matrices. In *Progress in Cryptology - INDOCRYPT 2001*, number 2247 in Lecture Notes in Computer Science, pages 254–266. Springer Verlag, 2001.

- [9] X. Guo-Zhen and J. Massey. A spectral characterization of correlation immune combining functions. *IEEE Transactions on Information Theory*, 34(3):569–571, May 1988.
- [10] S. Maitra and E. Pasalic. Further constructions of resilient Boolean functions with very high nonlinearity. *IEEE Transactions on Information Theory*, 48(7):1825–1834, July 2002.
- [11] S. Maitra. Autocorrelation Properties of correlation immune Boolean functions. INDOCRYPT 2001, number 2247 Lecture Notes in Computer Science. Pages 242–253. Springer Verlag, December 2001.
- [12] S. Maitra and E. Pasalic. A Maiorana-McFarland type construction for resilient Boolean functions on  $n$  variables ( $n$  even) with nonlinearity  $> 2^{n-1} - 2^{\frac{n}{2}} + 2^{\frac{n}{2}-2}$ . *Discrete Applied Mathematics* 154(2): 357-369, 2006.
- [13] S. Maity and T. Johansson. Construction of Cryptographically Important Boolean Functions. In *Progress in Cryptology - INDOCRYPT 2002*, number 2551 in Lecture Notes in Computer Science, pages 234–245. Springer Verlag, 2002.
- [14] S. Maity and S. Maitra. Minimum distance between bent and 1-resilient functions. In *Fast Software Encryption-FSE 2004*, number 3017 in Lecture Notes in Computer Science, pages 143-160 . Springer-Verlag, 2004.
- [15] B. Preneel, W. Van Leekwijck, L. Van Linden, R. Govaerts, and J. Vandewalle. Propagation characteristics of Boolean functions. In *Advances in Cryptology - EUROCRYPT'90*, number 473 in Lecture Notes in Computer Science, pages 161–173. Springer-Verlag, 1991.
- [16] O. S. Rothaus. On bent functions. *Journal of Combinatorial Theory, Series A*, 20:300–305, 1976.
- [17] P. Sarkar and S. Maitra. Construction of nonlinear Boolean functions with important cryptographic properties. In *Advances in Cryptology - EUROCRYPT 2000*, number 1807 in Lecture Notes in Computer Science, pages 485–506. Springer Verlag, 2000.
- [18] P. Sarkar and S. Maitra. Nonlinearity bounds and constructions of resilient Boolean functions. In *Advances in Cryptology - CRYPTO 2000*, number 1880 in Lecture Notes in Computer Science, pages 515–532. Springer Verlag, 2000.

- [19] J. Seberry, X. M. Zhang, and Y. Zheng. Nonlinearly balanced Boolean functions and their propagation characteristics. In *Advances in Cryptology - CRYPTO'93*, number 773 in Lecture Notes in Computer Science, pages 49–60. Springer-Verlag, 1994.
- [20] Y. V. Tarannikov. On resilient Boolean functions with maximum possible nonlinearity. In *Progress in Cryptology - INDOCRYPT 2000*, number 1977 in Lecture Notes in Computer Science, pages 19–30. Springer Verlag, 2000.
- [21] Y. V. Tarannikov. New constructions of resilient Boolean functions with maximal nonlinearity. In *Fast Software Encryption - FSE 2001*, number 2355 in Lecture Notes in Computer Science, pages 66–77. Springer Verlag, 2001.
- [22] Y. V. Tarannikov, P. Korolev and A. Botev. Autocorrelation coefficients and correlation immunity of Boolean functions. In *ASIACRYPT 2001*, number 2248 in Lecture Notes in Computer Science, pages 460–479. Springer Verlag, 2001.
- [23] Y. Zheng and X. M. Zhang. Improved upper bound on the nonlinearity of high order correlation immune functions. In *Selected Areas in Cryptography - SAC 2000*, number 2012 in Lecture Notes in Computer Science, pages 264–274. Springer Verlag, 2000.
- [24] X. M. Zhang and Y. Zheng. GAC - the criterion for global avalanche characteristics of cryptographic functions. *Journal of Universal Computer Science*, 1(5):316–333, 1995.
- [25] Y. Zheng and X. M. Zhang. On relationships among propagation degree, nonlinearity and correlation immunity. In *Advances in Cryptology - ASIACRYPT 2000*, number 1976 in Lecture Notes in Computer Science, pages 470–482. Springer Verlag, 2000.