

# Antipodal Labelings for Cycles

Justie Su-Tzu Juan\* and Daphne Der-Fen Liu†

Submitted: December 2006; revised: August 2007

## Abstract

Let  $G$  be a graph with diameter  $d$ . An antipodal labeling of  $G$  is a function  $f$  that assigns to each vertex a non-negative integer (label) such that for any two vertices  $u$  and  $v$ ,  $|f(u) - f(v)| \geq d - d(u, v)$ , where  $d(u, v)$  is the distance between  $u$  and  $v$ . The span of an antipodal labeling  $f$  is  $\max\{f(u) - f(v) : u, v \in V(G)\}$ . The antipodal number for  $G$ , denoted by  $\text{an}(G)$ , is the minimum span of an antipodal labeling for  $G$ . Let  $C_n$  denote the cycle on  $n$  vertices. Chartrand et al. [4] determined the value of  $\text{an}(C_n)$  for  $n \equiv 2 \pmod{4}$ . In this article we obtain the value of  $\text{an}(C_n)$  for  $n \equiv 1 \pmod{4}$ , confirming a conjecture in [4]. Moreover, we settle the case  $n \equiv 3 \pmod{4}$ , and improve the known lower bound and give an upper bound for the case  $n \equiv 0 \pmod{4}$ .

---

\*Department of Computer Science and Information Engineering, National Chi Nan University, Nantou 54561, Taiwan. Email: jsjuan@ncnu.edu.tw. Supported in part by the National Science Council under grant NSC95-2918-I-260-002.

†Corresponding author. Department of Mathematics, California State University, Los Angeles, CA 90032. Email: dliu@calstatela.edu. Supported by the National Science Foundation, USA, under grant DMS 0302456.

# 1 Introduction

Radio  $k$ -labeling was motivated by the frequency assignment problem (cf. [7]). Let  $k$  be a positive integer. A *radio  $k$ -labeling* (or  *$k$ -labeling* for short) for a graph  $G$  is a function,  $f : V(G) \rightarrow \{0, 1, 2, \dots\}$ , such that the following is satisfied for any vertices  $u$  and  $v$ :

$$|f(u) - f(v)| \geq k + 1 - d(u, v).$$

where  $d(u, v)$  denotes the distance between  $u$  and  $v$ . The *span* of such a function  $f$ , denoted by  $\text{sp}(f)$ , is defined as  $\text{sp}(f) = \max\{f(u) - f(v) : u, v \in V(G)\}$ . The minimum span over all  $k$ -labelings of a graph  $G$  is called the  $\Phi_k$ -number and denoted by  $\Phi_k(G)$ .

For the special case that  $k = 1$ , the 1-labeling is indeed the conventional vertex coloring and we have  $\Phi_1(G) = \chi(G) - 1$ , where  $\chi(G)$  is the chromatic number of  $G$ . Another special case is when  $k = 2$ , the 2-labeling is the same as the *distance two labeling* (or  *$L(2, 1)$ -labeling*) which has been studied extensively in the past years (cf. [1, 2, 3, 9, 10, 11, 12, 14]). The  $\Phi_2$ -number is known as the  $\lambda$ -number of  $G$ .

The radio  $k$ -labeling for large values of  $k$  has also been investigated by several authors. Let  $G$  be a connected graph. The maximum distance among all pairs of vertices in  $G$  is the *diameter* of  $G$ , denoted by  $\text{diam}(G)$ . The *radio labeling* (or *multi-level distance labeling*) is a radio  $k$ -labeling when  $k = \text{diam}(G)$ . The  $\Phi_{\text{diam}(G)}$ -number of  $G$  is called the *radio number* of  $G$ , denoted by  $\text{rn}(G)$ . The radio number for different families of graphs has been investigated in [6, 8, 15, 16, 17, 18, 19]. For instance, the radio number for paths and cycles has been studied in [6, 8, 19] and was recently settled in [18].

When  $k = \text{diam}(G) - 1$ , a  $k$ -labeling is called an *antipodal labeling*. That is, an *antipodal labeling* (or *radio antipodal coloring*) for  $G$  is a function,  $f : V(G) \rightarrow \{0, 1, 2, \dots\}$ , such that the following is satisfied for any two vertices  $u$  and  $v$ :

$$|f(u) - f(v)| \geq \text{diam}(G) - d(u, v).$$

The *antipodal number* for  $G$ , denoted by  $\text{an}(G)$ , is the minimum span of an antipodal labeling admitted by  $G$ . Notice that a radio labeling is a one-to-one function, while in an antipodal labeling, two vertices of distance  $\text{diam}(G)$  apart may receive the same label (this is where the name “antipodal” came from).

The antipodal labeling for graphs was first studied by Chartrand et al. [4, 5], in which, among other results, general bounds of  $\text{an}(G)$  were obtained. Khennoufa and Togni [13] determined the exact value of  $\text{an}(P_n)$  for paths  $P_n$ . The antipodal labeling for cycles  $C_n$  was studied in [4], in which lower bounds for  $\text{an}(C_n)$  were shown. In addition, the bound for the case  $n \equiv 2 \pmod{4}$  was proved to be the exact value of  $\text{an}(C_n)$ , and the bound for the case  $n \equiv 1 \pmod{4}$  was conjectured to be the exact value as well [4].

In this article, we confirm the conjecture mentioned above. Moreover, we determine the value of  $\text{an}(C_n)$  for the case  $n \equiv 3 \pmod{4}$ . For the case  $n \equiv 0 \pmod{4}$ , we improve the known lower bound [4] and give an upper bound. It is conjectured that the upper bound is the exact value.

## 2 Lower Bounds

In this section, we establish lower bounds for  $\text{an}(C_n)$ . These bounds were proved by Chartrand et al [4]. We present here a different proof which includes techniques that will be used in later sections.

In an antipodal labeling, the number assigned to a vertex is called a *label*. Notice that as we are seeking for the minimum span of an antipodal labeling, without loss of generality we assume that the label 0 is used by any antipodal labeling. Consequently, the span of  $f$  is the maximum label used.

In the following we introduce notations to be used throughout this article. Denote  $V(C_n) = \{v_0, v_1, \dots, v_{n-1}\}$ ,  $v_i v_{i+1} \in E(C_n)$  for  $0 \leq i \leq n-2$ , and  $v_{n-1} v_0 \in E(C_n)$ . The diameter of  $C_n$  is denoted by  $d$ , where  $d = \lfloor n/2 \rfloor$ . Every antipodal labeling  $f$  for  $C_n$  gives an ordering (which may not be unique) of the vertices according to the labels assigned. Denote the ordering by  $(x_0, x_1, \dots, x_{n-1})$ , where  $\{x_0, x_1, \dots, x_{n-1}\} = V(C_n)$  and

$$0 = f(x_0) \leq f(x_1) \leq f(x_2) \leq \dots \leq f(x_{n-1}).$$

Note, the span of  $f$  is  $f(x_{n-1})$ .

For  $i = 0, 1, \dots, n-2$ , we define the *distance gap* and *label gap*, respectively, by:

$$d_i = d(x_i, x_{i+1}), \quad f_i = f(x_{i+1}) - f(x_i).$$

By definition, it holds that  $f_i \geq d - d_i$ .

**Proposition 1** For any three vertices  $u, v$  and  $w$  on a cycle  $C_n$ ,

$$d(u, v) + d(v, w) + d(u, w) \leq n.$$

**Proof.** Without loss of generality, assume  $d(u, v), d(v, w) \leq d(u, w)$ . If all the three vertices lie on one half of the cycle, then  $d(u, v) + d(v, w) + d(u, w) = 2d(u, w) \leq n$ . Otherwise, we have  $d(u, v) + d(v, w) + d(u, w) = n$ .  $\square$

**Lemma 2** Let  $f$  be an antipodal labeling for  $C_n$ ,  $n \geq 3$ , with labels  $f(x_0) \leq f(x_1) \leq \dots \leq f(x_{n-1})$ . Let  $n = 4k + r$  for some  $0 \leq r \leq 3$ . Then for any  $0 \leq i \leq n - 3$ ,

$$f(x_{i+2}) - f(x_i) = f_i + f_{i+1} \geq \begin{cases} k, & \text{if } r = 0, 1, 3; \\ k + 1, & \text{if } r = 2. \end{cases}$$

**Proof.** By definition, we have  $f(x_{i+1}) - f(x_i) \geq d - d(x_{i+1}, x_i)$ ,  $f(x_{i+2}) - f(x_{i+1}) \geq d - d(x_{i+2}, x_{i+1})$ , and  $f(x_{i+2}) - f(x_i) \geq d - d(x_{i+2}, x_i)$ . Summing up these three inequalities and by Proposition 1, we get

$$\begin{aligned} 2(f(x_{i+2}) - f(x_i)) &\geq 3d - (d(x_i, x_{i+1}) + d(x_{i+1}, x_{i+2}) \\ &\quad + d(x_i, x_{i+2})) \\ &\geq 3d - n. \end{aligned}$$

Therefore,  $f_i + f_{i+1} = f(x_{i+2}) - f(x_i) \geq \lceil (3d - n)/2 \rceil$ . The results then follow by immediate calculations for different values of  $n$ .  $\square$

**Corollary 3** [4] Let  $n = 4k + r$  for some  $n \geq 3$  and  $0 \leq r \leq 3$ . Then

$$\text{an}(C_n) \geq \begin{cases} k(2k - 1), & \text{if } r = 0; \\ 2k^2, & \text{if } r = 1; \\ 2k(k + 1), & \text{if } r = 2; \\ k(2k + 1), & \text{if } r = 3. \end{cases}$$

**Proof.** Let  $f$  be an antipodal labeling for  $C_n$ . The span of  $f$  is

$$f(x_{n-1}) = f_0 + f_1 + \cdots + f_{n-2}.$$

By Lemma 2, the results follow by pairing up the terms in the above summation and leaving the last term  $f_{n-2}$  (if  $n$  is even) which is at least 0.  $\square$

In [4], it was proved that the equality in Corollary 3 holds for the case  $n \equiv 2 \pmod{4}$ , and conjectured that the equality also holds for the case  $n \equiv 1 \pmod{4}$ . This conjecture is confirmed in the next section.

### 3 $n = 4k + 1$

Let  $f$  be an antipodal labeling for a cycle  $C_n$  with  $0 = f(x_0) \leq f(x_1) \leq \cdots \leq f(x_{n-1})$ . In the rest of this article, we denote the permutation  $\pi$  on  $\{0, 1, 2, \dots, n-1\}$  generated from  $f$  with

$$x_i = v_{\pi(i)}.$$

For an integer  $x$  and a positive integer  $y$ , we denote " $x \bmod y$ " as a binary operation which outputs an integer  $z$  with  $z \equiv x \pmod{y}$  and  $0 \leq z \leq y-1$ .

In this section, we prove the following result:

**Theorem 4** *If  $n = 4k + 1$  for some integer  $k \geq 1$ , then*

$$\text{an}(C_n) = 2k^2.$$

**Proof.** By Corollary 3, it suffices to find an antipodal labeling with span  $2k^2$ . Two cases are considered. Recall  $d = \text{diam}(C_{4k+1}) = 2k$ .

**Case 1.  $k$  is odd** First, we label the  $2k + 1$  vertices  $x_0, x_2, \dots, x_{4k}$  by

$$\pi(2i) = ki \bmod n, \text{ and } f(x_{2i}) = ki, \text{ for } i = 0, 1, 2, \dots, 2k.$$

For instance,  $\pi(2) = k$  (i.e.,  $x_2 = v_k$ ) and  $f(x_2) = k$ ; and  $\pi(4k) = 2k - \frac{k-1}{2}$  and  $f(x_{4k}) = 2k^2$ .

Secondly, we label the remaining vertices  $x_1, x_3, \dots, x_{4k-1}$  by  $\pi(1) = \pi(4k) + k = 3k - \frac{k-1}{2}$ ; and  $\pi(2i+1) = (\pi(2i-1) + k)$

mod  $n$ , for  $i = 1, 2, \dots, 2k - 1$ , with labels  $f(x_{2i+1}) = (k - 1)/2 + ki$  for  $i = 0, 1, \dots, 2k - 1$ . See Figure 1 for an example. In Figure 1 (and all other figures), the number inside the circle for each vertex is the label assigned to that vertex.




Figure 1: An antipodal labeling for  $C_{13}$  with minimum span  $\text{an}(C_{13}) = 18$ .

To see that  $\pi$  is a permutation of  $\{0, 1, \dots, n - 1\}$ , we observe that  $\pi(0), \pi(2), \dots, \pi(4k), \pi(1), \pi(3), \dots, \pi(4k - 1)$  is a list of vertices winding around  $C_n$  by jumping  $k$  vertices between any two consecutive terms. Since  $\text{gcd}(n, k) = 1$ , so  $\pi$  is a permutation of  $\{0, 1, \dots, n - 1\}$ . In addition, one can easily check that for every  $i$  the following hold:

$$\begin{aligned} f(x_{i+1}) - f(x_i) &\geq d - d(x_{i+1}, x_i), \\ f(x_{i+2}) - f(x_i) &= k = 2k - k = d - d(x_{i+2}, x_i), \\ f(x_{i+s}) - f(x_i) &\geq 2k \geq d - d(x_{i+s}, x_i), \text{ for } s \geq 4. \end{aligned}$$

Hence, to show that  $f$  is an antipodal labeling, it suffices to verify  $f(x_{i+3}) - f(x_i) \geq 2k - d(x_{i+3}, x_i)$ . This is true since  $d(x_{i+3}, x_i) = (k + 1)/2$ , and  $f(x_{i+3}) - f(x_i) \in \{(3k - 1)/2, (3k + 1)/2\}$ .

**Case 2.  $k$  is even** Similar to Case 1, we first label the  $2k + 1$  vertices  $x_0, x_2, \dots, x_{4k}$ , by  $\pi(2i) = ki \bmod n$ , for  $i = 0, 1, \dots, 2k$ ,

using labels  $f(x_{2i}) = ki$ . Note that since  $2k^2 \equiv n - \frac{k}{2} \pmod{n}$ , we have  $x_{4k} = v_{n-(k/2)}$ .

Secondly, we label the remaining vertices by  $\pi(1) = 2k+1$ ,  $f(x_1) = 0$ , and

$$\pi(2i+1) = \begin{cases} (\pi(2i-1) + k) \pmod{n}, & \text{if } i \text{ is odd;} \\ (\pi(2i-1) + k + 1) \pmod{n}, & \text{if } i \text{ is even,} \end{cases}$$

with labels

$$f(x_{2i+1}) = \begin{cases} f(x_{2i-1}) + k, & \text{if } i \text{ is odd;} \\ f(x_{2i-1}) + k + 1, & \text{if } i \text{ is even.} \end{cases}$$

See Figure 2 for an example.




Figure 2: An antipodal labeling for  $C_9$  with minimum span  $\text{an}(C_9) = 8$ .

By calculation,  $\pi(1) = 2k + 1 \equiv -2k \pmod{n}$ , and for  $1 \leq i \leq 2k - 1$ ,

$$\pi(2i+1) \equiv \begin{cases} -ik \pmod{n}, & \text{if } i \text{ is odd;} \\ -(i+2)k \pmod{n}, & \text{if } i \text{ is even.} \end{cases}$$

Since  $\pi(2i) = ki \pmod{n}$ , for  $0 \leq i \leq 2k$ , we conclude

$$\{\pi(i) : 0 \leq i \leq 4k\} = \{jk \pmod{n} : -2k \leq j \leq 2k\}.$$

Since  $\gcd(n, k) = 1$ ,  $\pi$  is a permutation. Similar to Case 1, it is straightforward to check that  $f$  is an antipodal labeling, and we shall leave the details to the reader.  $\square$

## 4 $n = 4k + 3$

As it turned out (Theorem 5), the exact value of  $\text{an}(C_{4k+3})$  is greater than the lower bound established in Corollary 3.

**Theorem 5** For every integer  $k \geq 0$ ,  $\text{an}(C_{4k+3}) = 2k^2 + 2k$ .

Note, when  $k = 0$  in Theorem 5, it is trivial that  $\text{an}(C_3) = 0$ . The following lemma will be used to prove Theorem 5 for  $k \geq 1$ .

**Lemma 6** Let  $f$  be an antipodal labeling for  $C_n$  where  $n = 4k + 3$ ,  $k \geq 1$ . If  $f_i + f_{i+1} = k$  for some  $0 \leq i \leq n - 3$ , then the following hold:

$$(1) \quad d(x_i, x_{i+2}) = k + 1,$$

$$(2) \quad f_i = t, \quad d_{i+1} = k + t + 1, \quad \text{and} \quad d_i = 2k - t + 1, \quad \text{for some} \\ t \in \{0, 1, \dots, k\}.$$

**Proof.** Recall  $d = \text{diam}(C_{4k+3}) = 2k + 1$ . Assume  $f_i + f_{i+1} = k$  for some  $i$ . By definition,

$$\begin{aligned} d(x_i, x_{i+2}) &\geq d - (f(x_{i+2}) - f(x_i)) = d - (f_{i+1} + f_i) \\ &= (2k + 1) - k = k + 1. \end{aligned}$$

On the other hand, by Proposition 1 and definition, we have

$$\begin{aligned} d(x_i, x_{i+2}) &\leq (4k + 3) - (d_i + d_{i+1}) \\ &\leq (4k + 3) - (d - f_i + d - f_{i+1}) \\ &= (4k + 3) - (4k + 2 - k) \\ &= k + 1. \end{aligned}$$

This verifies (1).

Let  $f_i = t$  for some  $t \in \{0, 1, \dots, k\}$ . By (1), the second equality in the above holds, which implies that  $d_i = d - f_i$  and  $d_{i+1} = d - f_{i+1}$ . Therefore, (2) follows as  $d = 2k + 1$ .  $\square$

**Lemma 7** Let  $f$  be an antipodal labeling for  $C_n$  where  $n = 4k + 3$  for some integer  $k \geq 1$ . Then for any  $0 \leq i \leq n - 5$ ,

$$f_i + f_{i+1} + f_{i+2} + f_{i+3} \geq 2k + 1.$$



**Proof.** Assume to the contrary that for some  $i$ ,  $f_i + f_{i+1} + f_{i+2} + f_{i+3} \leq 2k$ . By Lemma 2,  $f_i + f_{i+1} = f_{i+2} + f_{i+3} = k$ . By symmetry and by Lemma 6 (1), without loss of generality, assume  $x_i = v_0$ ,  $x_{i+2} = v_{k+1}$  and  $x_{i+4} = v_{2(k+1)}$ . By Lemma 6 (2),  $f_i = t$  for some  $0 \leq t \leq k$  and  $d_i = 2k - t + 1$ . Note,  $x_{i+1} \neq v_{2k-t+1}$ , for otherwise it would be  $d_{i+1} = d(x_{i+1}, x_{i+2}) = k - t$ , a contradiction. Hence, we conclude  $x_{i+1} = v_{n-(2k-t+1)} = v_{2k+t+2}$ . This implies  $d(x_{i+4}, x_i) = t$ . Because  $f$  is an antipodal labeling, we have

$$\begin{aligned} 2k - t = f_{i+1} + f_{i+2} + f_{i+3} &= f(x_{i+4}) - f(x_{i+1}) \\ &\geq 2k + 1 - d(x_{i+4}, x_{i+1}) \\ &= 2k + 1 - t, \end{aligned}$$

a contradiction. □

**Theorem 8** For every integer  $k \geq 1$ ,  $\text{an}(C_{4k+3}) \geq 2k^2 + 2k$ .

**Proof.** By Lemmas 2 and 7, the span of an antipodal labeling  $f$  for  $C_{4k+3}$  has

$$\begin{aligned} &f_0 + f_1 + \cdots + f_{4k+1} \\ &= \sum_{i=0}^{k-1} (f_{4i} + f_{4i+1} + f_{4i+2} + f_{4i+3}) + f_{4k} + f_{4k+1} \\ &\geq k(2k + 1) + k = 2k^2 + 2k. \end{aligned}$$

□

**Proof of Theorem 5.** For  $k = 0$ ,  $\text{an}(C_3) = 0$  is trivial as mentioned earlier. For  $k \geq 1$ , it remains to find an antipodal labeling for  $C_{4k+3}$  with span equal to the desired number. First, we label the vertices  $x_0, x_2, \dots, x_{4k+2}$ , by  $\pi(0) = 0$  and  $f(x_0) = 0$ ; and for  $1 \leq i \leq 2k + 1$ ,

$$\pi(2i) = \begin{cases} (\pi(2i - 2) + k + 1) \pmod n, & \text{if } i \text{ is odd;} \\ (\pi(2i - 2) + k) \pmod n, & \text{if } i \text{ is even,} \end{cases}$$

$$f(x_{2i}) = \begin{cases} f(x_{2i-2}) + k, & \text{if } i \text{ is odd;} \\ f(x_{2i-2}) + k + 1, & \text{if } i \text{ is even.} \end{cases}$$

Secondly, we label the remaining vertices by  $\pi(1) = 2k + 2$  and  $f(x_1) = 0$ ; and for  $1 \leq i \leq 2k$ ,

$$\pi(2i + 1) = (\pi(2i - 1) + k + 1) \pmod n, \text{ and } f(x_{2i+1}) = i(k + 1).$$




Figure 3: An antipodal labeling for  $C_{11}$  with minimum span  $\text{an}(C_{11}) = 12$ .

See Figure 3 for an example.

By some calculation, one gets

$$\pi(2i + 1) \equiv (i + 2)(k + 1) \pmod{n}, \text{ for } 0 \leq i \leq 2k, \text{ and}$$

$$\pi(2i) \equiv \begin{cases} -(i - 2)(k + 1) \pmod{n}, & \text{if } i \text{ is odd;} \\ -i(k + 1) \pmod{n}, & \text{if } i \text{ is even,} \end{cases}$$

for  $0 \leq i \leq 2k + 1$ . Hence, we conclude

$$\{\pi(i) : 0 \leq i \leq 4k + 2\} = \{j(k + 1) \pmod{n} : -2k \leq j \leq 2k + 2\}.$$

Because  $\gcd(n, k + 1) = 1$ ,  $\pi$  is a permutation. Similar to the proof of Theorem 4, it is straightforward to show that  $f$  is an antipodal labeling, and we shall leave the details to the reader. This completes the proof of Theorem 5.  $\square$

## 5 $n = 4k$

Note, it is trivial that  $\text{an}(C_4) = 1$ . For cycles with  $n = 4k$  nodes,  $k \geq 2$ , we improve the lower bound in Corollary 3 and give an upper bound.

**Theorem 9** For every integer  $k \geq 2$ ,

$$2k^2 - \lfloor k/2 \rfloor \leq \text{an}(C_{4k}) \leq 2k^2 - 1.$$

The following lemma will be used to prove the lower bound for  $\text{an}(C_{4k})$  in Theorem 9. Recall that  $d = \text{diam}(C_{4k}) = 2k$ .

**Lemma 10** Let  $f$  be an antipodal labeling of  $C_{4k}$ , for some integer  $k \geq 2$ . If  $f_i + f_{i+1} = k$  for some  $0 \leq i \leq n - 3$ , then  $d(x_i, x_{i+2}) = k$ .

**Proof.** Assume  $f_i + f_{i+1} = k$  for some  $0 \leq i \leq n - 3$ . Then  $d(x_i, x_{i+2}) \geq d - (f_i + f_{i+1}) = k$ . On the other hand, by Proposition 1 and definition,

$$\begin{aligned} d(x_i, x_{i+2}) &\leq n - (d_i + d_{i+1}) \\ &\leq 4k - (d - f_i + d - f_{i+1}) \\ &= k. \end{aligned}$$

□

**Lemma 11** Let  $f$  be an antipodal labeling of  $C_{4k}$ ,  $k \geq 2$ . Then for any  $0 \leq i \leq n - 9$ ,

$$\sum_{j=0}^7 f_{i+j} \geq 4k + 1.$$

**Proof.** Assume to the contrary, for some  $0 \leq i \leq n - 9$ , we have  $\sum_{j=0}^7 f_{i+j} \leq 4k$ . By Lemma 2,  $f_i + f_{i+1} = f_{i+2} + f_{i+3} = f_{i+4} + f_{i+5} = f_{i+6} + f_{i+7} = k$ . By Lemma 10,  $d(x_i, x_{i+2}) = d(x_{i+2}, x_{i+4}) = d(x_{i+4}, x_{i+6}) = d(x_{i+6}, x_{i+8}) = k$ . Since  $n = 4k$ , it is impossible that all these four equations hold. So the result follows. □

**Lemma 12** Let  $f$  be an antipodal labeling of  $C_{4k}$ ,  $k \geq 2$ . The following are true.

- (1) If  $f_i + f_{i+1} = k$  for some  $0 \leq i \leq n - 4$ , then  $f_{i+2} \geq f_i$ .
- (2) If  $f_i + f_{i+1} = k$  for some  $1 \leq i \leq n - 3$ , then  $f_{i-1} \geq f_{i+1}$ .

(3) If  $f_i + f_{i+1} + f_{i+2} + f_{i+3} = 2k$  for some  $0 \leq i \leq n-6$ , then  $f_{i+4} \geq f_i \geq 1$ .

(4) If  $\sum_{j=0}^7 f_{i+j} = 4k+1$  for some  $0 \leq i \leq n-10$ , then  $f_{i+8} \geq f_i$ .

(5) If  $\sum_{j=0}^7 f_{i+j} = 4k+1$  for some  $0 \leq i \leq n-10$ , then  $f_{i+8} \geq 1$ .

(6) For any  $0 \leq i \leq n-6$ ,  $\sum_{j=0}^4 f_{i+j} \geq 2k+1$ .

(7) For any  $0 \leq i \leq n-10$ ,  $\sum_{j=0}^8 f_{i+j} \geq 4k+2$ .

**Proof.** To prove (1), assume  $f_i + f_{i+1} = k$  for some  $0 \leq i \leq n-4$ . By Lemma 2,  $f_{i+2} + f_{i+1} \geq k = f_{i+1} + f_i$ , hence  $f_{i+2} \geq f_i$ . (2) follows by a similar argument.

To prove (3), assume  $f_i + f_{i+1} + f_{i+2} + f_{i+3} = 2k$  for some  $0 \leq i \leq n-6$ . Then by Lemma 2,  $f_i + f_{i+1} = f_{i+2} + f_{i+3} = k$ . By Lemma 10,  $d(x_i, x_{i+2}) = d(x_{i+2}, x_{i+4}) = k$ , so  $d(x_i, x_{i+4}) = 2k$ . This implies that  $d_i < 2k$ , as  $n = 4k$ . By definition of antipodal labeling,  $f_i \geq 1$ . Hence, by (1), we have  $f_{i+4} \geq f_{i+2} \geq f_i \geq 1$ .

To prove (4), assume  $\sum_{j=0}^7 f_{i+j} = 4k+1$  for some  $0 \leq i \leq n-10$ . By Lemma 11,  $\sum_{j=1}^8 f_{i+j} \geq 4k+1 = \sum_{j=0}^7 f_{i+j}$ , hence  $f_{i+8} \geq f_i$ .

To prove (5), assume  $\sum_{j=0}^7 f_{i+j} = 4k+1$  for some  $0 \leq i \leq n-10$ . By Lemma 2, we have  $f_i + f_{i+1} = f_{i+2} + f_{i+3} = k$  or  $f_{i+4} + f_{i+5} = f_{i+6} + f_{i+7} = k$ . For the former case, the result follows by (4) and (3); for the latter case, the results follows by (3).

(6) follows by (3) and Lemma 2; and (7) follows by (5) and Lemma 11.  $\square$

**Corollary 13** For any integer  $k \geq 2$ ,  $\text{an}(C_{4k}) \geq 2k^2 - \lfloor k/2 \rfloor$ .

**Proof.** For  $k = 2$ , by Lemma 2 and Lemma 12 (6), the span of an antipodal labeling  $f$  for  $C_8$  has  $f(x_7) = (f_0 + f_1 + \cdots + f_4) + (f_5 + f_6) \geq 5 + 2 = 2k^2 - \lfloor k/2 \rfloor$ .

For  $k \geq 3$ , by Lemmas 2, 11 and 12 (7), the span of an antipodal labeling  $f$  for  $C_{4k}$  has

$$\begin{aligned} f(x_{4k-1}) &= \sum_{i=0}^8 f_i + \sum_{i=1}^{\lfloor (4k-9)/8 \rfloor} \left( \sum_{j=1}^8 (f_{8i+j}) \right) \\ &\quad + f_{8\lfloor (4k-1)/8 \rfloor + 1} + f_{8\lfloor (4k-1)/8 \rfloor + 2} + \cdots + f_{4k-2} \\ &\geq \begin{cases} (4k+2) + [2k^2 - (11/2)k - 3/2] + k, & k \text{ is odd} \\ (4k+2) + [2k^2 - (15/2)k - 2] + 3k, & k \text{ is even} \end{cases} \\ &= 2k^2 - \lfloor k/2 \rfloor. \end{aligned}$$

□

**Proof of Theorem 9.** It remains to find an antipodal labeling for  $C_{4k}$  with span  $2k^2 - 1$ . First, we label the vertices  $x_0, x_2, \dots, x_{4k-2}$ , by  $\pi(0) = 0$  and  $f(x_0) = 0$ ; and for  $1 \leq i \leq 2k - 1$ ,

$$\pi(2i) = \begin{cases} (\pi(2i-2) + k) \pmod n, & \text{if } i \text{ is odd;} \\ (\pi(2i-2) + k + 1) \pmod n, & \text{if } i \text{ is even,} \end{cases}$$

$$f(x_{2i}) = \begin{cases} f(x_{2i-2}) + k, & \text{if } i \text{ is odd;} \\ f(x_{2i-2}) + k + 1, & \text{if } i \text{ is even.} \end{cases}$$

Secondly, we label the remaining vertices by: For  $0 \leq i \leq 2k - 1$ ,

$$\pi(2i+1) = (\pi(2i) + 2k) \pmod n, \text{ and } f(x_{2i+1}) = f(x_{2i}).$$

See Figure 4 for an example.

By calculation, one gets the following for  $0 \leq i \leq k - 1$ :

$$\begin{aligned} \pi(4i) &\equiv i(2k+1) \pmod n. \\ \pi(4i+1) &\equiv (i+2k)(2k+1) \pmod n. \\ \pi(4i+2) &\equiv \begin{cases} (i+3k)(2k+1) \pmod n, & \text{if } k \text{ is odd;} \\ (i+k)(2k+1) \pmod n, & \text{if } k \text{ is even.} \end{cases} \\ \pi(4i+3) &\equiv \begin{cases} (i+k)(2k+1) \pmod n, & \text{if } k \text{ is odd;} \\ (i+3k)(2k+1) \pmod n, & \text{if } k \text{ is even.} \end{cases} \end{aligned}$$




Figure 4: An antipodal labeling for  $C_{12}$  with minimum span  $\text{an}(C_{12}) = 17$ .

Therefore, we conclude

$$\{\pi(i) : 0 \leq i \leq 4k - 1\} = \{j(2k + 1) \bmod n : 0 \leq j \leq 4k - 1\}.$$

Because  $\gcd(n, 2k + 1) = 1$ ,  $\pi$  is a permutation. Similar to the proof of Theorem 4, it is straightforward to show that  $f$  is an antipodal labeling, and we shall leave the details to the reader. This completes the proof of Theorem 9.  $\square$

We conjecture that  $\text{an}(C_{4k})$  is equal to the upper bound in Theorem 9.

**Conjecture 1** For any  $k \geq 1$ ,  $\text{an}(C_{4k}) = 2k^2 - 1$ .

A case analysis has confirmed the above conjecture for  $k \leq 5$ .

## References

- [1] T. Calamoneri and R. Petreschi, “ $L(2, 1)$ -labeling of planar graphs,” *ACM* (2001), 28–33.
- [2] G. J. Chang, and C. Lu, “Distance-two labelings of graphs,” *European J. of Combin.*, 24 (2003), 53–58.

- [3] G. J. Chang and D. Kuo, "The  $L(2, 1)$ -labeling problem on graphs," *SIAM J. Discrete Math.*, 9 (1996), 309–316.
- [4] G. Chartrand, D. Erwin, and P. Zhang, "Radio antipodal colorings of cycles," *Cong. Numer.*, 144 (2000), 129–141.
- [5] G. Chartrand, D. Erwin, and P. Zhang, "Radio antipodal colorings of graphs," *Math. Bohem.*, 127 (2002), 57–69.
- [6] G. Chartrand, D. Erwin, and P. Zhang, "Radio labelings of graphs," *Bull. Inst. Combin. Appl.*, 33 (2001), 77–85.
- [7] G. Chartrand, D. Erwin, and P. Zhang, "Radio  $k$ -colorings of paths," *Discuss Math. Graph Theory*, 24 (2004), 5–21.
- [8] G. Chartrand, D. Erwin, and P. Zhang, "A graph labeling problem suggested by FM channel restrictions," *Bull. Inst. Combin. Appl.*, 43 (2005), 43–57.
- [9] Z. Furedi, J. Griggs, and D. Kleitman, "Pair labelings with given distance," *SIAM J. Discrete Math.*, 4 (1989), 491–499.
- [10] J. Georges, D. Mauro, and M. Whittlesey, "Relating path covering to vertex labelings with a condition at distant two," *Discrete Math.*, 135 (1994), 103–111.
- [11] J. Georges, D. Mauro, and M. Whittlesey, "On the  $\lambda$ -number of  $Q_n$  and related graphs," *SIAM J. Discrete Math.*, 8 (1995), 499–506.
- [12] J. Griggs and R. Yeh, "Labeling graphs with a condition at distance 2," *SIAM J. Discrete Math.*, 5 (1992), 586–595.
- [13] R. Khennoufa and O. Tongni, "A note on radio antipodal colourings of paths," *Math. Bohem.*, 130 (2005), 277–282.
- [14] D. Liu, and R.K. Yeh, "On distance two labelings of graphs," *Ars Combin.*, 47 (1997), 13–22.
- [15] D. Liu, "Radio number for trees," *Discrete Math.*, to appear.