

On the factorization numbers of some finite p -groups

Marius Tărnăuceanu

March 8, 2012

Abstract

This note deals with the computation of the factorization number $F_2(G)$ of a finite group G . By using the Möbius inversion formula, explicit expressions of $F_2(G)$ are obtained for two classes of finite abelian groups, improving the results of *Factorization numbers of some finite groups*, Glasgow Math. J. (2012).

MSC (2010): Primary 20D40; Secondary 20D60.

Key words: factorization number, subgroup commutativity degree, Möbius function, finite abelian group.

1 Introduction

Let G be a group, $L(G)$ be the subgroup lattice of G and H, K be two subgroups of G . If $G = HK$, then G is said to be *factorized* by H and K and the expression $G = HK$ is said to be a *factorization* of G . Denote by $F_2(G)$ the *factorization number* of G , that is the number of all factorizations of G .

The starting point for our discussion is given by the paper [3], where $F_2(G)$ has been computed for certain classes of finite groups. The connection between $F_2(G)$ and the subgroup commutativity degree $sd(G)$ of G (see [5, 7]) has been also established, namely

$$sd(G) = \frac{1}{|L(G)|^2} \sum_{H \leq G} F_2(H).$$

Obviously, by applying the well-known Möbius inversion formula to the above equality, one obtains

$$(1) \quad F_2(G) = \sum_{H \leq G} sd(H) |L(H)|^2 \mu(H, G).$$

In particular, if G is abelian, then we have $sd(H) = 1$ for all $H \in L(G)$, and consequently

$$(2) \quad F_2(G) = \sum_{H \leq G} |L(H)|^2 \mu(H, G) = \sum_{H \leq G} |L(G/H)|^2 \mu(H).$$

This formula will be used in the following to calculate the factorization numbers of an elementary abelian p -group and of a rank 2 abelian p -group, improving Theorem 1.2 and Corollary 2.5 of [3]. An interesting conjecture about the maximum value of $F_2(G)$ on the class of p -groups of the same order will be also presented.

First of all, we recall a theorem due to P. Hall [1] (see also [2]), that permits us to compute explicitly the Möbius function of a finite p -group.

Theorem 1. *Let G be a finite p -group of order p^n . Then $\mu(G) = 0$ unless G is elementary abelian, in which case we have $\mu(G) = (-1)^n p^{\binom{n}{2}}$.*

In contrast with Theorem 1.2 of [3] that gives only a recurrence relation satisfied by $F_2(\mathbb{Z}_p^n)$, $n \in \mathbb{N}$, we are able to determine precise expressions of these numbers.

Theorem 2. *We have*

$$(3) \quad F_2(\mathbb{Z}_p^n) = \sum_{i=0}^n (-1)^i a_{n,p}(i) a_{n-i,p}^2 p^{\binom{i}{2}},$$

where $a_{n,p}(i)$ is the number of subgroups of order p^i of \mathbb{Z}_p^n , $a_{n,p}$ is the total number of subgroups of \mathbb{Z}_p^n , and, by convention, $\binom{i}{2} = 0$ for $i = 0, 1$.

Since the numbers $a_{n,p}(i)$, $i = 0, 1, \dots, n$, are well-known, namely

$$a_{n,p}(i) = \frac{(p^n - 1) \cdots (p - 1)}{(p^i - 1) \cdots (p - 1)(p^{n-i} - 1) \cdots (p - 1)},$$

the equality (3) easily leads to the following values of $F_2(\mathbb{Z}_p^n)$ for $n = 1, 2, 3, 4$.

Examples.

a) $F_2(\mathbb{Z}_p) = 3.$

b) $F_2(\mathbb{Z}_p^2) = p^2 + 3p + 5.$

c) $F_2(\mathbb{Z}_p^3) = 3p^4 + 4p^3 + 8p^2 + 5p + 7.$

d) $F_2(\mathbb{Z}_p^4) = p^8 + 3p^7 + 9p^6 + 11p^5 + 14p^4 + 15p^3 + 12p^2 + 23p + 9.$

Next we compute the factorization number of a rank 2 abelian p -group.

Theorem 3. *The factorization number of the finite abelian p -group $\mathbb{Z}_{p^{\alpha_1}} \times \mathbb{Z}_{p^{\alpha_2}}$, $\alpha_1 \leq \alpha_2$, is given by the following equality:*

$$F_2(\mathbb{Z}_{p^{\alpha_1}} \times \mathbb{Z}_{p^{\alpha_2}}) = \frac{1}{(p-1)^4} \left[(2\alpha_2 - 2\alpha_1 + 1)p^{2\alpha_1+4} - (6\alpha_2 - 6\alpha_1 + 1)p^{2\alpha_1+3} + \right. \\ \left. + (6\alpha_2 - 6\alpha_1 - 1)p^{2\alpha_1+2} - (2\alpha_2 - 2\alpha_1 - 1)p^{2\alpha_1+1} - (2\alpha_1 + 2\alpha_2 + 3)p^3 + \right. \\ \left. + (6\alpha_1 + 6\alpha_2 + 7)p^2 - (6\alpha_1 + 6\alpha_2 + 5)p + (2\alpha_1 + 2\alpha_2 + 1) \right].$$

We remark that Theorem 3 gives a generalization of Corollary 2.5 of [3]. Indeed, by taking $\alpha_1 = 1$ and $\alpha_2 = n$ in the above formula, one obtains:

Corollary 4. $F_2(\mathbb{Z}_p \times \mathbb{Z}_{p^n}) = (2n - 1)p^2 + (2n + 1)p + (2n + 3).$

Finally, we will focus on the minimum/maximum of $F_2(G)$ when G belongs to the class of p -groups of order p^n . It is easy to see that

$$2n + 1 = F_2(\mathbb{Z}_{p^n}) \leq F_2(G).$$

For $n \leq 3$ the greatest value of $F_2(G)$ is obtained for $G \cong \mathbb{Z}_p^n$, as shows the following result.

Theorem 5. *Let G be a finite p -group of order p^n . If $n \leq 3$, then*

$$F_2(G) \leq F_2(\mathbb{Z}_p^n).$$

Inspired by Theorem 5, we came up with the following conjecture, which we also have verified for several $n \geq 4$ and particular values of p .

Conjecture 6. For every finite p -group G of order p^n , we have

$$F_2(G) \leq F_2(\mathbb{Z}_p^n).$$

We end our note by indicating a natural problem concerning the factorization number of abelian p -groups.

Open problem. Compute explicitly $F_2(G)$ for an arbitrary finite abelian p -group G . Given a positive integer n , two partitions τ, τ' of n and denoting by G, G' the abelian p -groups of order p^n induced by τ and τ' , respectively, is it true that $F_2(G) \geq F_2(G')$ if and only if $\tau \preceq \tau'$ (where \preceq denotes the lexicographic order)?

2 Proofs of the main results

Proof of Theorem 2. By using Theorem 1 in (2), it follows that

$$\begin{aligned} F_2(\mathbb{Z}_p^n) &= \sum_{H \leq \mathbb{Z}_p^n} |L(\mathbb{Z}_p^n/H)|^2 \mu(H) = \sum_{i=0}^n \sum_{\substack{H \leq \mathbb{Z}_p^n \\ |H|=p^i}} |L(\mathbb{Z}_p^n/H)|^2 \mu(H) = \\ &= \sum_{i=0}^n a_{n,p}(i) |L(\mathbb{Z}_p^{n-i})|^2 (-1)^i p^{\binom{i}{2}} = \sum_{i=0}^n (-1)^i a_{n,p}(i) a_{n-i,p}^2 p^{\binom{i}{2}}, \end{aligned}$$

as desired. ■

Proof of Theorem 3. It is well-known that $G = \mathbb{Z}_{p^{\alpha_1}} \times \mathbb{Z}_{p^{\alpha_2}}$ has a unique elementary abelian subgroup of order p^2 , say M , and that

$$G/M \cong \mathbb{Z}_{p^{\alpha_1-1}} \times \mathbb{Z}_{p^{\alpha_2-1}}.$$

Moreover, all elementary abelian subgroups of G are contained in M . Denote by $M_i, i = 1, 2, \dots, p+1$, the minimal subgroups of G . Then every quotient G/M_i is isomorphic to a maximal subgroup of G and therefore we may assume that

$$G/M_i \cong \mathbb{Z}_{p^{\alpha_1-1}} \times \mathbb{Z}_{p^{\alpha_2}} \text{ for } i = 1, 2, \dots, p$$

and

$$G/M_{p+1} \cong \mathbb{Z}_{p^{\alpha_1}} \times \mathbb{Z}_{p^{\alpha_2-1}}.$$

Clearly, the equality (2) becomes

$$F_2(G) = |L(G/M)|^2 \mu(M) + \sum_{i=1}^{p+1} |L(G/M_i)|^2 \mu(M_i) + |L(G)|^2 \mu(1),$$

in view of Theorem 1. Since by Theorem 2 we have $\mu(M) = \mu(\mathbb{Z}_p^2) = p$, $\mu(M_i) = \mu(\mathbb{Z}_p) = -1$, for all $i = \overline{1, p+1}$, and $\mu(1) = 1$, one obtains

$$(4) \quad F_2(G) = p |L(\mathbb{Z}_{p^{\alpha_1-1}} \times \mathbb{Z}_{p^{\alpha_2-1}})|^2 - p |L(\mathbb{Z}_{p^{\alpha_1-1}} \times \mathbb{Z}_{p^{\alpha_2}})|^2 - \\ - |L(\mathbb{Z}_{p^{\alpha_1}} \times \mathbb{Z}_{p^{\alpha_2-1}})|^2 + |L(\mathbb{Z}_{p^{\alpha_1}} \times \mathbb{Z}_{p^{\alpha_2}})|^2.$$

The total number of subgroups of $\mathbb{Z}_{p^{\alpha_1}} \times \mathbb{Z}_{p^{\alpha_2}}$ has been computed in Theorem 3.3 of [6], namely

$$\frac{1}{(p-1)^2} [(\alpha_2 - \alpha_1 + 1)p^{\alpha_1+2} - (\alpha_2 - \alpha_1 - 1)p^{\alpha_1+1} - (\alpha_1 + \alpha_2 + 3)p + (\alpha_1 + \alpha_2 + 1)].$$

Then the desired formula follows immediately by a direct calculation in the right side of (4). ■

Proof of Theorem 5. For $n = 2$ we obviously have

$$F_2(\mathbb{Z}_{p^2}) = 5 < F_2(\mathbb{Z}_p^2) = p^2 + 3p + 5.$$

For $n = 3$ it is well-known (see e.g. (4.13), [4], II) that G can be one of the following groups:

- $\mathbb{Z}_2^3, \mathbb{Z}_2 \times \mathbb{Z}_4, \mathbb{Z}_8, D_8$ and Q_8 if $p = 2$;
- $\mathbb{Z}_p^3, \mathbb{Z}_p \times \mathbb{Z}_{p^2}, \mathbb{Z}_{p^3}, M(p^3) = \langle x, y \mid x^{p^2} = y^p = 1, y^{-1}xy = x^{p+1} \rangle$ and $E(p^3) = \langle x, y \mid x^p = y^p = [x, y]^p = 1, [x, y] \in Z(E(p^3)) \rangle$ if $p \geq 3$.

By using the results in Section 2 of [3], one obtains

for $p = 2$:

$$F_2(\mathbb{Z}_2^3) = 129 > F_2(\mathbb{Z}_2 \times \mathbb{Z}_4) = 29, F_2(\mathbb{Z}_8) = 7, F_2(D_8) = 41, F_2(Q_8) = 17$$

and

for $p \geq 3$:

$$F_2(\mathbb{Z}_p^3) = 3p^4 + 4p^3 + 8p^2 + 5p + 7 > F_2(\mathbb{Z}_p \times \mathbb{Z}_{p^2}) = F_2(M(p^3)) = 3p^2 + 5p + 7,$$

$$F_2(\mathbb{Z}_{p^3}) = 7.$$

We also observe that $E(p^3)$ has $p + 1$ elementary abelian subgroups of order p^2 , say M_1, M_2, \dots, M_{p+1} , and that every M_i contains $p + 1$ subgroups of order p , namely $\Phi(E(p^3))$ and M_{ij} , $j = 1, 2, \dots, p$. Then $|L(E(p^3))| = p^2 + 2p + 4$ and so

$$F_2(E(p^3)) < |L(E(p^3))|^2 = p^4 + 4p^3 + 12p^2 + 16p + 16.$$

On the other hand, we can easily see that this quantity is less than $F_2(\mathbb{Z}_p^3)$ for all primes $p \geq 3$, completing the proof. \blacksquare

Remark. It is clear that an explicit formula for $F_2(E(p^3))$ cannot be obtained by applying (2), but we are able to determine it by a direct computation. The factorization pairs of $E(p^3)$ are:

- $(1, E(p^3)), (E(p^3), 1)$;
- $(M_{ij}, M_{i'}) \forall i' \neq i, (M_{ij}, E(p^3)), (E(p^3), M_{ij}), i = \overline{1, p+1}, j = \overline{1, p}$;
- $(\Phi(E(p^3)), E(p^3)), (E(p^3), \Phi(E(p^3)))$;
- $(M_i, M_{i'}) \forall i' \neq i, j = 1, 2, \dots, p, (M_i, M_{i'}) \forall i' \neq i, (M_i, E(p^3))$ and $(M_i, E(p^3)), i = \overline{1, p+1}$;
- $(E(p^3), E(p^3))$.

Hence

$$F_2(E(p^3)) = 2 + p(p+1)(p+2) + 2 + (p+1)(p^2 + p + 2) + 1 =$$

$$= 2p^3 + 5p^2 + 5p + 7.$$

Acknowledgements. The author is grateful to the reviewer for its remarks which improve the previous version of the paper.

References

- [1] Hall, P., *A contribution to the theory of groups of prime-power order*, Proc. London Math. Soc. **36** (1933), 29-95.
- [2] Hawkes, T., Isaacs, I.M., Özaydin, M., *On the Möbius function of a finite group*, Rocky Mountain J. Math. **19** (1989), 1003-1033.
- [3] Saeedi, F., Farrokhi D.G., M., *Factorization numbers of some finite groups*, to appear in Glasgow Math. J. (2012).
- [4] Suzuki, M., *Group theory*, I, II, Springer Verlag, Berlin, 1982, 1986.
- [5] Tărnăuceanu, M., *Subgroup commutativity degrees of finite groups*, J. Algebra **321** (2009), 2508-2520, doi: 10.1016/j.jalgebra.2009.02.010.
- [6] Tărnăuceanu, M., *An arithmetic method of counting the subgroups of a finite abelian group*, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) **53/101** (2010), 373-386.
- [7] Tărnăuceanu, M., *Addendum to "Subgroup commutativity degrees of finite groups"*, J. Algebra **337** (2011), 363-368, doi: 10.1016/j.jalgebra.2011.05.001.

Marius Tărnăuceanu
Faculty of Mathematics
"A.I. Cuza" University
Iași, Romania
e-mail: tarnauc@uaic.ro