

The Zero Forcing Number of Quartic Circulant Graphs of Prime Order

Huixian Li^{1,✉}, Guang Li¹, Shengjin Ji¹

¹ School of Mathematics and Statistics Shandong University of Technology, Zibo, China

ABSTRACT

Let G be a graph, the zero forcing number $Z(G)$ is the minimum of $|Z|$ over all zero forcing sets $Z \subseteq V(G)$. In this paper, we are interested in studying the zero forcing number of quartic circulant graphs $C_p(s, t)$, where p is an odd prime. Based on the fact that $C_p(s, t) \cong C_p(1, q)$, we give the exact values of the zero forcing number of some specific quartic circulant graphs.

Keywords: Zero forcing set, Zero forcing number, Circulant graphs

2020 Mathematics Subject Classification: 05C30

1. Introduction

In this paper, all graphs considered are finite, undirected and simple. For any terminology used but not defined here, one may refer to [5].

Graph parameters have many application in some science trends. As one of the most important, it can be *zero forcing number*. Also, zero forcing number have applications in power domination problem, logic circuits, information in social networks and so on, see [6, 4, 14]. We now formally introduce its definition. Let G be a graph and Z be an subset of $V(G)$. Color the vertices of Z and $V(G) \setminus Z$ black and white initially, respectively. A *forcing process* is defined as follows: for a black vertex u of $V(G)$, if u has exactly one

✉ Corresponding author.

E-mail addresses: jishengjin2013@163.com (Shengjin Ji).

Received 9 March 2024; accepted 5 October 2024; published 31 December 2024.

DOI: [10.61091/ars161-01](https://doi.org/10.61091/ars161-01)

© 2024 The Author(s). Published by Combinatorial Press. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>).

white neighbor v , then reassign v black. At the time, we say u forces v and denote it by $u \rightarrow v$. The subset Z is called a *zero forcing set*, if all vertices in $V(G) \setminus Z$ are forced black by repeatedly applying the forcing process. The *zero forcing number* of G is the minimum of $|Z|$ over all zero forcing sets $Z \subseteq V(G)$, denoted by $Z(G)$.

The computation of zero forcing number is NP-hard, see [9]. It has attracted many scholars, and a lot of results are obtained in the field. In [1], some elementary results of graphs regarding zero forcing are obtained, such as, trees, Cartesian products of two paths. Kalinowski et al. ([16]) showed that $Z(G) \geq n(1 + \frac{2\lambda_{min}}{d-\lambda_{min}})$ if G is d -regular. In [2, 3], if a graph G has the minimum and maximum degree δ and Δ respectively, then the zero forcing number of a graph G with n vertices can be as $\delta \leq Z(G) \leq \frac{n\Delta}{\Delta+1}$. In addition, Meyer [20] obtained upper and lower bounds for $Z(G)$ when G is a bipartite circulant and showed which graphs achieve the equality in the lower bound. From these references, we can find that, even for some special graphs, it seems to be impossible to determine the exact number of zero forcing number. Note that, Riddell [23] obtained some results of the zero forcing number on cubic circulant graphs. Hence determining the zero forcing number for all quartic circulant graphs is an interesting and challenging problem. For more details on zero forcing, see [9, 10, 12, 13, 19, 24, 7, 25].

Given an integer $n \geq 2$ and a subset $S \subseteq \left\{1, 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor\right\}$, a circulant graph $G = C_n(S)$ is formally defined as a graph with vertex set $V(G) = \{0, 1, \dots, n-1\}$ and edge set $E(G) = \{\{i, i \pm j\} | i \in \{0, \dots, n-1\} \text{ and } j \in S\}$, where i, j and $i \pm j$ are taken modulo n . Note that if $S = \{s_1, \dots, s_t\}$, then we can always assume that $s_1 < s_2 < \dots < s_t$. In particular, if $S = \{s, t\}$, the circulant graph $C_n(s, t)$ is a quartic circulant graph, where $1 \leq s < t \leq \left\lfloor \frac{n}{2} \right\rfloor$, that is to say, for any $i \in \{0, \dots, n-1\}$, the neighbours of i be $\{i \pm s, i \pm t\}$, where $\{i \pm s, i \pm t\}$ are taken modulo n .

The *distance* between two vertices u and v is the length of the shortest path from u to v , denoted by $d(u, v)$. The *diameter* of G is the maximum distance among all pairs of vertices in G , denoted by $diam(G)$. If u is a vertex of G , then let $N_i(u)$ denote the set of vertices at distance i from u . The partition $\{N_0(u), N_1(u), \dots, N_d(u)\}$ is called the *distance partition* with respect to u . Clearly, $N_0(u) = \{u\}$. The set of neighbours of a vertex v in a graph G is denoted by $N(v)$. Studying the properties of circulant graphs has become an active topic in algebraic graph theory for a long time. Actually, they are close related with graph parameters and networks etc, such as, [15, 18, 22, 27]. Let Z_p be the additive group of integers modulo p , where p be an odd prime, $Z_p^* = Z_p \setminus \{0\}$. Some examples of circulant graphs are given in Figure 1.




Fig. 1. The circulant graphs $C_7(1, 2)$ and $C_{11}(2, 5)$

We first characterize the zero forcing number of quartic circulant graphs $C_p(1, q)$ on

groups G with some special q .

Theorem 1.1. *Let $G = C_p(1, q)$ with $2 \leq q \leq \lfloor \frac{p}{2} \rfloor$.*

(1) *If $q = 2$, then $Z(C_p(1, 2)) = 4$.*

(2) *If $q = 3$, then*

$$Z(C_p(1, 3)) = \begin{cases} 4, & p = 7, \\ 6, & p \geq 11. \end{cases}$$

(3) *If $q = 4$, then*

$$Z(C_p(1, 4)) = \begin{cases} 6, & p = 11 \text{ or } 13, \\ 8, & p \geq 17. \end{cases}$$

(4) *If $q = 5$, then*

$$Z(C_p(1, 5)) = \begin{cases} 4, & p = 11, \\ 6, & p = 13 \text{ or } 17, \\ 8, & p = 19 \text{ or } 23, \\ 10, & p \geq 29. \end{cases}$$

Motivated by the above result, we have the following conjecture: when $|p - q|$ is sufficiently large, $Z(C_p(1, q)) = 2q$.

The rest of the paper is structured as follows. In Section 2 we give some preliminary results. We focus on the isomorphism of quartic circulant graphs, and conclude that $C_p(s, t) \cong C_p(1, q)$ for some $q \in Z_p^*$. Inspired by the idea of the isomorphism of those graphs, the zero forcing number of $C_p(1, q)$ can help us research the zero forcing number of $C_p(s, t)$. In Section 3 we prove Theorem 1.1.

2. Preliminary Results

In this section, we will characterize the isomorphism of (quartic) circulant graphs. Assume that $C_p(s_1, s_2, \dots, s_t)$ be a circulant graph of prime order p with $1 \leq s_i \leq \lfloor \frac{p}{2} \rfloor$, $1 \leq i \leq t$. We first propose the following result.

Lemma 2.1. [17] *Let Z_p be the additive group of integers modulo p , where p be an odd prime. Then $\text{Aut}((Z_p, +)) \cong (Z_p^*, \times)$.*

By [8] and [21], together with Lemma 2.1, we have the following useful result, which characterize the isomorphism of circulant graphs with prime vertices.

Theorem 2.2. [8, 21] *Let $G = C_p(s_1, s_2, \dots, s_t)$ be a circulant graph, where p be an odd prime, $2 \leq s_i \leq \lfloor \frac{p}{2} \rfloor$. Then $C_p(s_1, s_2, \dots, s_t) \cong C_p(1, q_2, \dots, q_t)$, for some $q_i \in Z_p^*$.*

Based on Theorem 2.2, for quartic circulant graphs, we can get the following conclusion.

Theorem 2.3. Let $C_p(s, t)$ be a quartic circulant graph with $2 \leq s < t \leq \lfloor \frac{p}{2} \rfloor$. Then $C_p(s, t) \cong C_p(1, q)$ for some $q \in Z_p^*$.

As an example of Theorem 2.3, if $p = 11$ and $s = 2, t = 5$, then $C_{11}(2, 5) \cong C_{11}(1, 3)$. Therefore, the characterization of the zero forcing number of quartic circulant graphs $C_p(s, t)$ can be converted into the zero forcing number of quartic circulant graphs $C_p(1, q)$.




Fig. 2. The circulant graph $C_{11}(2, 5) \cong C_{11}(1, 3)$

3. The Zero Forcing Number of Circulant Graphs $C_p(1, q)$

Motivated by Theorem 2.3, we will characterize the zero forcing number of circulant graphs $C_p(1, q)$. In this section, we first give the upper bound of the zero forcing number of $C_p(1, q)$. For the rest of this section, we will give the exact values of the zero forcing number for $C_p(1, 2)$, $C_p(1, 3)$, $C_p(1, 4)$, and $C_p(1, 5)$, respectively.

3.1. The zero forcing number of $C_p(1, q)$ with $q = 2, 3$

In this subsection, we characterize the exact values of the zero forcing number of the circulant graphs $C_p(1, 2)$ and $C_p(1, 3)$. For consecutive circulant graphs, we have the following conclusions.

Theorem 3.1. [26] If $G = C_p(1, q)$, then $Z(G) \leq 2q$.

Lemma 3.2. [8] If $G = C_n(1, 2, \dots, d)$, $1 \leq d \leq \lfloor \frac{n}{2} \rfloor$, then $Z(G) = 2d$.

By Lemma 3.2, we can get the following theorem immediately.

Theorem 3.3. If $G = C_p(1, 2)$, then $Z(C_p(1, 2)) = 4$.

Remark 3.4. Observe that $\{0, 1, 2, p-1\}$ is a zero forcing set of $C_p(1, 2)$.

Theorem 3.5. Let $G = C_n(1, m)$ be a circulant graph, where $n=2m+1$, then $Z(C_n(1, m)) = 4$.

Proof. By [8, Lemma 1.2], $Z(C_n(1, m)) \geq 4$. In the following, we can get a forcing set

of size 4, that is to say $Z(C_n(1, m)) \leq 4$.

Step 1. Black four vertices $\{0, 1, m, 2m\}$, then $m + 1$ be the unique neighbour not be colored black of vertex 0, thus $0 \rightarrow m + 1$, see Figure 3.




Fig. 3. Step 1

Step 2. After the above forcing process, $m + 1$ has the unique neighbour $m + 2$ not be colored black, thus $m + 1 \rightarrow m + 2$, see Figure 4.




Fig. 4. Step 2

Step 3. After this forcing process, 1 has the unique neighbour 2 not be colored black, thus $1 \rightarrow 2$.

Step 4. Similarly, the black vertex $m + 2$ has the unique neighbour $m + 3$ not be colored black, thus $m + 2 \rightarrow m + 3$. Then the black vertex 2 has the unique neighbour 3 not be colored black, thus $2 \rightarrow 3$. Continue the similiar analysis, the vertices $m + 4, 4, \dots, 2m - 1, m - 1$ will be colored black by the forcing process of $m + 3, 3, \dots, 2m - 2, m - 2$, respectively.

Then all the vertices of G be colored black, thus $\{0, 1, m, 2m\}$ be a forcing set and $Z(C_n(1, m)) \leq 4$. □

The girth is the length of the shortest cycle of the graph. It is clearly that the girth of quartic circulant graphs $C_p(1, q)$ is 3 or 4. By [11, Theorem 2.4], if G is a graph of girth g at least 3 and minimum degree δ at least 2, then $Z(G) \geq (g - 2)(\delta - 2) + 2$. The next theorem give the exact value of the zero forcing number of $C_p(1, 3)$.

Theorem 3.6. *If $G = C_p(1, 3)$, then*

$$Z(C_p(1, 3)) = \begin{cases} 4, & p = 7, \\ 6, & p \geq 11. \end{cases}$$

Proof. By Theorem 3.5, we have $Z(C_7(1, 3)) = 4$.




Fig. 5. $C_7(1, 3)$ with a zero forcing set $\{0, 1, 4, 6\}$

We are going to prove that when $p \geq 11$, $Z(C_p(1, 3)) = 6$. By Theorem 3.1, $Z(C_p(1, 3)) \leq 2 \times 3 = 6$. In the following, we prove that $Z(C_p(1, 3)) \geq 6$. We claim that when $p \geq 11$, the girth of $C_p(1, 3)$ is 4, that is, $C_p(1, 3)$ is triangle-free. Otherwise, without loss of generality, we can assume that $C = (0, x, y, 0)$ is the 3-cycle containing the identity element 0 (the vertex-transitivity guarantees the existence of this 3-cycle). That is to say, x and y are adjacent. Since $N(0) = \{1, 3, p-3, p-1\}$, under the assumption that $p \geq 11$, $3, p-3, p-1$ are not adjacent. Thus $g \geq 4$. Also we can find a 4-cycle $(0, 1, p-2, p-3, 0)$. So when $p \geq 11$, the girth of $C_p(1, 3)$ equal to 4, and $Z(C_p(1, 3)) \geq (g-2)(\delta-2) + 2 = (4-2)(4-2) + 2 = 6$.

Hence we conclude that $Z(C_p(1, 3)) = 6$. \square

3.2. The zero forcing number of $C_p(1, 4)$

In this section, we characterize the exact values of the zero forcing number of the circulant graphs $C_p(1, 4)$. We first propose the following results.

Lemma 3.7. *Let G be a graph, u and v be the first and second vertices performing the forcing process with the property that $d(u, v) \geq 3$, then $|Z| \geq d(u) + d(v)$.*

Proof. Since u and v be the first and second vertices performing the forcing process and $d(u, v) \geq 3$, $|N(u) \cap N(v)| = 0$, then $|Z| \geq 1 + (d(u) - 1) + 1 + (d(v) - 1) = d(u) + d(v)$. \square

We will present the first three parts of distance partition $\{N_0(0), N_1(0), \dots, N_d(0)\}$ of $C_p(1, 4)$ for $p > 17$, see Figure 6.




Fig. 6. Distance partition of $C_p(1,4)$, where $p > 17$

According to the graphs definition, when $p > 17$, $N_0(0)$, $N_1(0)$, $N_2(0)$ are fixed. Note that when $p = 19$, $7 = p - 12$ and $12 = p - 7$, then $N_3(0)$ has 6 vertices instead of 8. But this does not affect the following proof, and the proof of $p = 19$ and $p > 19$ is the same. So in the following, we use the Figure 6 as the example graph.

Let Z be a zero forcing set for $C_p(1,4)$. Note that $C_p(1,4)$ is the vertex-transitive graph, with no loss of generality, we assume that 0 is the first black vertex which performs the forcing process. Then all but one of the neighbors of 0 must be contained in Z . Suppose that $4 \notin Z$, where $4 \in N_1(0)$. By applying the forcing process, $0 \rightarrow 4$.

Let v be the second black vertex which performs the forcing process. According to the symmetry of $C_p(1,4)$, we only discuss vertex v as one vertex of the first half of Figure 6, i.e., $\{1, 4, 2, 3, 5, 8, 6, 7, \dots\}$. In the following, we say that a set is black, which means that its all vertices are black.

Based on the distance partition, we can get the following lemma of the zero forcing number of $C_p(1,4)$.

Lemma 3.8. *Let $G = C_p(1,4)$ with $p > 17$, 0 be the first black vertex which performs the forcing process, v be the second black vertex which performs the forcing process and $v \in N_2(0)$, then $|Z| \geq 8$.*

Proof. By the symmetry of $C_p(1,4)$, we only discuss vertex v as one vertex of 2, 3, 5, 8.

(1) When $v = 2$.

In this case, $|N(v) \cap N_1(0)| = |\{1\}| = 1$, $|N(v) \cap N_2(0)| = |\{3, p-2\}| = 2$, $|N(v) \cap N_3(0)| = |\{6\}| = 1$. Since v be the second black vertex which performs the forcing process, v and its neighbors except one must be included in Z , that is, $|Z| \geq 7$, and $0, 1, 4, p-4, p-1, 2, 3, p-2, 6$ are black. Let $W_1 = \{0, 1, 4, p-4, p-1, 2, 3, p-2, 6\}$.

After the above forcing process, the black vertex $3, p-1$ have the unique neighbour $7, p-5$ not be colored black, respectively. Thus $3 \rightarrow 7, p-1 \rightarrow p-5$, and $W_1 \cup \{7, p-5\}$ is black, see Figure 7.




Fig. 7.

Let w be the next black vertex which performs the forcing process, w has the property that $|(N(w) \cup w) \cap (W_1 \cup \{7, p-5\})| \leq 3$, so we have at least one more vertex belong to Z , which infers that $|Z| \geq 7 + 1 = 8$.

When $v = 8$, $|N(v) \cap N_1(0)| = |\{4\}| = 1$, $|N(v) \cap N_3(0)| = |\{7, 9, 12\}| = 3$, Since v be the second black vertex which performs the forcing process, v and its neighbors except one must be included in Z , that is, $|Z| \geq 7$. Then we can get the result of $|Z| \geq 8$ by a similar analysis.

(2) When $v = 3$.

In this case, $|N(v) \cap N_1(0)| = |\{4, p-1\}| = 2$, $|N(v) \cap N_2(0)| = |\{2\}| = 1$, $|N(v) \cap N_3(0)| = |\{7\}| = 1$. Since v be the second black vertex which performs the forcing process, v and its neighbors except one must be contained in Z , that is, $|Z| \geq 6$, and $0, 1, 4, p-4, p-1, 2, 3, 7$ are black. Let $W_2 = \{0, 1, 4, p-4, p-1, 2, 3, 7\}$, see Figure 8.




Fig. 8.

Let w be the next black vertex which performs the forcing process. The following cases will be considered.

Case i. $w \in V \setminus \{1, 2, 4, p-1\}$.

In this case, w has the property that $|(N(w) \cup w) \cap W_2| \leq 2$, there are at least two more vertices belong to Z , which implies that $|Z| \geq 6 + 2 = 8$.

Case ii. $w \in \{1, 2, 4, p-1\}$.

When $w = 1$. Without loss of generality, let $5 \in Z$, then $1 \rightarrow p-3$. At this time, $|Z| \geq 6 + 1 = 7$. Then the black vertex 4 has the unique neighbour 8 not be colored black,

thus $4 \rightarrow 8$. Let u be the next black vertex which performs the forcing process, u has the property that $|(N(u) \cup u) \cap (W_2 \cup \{5, 8, p - 3\})| \leq 3$. So there is at least one more vertex contained in Z , and $|Z| \geq 7 + 1 = 8$.

When $w = 2$. Without loss of generality, let $6 \in Z$, then $2 \rightarrow p - 2$. At this time, $|Z| \geq 6 + 1 = 7$. Then the black vertex $p - 1$ has the unique neighbour $p - 5$ not be colored black, thus $p - 1 \rightarrow p - 5$. Let u be the next black vertex which performs the forcing process, u has the property that $|(N(u) \cup u) \cap (W_2 \cup \{6, p - 2, p - 5\})| \leq 3$, so at least one more vertex is added to Z , then $|Z| \geq 7 + 1 = 8$.

When $w = 4$. Let $5 \in Z$, then $4 \rightarrow 8$ and $|Z| \geq 6 + 1 = 7$. The black vertex 1 has the unique neighbour $p - 3$ not be colored black, thus $1 \rightarrow p - 3$. Let u be the next black vertex which performs the forcing process, u has the property that $|(N(u) \cup u) \cap (W_2 \cup \{5, 8, p - 3\})| \leq 3$, so we have at least one more vertex belonging to Z , which results in $|Z| \geq 7 + 1 = 8$.

When $w = p - 1$. Let $p - 2 \in Z$, then $p - 1 \rightarrow p - 5$ and $|Z| \geq 6 + 1 = 7$. The black vertex 2 has the unique neighbour 6 not be colored black, thus $2 \rightarrow 6$. Let u be the next black vertex which performs the forcing process, u has the property that $|(N(u) \cup u) \cap (W_2 \cup \{6, p - 2, p - 5\})| \leq 3$, then there is at least one more vertex that is included in Z , and $|Z| \geq 7 + 1 = 8$.

When $v = 5$, $|N(v) \cap N_1(0)| = |\{1, 4\}| = 2$, $|N(v) \cap N_3(0)| = |\{6, 9\}| = 2$. Since v be the second black vertex which performs the forcing process, v and its neighbors except one must be included in Z , that is, $|Z| \geq 6$. Then we can get the result of $|Z| \geq 8$ by a similar analysis. □

Lemma 3.9. *Let $G = C_p(1, 4)$, where $p > 17$, 0 be the first black vertex which performs the forcing process, v be the second black vertex which performs the forcing process and $v \in N_1(0)$, then $|Z| \geq 8$.*

Proof. By the symmetry of $C_p(1, 4)$, v can be regarded as 1 or 4.

When $v = 1$. In this case, $|N(v) \cap N_0(0)| = |\{0\}| = 1$, $|N(v) \cap N_2(0)| = |\{2, 5, p - 3\}| = 3$. Since v be the second black vertex which performs the forcing process, v and its neighbors except one must be contained in Z , that is, $|Z| \geq 4 + 2 = 6$, and $0, 1, 4, p - 4, 2, 5, p - 3$ are black. Let $W_3 = \{0, 1, 4, p - 4, 2, 5, p - 3\}$.




Fig. 9.

Let w be the next black vertex which performs the forcing process. The following cases will be considered.

Case i. $w \in V \setminus \{3, 4, p-2, p-4\}$.

In this case, $|(N(w) \cup w) \cap W_3| \leq 2$, so there are at least two extra vertices that are added to Z , we thus conclude that $|Z| \geq 6 + 2 = 8$.

Case ii. $w \in \{3, 4, p-2, p-4\}$.

When $w = 3$. Let $3 \in Z$, then $3 \rightarrow 7$. At this time, $|Z| \geq 6 + 1 = 7$. The black vertex 4 has the unique neighbour 8 not be colored black, thus $4 \rightarrow 8$. Let u be the next black vertex which performs the forcing process, u has the property that $|(N(u) \cup u) \cap (W_3 \cup \{3, 7, 8\})| \leq 3$, then there is at least one more vertex belonging to Z , and $|Z| \geq 7 + 1 = 8$.

When $w = 4$. Without loss of generality, let $3 \in Z$, then $4 \rightarrow 8$ and $|Z| \geq 6 + 1 = 7$. The black vertex 3 has the unique neighbour 7 not be colored black, thus $3 \rightarrow 7$. Let u be the next black vertex performs the forcing process, u satisfies the property that $|(N(u) \cup u) \cap (W_3 \cup \{3, 7, 8\})| \leq 3$, so at least one more vertex belongs to Z , which infers that $|Z| \geq 7 + 1 = 8$.

When $w = p-2$. Let $p-2 \in Z$, then $p-2 \rightarrow p-6$. At this time, $|Z| \geq 6 + 1 = 7$. Then the black vertex $p-3$ has the unique neighbour $p-7$ not be colored black, thus $p-3 \rightarrow p-7$. Let u be the next black vertex which performs the forcing process, we can find that $|(N(u) \cup u) \cap (W_3 \cup \{p-2, p-6, p-7\})| \leq 3$, so there is at least one extra vertex that is added to Z , we get that $|Z| \geq 7 + 1 = 8$.

When $w = p-4$. Without loss of generality, let $p-5 \in Z$, then $p-4 \rightarrow p-8$ and $|Z| \geq 6 + 1 = 7$. Let u be the next black vertex which performs the forcing process, u has the property that $|(N(u) \cup u) \cap (W_3 \cup \{p-5, p-8\})| \leq 3$, so we have at least one more vertex belonging to Z , and $|Z| \geq 7 + 1 = 8$.

When $v = 4$, $|N(v) \cap N_0(0)| = |\{0\}| = 1$, $|N(v) \cap N_2(0)| = |\{3, 5, 8\}| = 3$. Since v be the second black vertex which performs the forcing process, v and its neighbors except one must be included in Z , that is, $|Z| \geq 6$. Then we can obtain the result of $|Z| \geq 8$ by a similar analysis. \square

With the above preparations, in the following, we can completely characterize the zero forcing number of $C_p(1, 4)$.

Theorem 3.10. *If $G = C_p(1, 4)$, then*

$$Z(C_p(1, 4)) = \begin{cases} 6, & p = 11 \text{ or } 13, \\ 8, & p \geq 17. \end{cases}$$

Proof. First, we prove that when $p = 11$, $Z(C_p(1, 4)) = 6$. The circulant graph $C_{11}(1, 4)$ as follows:




Fig. 10. The circulant graph $C_{11}(1,4)$

It is easy to check that $\{0, 1, 2, 7, 8, 10\}$ is a zero forcing set for $C_{11}(1,4)$, that is $Z(C_{11}(1,4)) \leq 6$. And $Z(C_{11}(1,4)) \geq (g-2)(\delta-2) + 2 \geq 6$, hence we conclude that $Z(C_{11}(1,4)) = 6$. The forcing processes as follows: $0 \rightarrow 4$, $1 \rightarrow 5$, $4 \rightarrow 3$, $7 \rightarrow 6$ and $8 \rightarrow 9$. Similarly, $Z(C_{13}(1,4)) = 6$.

Now we prove that when $p \geq 17$, $Z(C_p(1,4)) = 8$. By Theorem 3.1, $Z(C_p(1,4)) \leq 8$. In the following, we prove that $Z(C_p(1,4)) \geq 8$, and hence we conclude that $Z(C_p(1,4)) = 8$.

Firstly, we show that $Z(C_p(1,4)) \geq 8$ for $p > 17$. Let 0 be the first black vertex which performs the forcing process, and v be the second black vertex which performs the forcing process. The following cases will be considered.

Case 1. $v \in N_i(0)$, $i \geq 3$.

Since $C_p(1,4)$ is 4-regular, and $d(0, v) \geq 3$, by Lemma 3.7, $|Z| \geq d(0) + d(v) = 4 + 4 = 8$.

Case 2. $v \in N_2(0)$.

By Lemma 3.3, $|Z| \geq 8$.

Case 3. $v \in N_1(0)$.

By Lemma 3.4, $|Z| \geq 8$.

Note that when $p > 17$, 8 and $p-8$ are not adjacent, 8 and $p-5$ are not adjacent, 5 and $p-8$ are not adjacent. On the contrary, when $p = 17$, they are adjacent. Similarly, based on the distance partition, we can prove that when $p = 17$, $Z(C_p(1,4)) \geq 8$.

Therefore, when $p \geq 17$, $Z(C_p(1,4)) = 8$. □

3.3. The zero forcing number of $C_p(1,5)$

Using the similar way of showing $Z(C_p(1,4))$, we can determine the exact value of $Z(C_p(1,5))$, but the process of the proof is long and tedious. Hence, we omit it here. But we present the outline of the proof as follows: $Z(C_{11}(1,5))$ can be easily obtained through Theorem 3.5; the proof of $Z(C_{13}(1,5))$ and $Z(C_{17}(1,5))$ is similar to $Z(C_{11}(1,4))$; when $p \geq 19$, the proof of $Z(C_p(1,5))$ adopts the method of distance partition, and the proof process can be referred to $C_p(1,4)$, $p \geq 17$.

Theorem 3.11. *If $G = C_p(1,5)$ with the prime p , then $Z(C_p(1,5)) = 10$ for $p \geq 29$. In particular, for $p < 29$, $Z(C_p(1,5))$ is presented in the following table.*

$C_p(1, q)$	diameter	$ Z $	Z
$C_{11}(1, 5)$	3	4	$\{0, 1, 5, 10\}$
$C_{13}(1, 5)$	2	6	$\{0, 1, 2, 7, 8, 12\}$
$C_{17}(1, 5)$	3	6	$\{0, 1, 6, 7, 12, 16\}$
$C_{19}(1, 5)$	3	8	$\{0, 1, 2, 4, 14, 15, 17, 18\}$
$C_{23}(1, 5)$	3	8	$\{0, 1, 5, 6, 7, 12, 13, 22\}$

Table 1.

4. Concluding Remarks

In the paper, we have explored the zero forcing number of quartic circulant graphs $C_p(s, t)$. Firstly, we focus on the isomorphism of quartic circulant graphs, and conclude that $C_p(s, t) \cong C_p(1, q)$ for some $q \in Z_p^*$. In addition, for $q = 2, 3, 4, 5$, we obtain the zero forcing number of $C_p(1, q)$. When $|p - q|$ is small, we can use the method of distance partition to get $Z(G)$. However, for $|p - q|$ is large, analyzing $Z(G)$ with distance partition is very complicated. In this case, $Z(G)$ is still unsolved. For further study, it would be an interesting problem to verify when $|p - q|$ is sufficiently large, $Z(C_p(1, q)) = 2q$.

References

- [1] AIM Minimum Rank - Special Graphs Work Group (F. Barioli, W. Barrett, S. Butler, S. M. Cioabă, D. Cvetković, S. M. Fallat, C. Godsil, W. Haemers, L. Hogben, R. Mikkelsen, S. Narayan, O. Pryporova, I. Sciriha, W. So, D. Stevanović, H. van der Holst, K. Vander Meulen, A. Wangsness). Zero forcing sets and the minimum rank of graphs. *Linear Algebra and its Applications*, 428:1628–1648, 2008. <https://doi.org/10.1016/j.laa.2007.10.009>.
- [2] D. Amos, Y. Caro, R. Davila, and R. Pepper. Upper bounds on the k-forcing number of a graph. *Discrete Applied Mathematics*, 181:1–10, 2015. <https://doi.org/10.1016/j.dam.2014.08.029>.
- [3] F. Barioli, W. Barrett, S. M. Fallat, H. T. Hall, L. Hogben, B. Shader, H. van der Holst, and P. van der Driessche. Parameters related to tree-width, zero forcing, and maximum nullity of a graph. *Journal of Graph Theory*, 72(2):146–177, 2013. <https://doi.org/10.1002/jgt.21637>.
- [4] K. F. Benson, D. Ferrero, M. Flagg, V. Furst, L. Hogben, V. Vasilevska, and B. Wissman. Zero forcing and power domination for graph products. *Australas. Combin.*, 70, 2018.
- [5] J. A. Bondy and U. S. R. Murty. *Graph theory*. Springer Publishing Company, Incorporated, 2008.
- [6] D. Burgarth, V. Giovannetti, L. Hogben, S. Severini, and M. Young. Logic circuits from zero forcing. *Natural Computing*, 14:485–490, 2015. <https://doi.org/10.1007/s11047-014-9438-5>.

-
- [7] R. Davila, T. Kalinowski, and S. Stephen. A lower bound on the zero forcing number. *Discrete Applied Mathematics*, 250:363–367, 2018. <https://doi.org/10.1016/j.dam.2018.04.015>.
- [8] L. Duong, B. K. Kroschel, M. Riddell, K. N. Vander Meulen, and A. Tuyl. Maximum nullity and zero forcing of circulant graphs. *Special Matrices*, 8(1):221–234, 2020. <https://doi.org/10.1515/spma-2020-0106>.
- [9] C. J. Edholm, L. Hogben, M. Huynh, J. LaGrange, and D. D. Row. Vertex and edge spread of zero forcing number, maximum nullity, and minimum rank of a graph. *Linear Algebra and its Applications*, 436(12):4352–4372, 2012. <https://doi.org/10.1016/j.laa.2010.10.015>.
- [10] L. Eroh, C. X. Kang, and E. Yi. A comparison between the metric dimension and zero forcing number of trees and unicyclic graphs. *Acta Mathematica Sinica. English Series*, 33:731–747, 2017. <https://doi.org/10.1007/s10114-017-4699-4>.
- [11] M. Fürst and D. Rautenbach. A short proof for a lower bound on the zero forcing number. *Discussiones Mathematicae Graph Theory*, 1(2).
- [12] M. Gentner, L. D. Penso, D. Rautenbach, and U. S. Souza. Extremal values and bounds for the zero forcing number. *Discrete Applied Mathematics*, 214:196–200, 2016. <https://doi.org/10.1016/j.dam.2016.06.004>.
- [13] M. Gentner and D. Rautenbach. Some bounds on the zero forcing number of a graph. *Discrete Applied Mathematics*, 236:203–213, 2018. <https://doi.org/10.1016/j.dam.2017.11.015>.
- [14] T. W. Haynes, S. M. Hedetniemi, S. T. Hedetniemi, and M. A. Henning. Domination in graphs applied to electric power networks. *SIAM Journal on Discrete Mathematics*, 15(4):519–529, 2002. <https://doi.org/10.1137/S0895480100375831>.
- [15] C. Heuberger. On planarity and colorability of circulant graphs. *Discrete Mathematics*, 268(1–3):153–169, 2003. [https://doi.org/10.1016/S0012-365X\(02\)00685-4](https://doi.org/10.1016/S0012-365X(02)00685-4).
- [16] T. Kalinowski, N. Kamcev, and B. Sudakov. The zero forcing number of graphs. *SIAM Journal on Discrete Mathematics*, 33(1):95–115, 2019. <https://doi.org/10.1137/17M1133051>.
- [17] S. Lang. *Algebra*. Springer Science and Business Media, 2012.
- [18] L. Loudiki, M. Kchkech, and E. H. Essaky. A new approach for computing the distance and the diameter in circulant graphs. arXiv preprint, 2022. <https://doi.org/10.48550/arXiv.2210.11116>. (Cornell Univ., Ithaca, NY).
- [19] L. Lu, B. Wu, and Z. Tang. Proof of a conjecture on the zero forcing number of a graph. *Discrete Applied Mathematics*, 213:233–237, 2016. <https://doi.org/10.1016/j.dam.2016.05.009>.
- [20] S. A. Meyer. Zero forcing sets and bipartite circulants. *Linear Algebra and its Applications*, 436(4):888–900, 2012. <https://doi.org/10.1016/j.laa.2011.09.022>.
- [21] M. Muzychuk. \acute{A} dám’s conjecture is true in the square-free case. *Journal of Combinatorial Theory, Series A*, 72(1):118–134, 1995. [https://doi.org/10.1016/0097-3165\(95\)90031-4](https://doi.org/10.1016/0097-3165(95)90031-4).

-
- [22] N. Obradović, J. Peters, and G. Ružić. Efficient domination in circulant graphs with two chord lengths. *Information Processing Letters*, 102(6):253–258, 2007. <https://doi.org/10.1016/j.ipl.2007.02.004>.
- [23] M. Riddell. *The zero forcing number of circulant graphs*. msc, McMaster University, 2017.
- [24] M. Trefois and J. C. Delvenne. Zero forcing sets, constrained matchings and minimum rank. *Linear and Multilinear Algebra*, 484:199–218, 2013.
- [25] X. Wang, D. Wong, and Y. Zhang. Zero forcing number of a graph in terms of the number of pendant vertices. *Linear and Multilinear Algebra*, 68(7):1424–1433, 2020. <https://doi.org/10.1080/03081087.2018.1545829>.
- [26] D. Young. *Techniques for determining equality of the maximum nullity and the zero forcing number of a graph*. PhD thesis, Iowa State University, 2019.
- [27] Y. Zhang, X. Yong, and M. J. Golin. The number of spanning trees in circulant graphs. *Discrete Mathematics*, 223(1–3):337–350, 2000.