

An Extremal Problem On Potentially $K_m - C_4$ -graphic Sequences *

Chunhui Lai

Department of Mathematics

Zhangzhou Teachers College, Zhangzhou

Fujian 363000, P. R. of CHINA.

e-mail: zjlaichu@public.zzptt.fj.cn

Abstract

A sequence S is potentially $K_m - C_4$ -graphical if it has a realization containing a $K_m - C_4$ as a subgraph. Let $\sigma(K_m - C_4, n)$ denote the smallest degree sum such that every n -term graphical sequence S with $\sigma(S) \geq \sigma(K_m - C_4, n)$ is potentially $K_m - C_4$ -graphical. In this paper, we prove that $\sigma(K_m - C_4, n) \geq (2m - 6)n - (m - 3)(m - 2) + 2$, for $n \geq m \geq 4$. We conjecture that equality holds for $n \geq m \geq 4$. We prove that this conjecture is true for $m = 5$.

Key words: graph; degree sequence; potentially $K_m - C_4$ -graphic sequence

AMS Subject Classifications: 05C07, 05C35

1 Introduction

If $S = (d_1, d_2, \dots, d_n)$ is a sequence of non-negative integers, then it is called graphical if there is a simple graph G of order n , whose degree sequence $(d(v_1), d(v_2), \dots, d(v_n))$ is precisely S . If G is such a graph then G is said to realize S or be a realization of S . A graphical sequence S is potentially H -graphical if there is a realization of S containing H as a subgraph, while S is forcibly H -graphical if every realization of S contains H as a subgraph. Let $\sigma(S) = d(v_1) + d(v_2) + \dots + d(v_n)$, and $[x]$ denote the largest integer less than or equal to x . We denote $G+H$ as the graph with $V(G+H) = V(G) \cup V(H)$ and $E(G+H) = E(G) \cup E(H) \cup \{xy : x \in V(G), y \in V(H)\}$. Let K_k ,

*Project Supported by NNSF of China(10271105), NSF of Fujian(Z0511034), Science and Technology Project of Fujian, Fujian Provincial Training Foundation for "Bai-Quan-Wan Talents Engineering", Project of Fujian Education Department and Project of Zhangzhou Teachers College.

and C_k denote a complete graph on k vertices, and a cycle on k vertices, respectively. Let $K_m - C_4$ be the graph obtained from K_m by removing four edges of a 4 cycle C_4 .

Given a graph H , what is the maximum number of edges of a graph with n vertices not containing H as a subgraph? This number is denoted $ex(n, H)$, and is known as the Turán number. This problem was proposed for $H = C_4$ by Erdős [2] in 1938 and in general by Turán [10]. In terms of graphic sequences, the number $2ex(n, H) + 2$ is the minimum even integer l such that every n -term graphical sequence S with $\sigma(S) \geq l$ is forcibly H -graphical. Here we consider the following variant: determine the minimum even integer l such that every n -term graphical sequence S with $\sigma(S) \geq l$ is potentially H -graphical. We denote this minimum l by $\sigma(H, n)$. Erdős, Jacobson and Lehel [3] showed that $\sigma(K_k, n) \geq (k-2)(2n-k+1)+2$ and conjectured that equality holds. They proved that if S does not contain zero terms, this conjecture is true for $k = 3$, $n \geq 6$. The conjecture is confirmed in [4],[6],[7],[8] and [9].

Gould, Jacobson and Lehel [4] also proved that $\sigma(pK_2, n) = (p-1)(2n-2) + 2$ for $p \geq 2$; $\sigma(C_4, n) = 2\lfloor \frac{3n-1}{2} \rfloor$ for $n \geq 4$. Lai [5] proved that $\sigma(K_4 - e, n) = 2\lfloor \frac{3n-1}{2} \rfloor$ for $n \geq 7$. In this paper, we prove that $\sigma(K_m - C_4, n) \geq (2m-6)n - (m-3)(m-2) + 2$, for $n \geq m \geq 4$. We conjectured that equality holds for $n \geq m \geq 4$. We prove that this conjecture is true for $m = 5$.

2 Main results.

Theorem 1. $\sigma(K_m - C_4, n) \geq (2m-6)n - (m-3)(m-2) + 2$, for $n \geq m \geq 4$.

Proof. Let

$$H = K_{m-3} + \overline{K_{n-m+3}}$$

Then H is a uniquely realization of $((n-1)^{m-3}, (m-3)^{n-m+3})$ and H clearly does not contain $K_m - C_4$. Thus

$$\sigma(K_m - C_4, n) \geq (m-3)(n-1) + (m-3)(n-m+3) + 2 = (2m-6)n - (m-3)(m-2) + 2.$$

Theorem 2. For $n \geq 5$, $\sigma(K_5 - C_4, n) = 4n - 4$.

Proof. By theorem 1, for $n \geq 5$, $\sigma(K_5 - C_4, n) \geq 4n - 4$. We need to show that if S is an n -term graphical sequence with $\sigma(S) \geq 4n - 4$, then there is a realization of S containing a $K_5 - C_4$. Let $d_1 \geq d_2 \geq \dots \geq d_n$, and let G be a realization of S .

Case: $n = 5$, if a graph has size $q \geq 8$, then clearly it contains a $K_5 - C_4$, so that $\sigma(K_5 - C_4, 5) \leq 4n - 4$.

Case: $n = 6$. If $\sigma(S) = 20$, we first consider $d_6 \leq 2$. Let S' be the degree sequence of $G - v_6$, so $\sigma(S') \geq 20 - 2 \times 2 = 16$. Then, by induction S' has a

realization containing a $K_5 - C_4$. Therefore S has a realization containing a $K_5 - C_4$. Now we consider $d_6 \geq 3$. It is easy to see that $S = (5^1, 3^5)$ or $S = (4^2, 3^4)$. Obviously, each is potentially $K_5 - C_4$ -graphic. Next, if $\sigma(S) = 22$ then it must be that $d_6 \leq 3$. Let S' be the degree sequence of $G - v_6$, so $\sigma(S') \geq 22 - 3 \times 2 = 16$. Then S' has a realization containing a $K_5 - C_4$. Therefore S has a realization containing a $K_5 - C_4$. Finally, suppose that $\sigma(S) \geq 24$. We first consider $d_6 \leq 4$. Let S' be the degree sequence of $G - v_6$, so $\sigma(S') \geq 24 - 2 \times 4 = 16$. Then S' has a realization containing a $K_5 - C_4$. Therefore S has a realization containing a $K_5 - C_4$. Now we consider $d_6 \geq 5$. It is easy to see that $S = (5^6)$. Obviously, (5^6) is potentially $K_5 - C_4$ -graphic.

Case: $n = 7$. First we assume that $\sigma(S) = 24$. Suppose $d_7 \leq 2$ and let S' be the degree sequence of $G - v_7$, so $\sigma(S') \geq 24 - 2 \times 2 = 20$. Then S' has a realization containing a $K_5 - C_4$. Therefore S has a realization containing a $K_5 - C_4$. Now we assume that $d_7 \geq 3$. It is easy to see that S is one of $(6^1, 3^6)$, $(5^1, 4^1, 3^5)$, or $(4^3, 3^4)$. Obviously, all of them are potentially $K_5 - C_4$ -graphic. Next, if $\sigma(S) = 26$, It is easy to see that $d_7 \leq 3$. Let S' be the degree sequence of $G - v_7$, so $\sigma(S') \geq 26 - 3 \times 2 = 20$. Then S' has a realization containing a $K_5 - C_4$. Therefore S has a realization containing a $K_5 - C_4$. Finally, suppose that $\sigma \geq 28$. If $d_7 \leq 4$. Let S' be the degree sequence of $G - v_7$, so $\sigma(S') \geq 28 - 2 \times 4 = 20$. Then S' has a realization containing a $K_5 - C_4$. Therefore S has a realization containing a $K_5 - C_4$. Now we consider $d_7 \geq 5$. It is easy to see that $\sigma(S) > 5 \times 7 = 35$. Clearly, $d_7 \leq 6$. Let S' be the degree sequence of $G - v_7$, so $\sigma(S') > 35 - 6 \times 2 = 23$. Then, by induction S' has a realization containing a $K_5 - C_4$. Therefore S has a realization containing a $K_5 - C_4$.

We proceed by induction on n . Take $n \geq 8$ and make the inductive assumption that for $7 \leq t < n$, whenever S_1 is a t -term graphical sequence such that

$$\sigma(S_1) \geq 4t - 4$$

then S_1 has a realization containing a $K_5 - C_4$. Let S be an n -term graphical sequence with $\sigma(S) \geq 4n - 4$. If $d_n \leq 2$, let S' be the degree sequence of $G - v_n$. Then $\sigma(S') \geq 4n - 4 - 2 \times 2 = 4(n - 1) - 4$. By induction, S' has a realization containing a $K_5 - C_4$. Therefore S has a realization containing a $K_5 - C_4$. Hence, we may assume that $d_n \geq 3$. By Proposition 2 and Theorem 4 of [4] (or Theorem 3.3 of [6]) S has a realization containing a K_4 . By Lemma 1 of [4], there is a realization G of S with v_1, v_2, v_3, v_4 , the four vertices of highest degree containing a K_4 . If $d(v_2) = 3$, then $4n - 4 \leq \sigma(S) \leq n - 1 + 3(n - 1) = 4n - 4$. Hence, $S = ((n - 1)^1, 3^{n-1})$. Obviously, $((n - 1)^1, 3^{n-1})$ is potentially $K_5 - C_4$ -graphic. Therefore, we may assume that $d(v_2) \geq 4$. Let v_1 be adjacent to v_2, v_3, v_4, y_1 . If y_1 is adjacent to one of v_2, v_3, v_4 , then G contains a $K_5 - C_4$. Hence, we may assume that

y_1 is not adjacent to v_2, v_3, v_4 . Let v_2 be adjacent to v_1, v_3, v_4, y_2 . If y_2 is adjacent to one of v_1, v_3, v_4 , then G contains a $K_5 - C_4$. Hence, we may assume that y_2 is not adjacent to v_1, v_3, v_4 . Since $d(y_1) \geq d_n \geq 3$, there is a new vertex y_3 , such that $y_1 y_3 \in E(G)$.

If $y_3 v_1 \in E(G)$, then G contains a $K_5 - C_4$. Hence, we may assume that $y_3 v_1 \notin E(G)$. Then the edge interchange that removes the edges $y_1 y_3, v_1 v_4$ and $v_2 y_2$ and inserts the non-edges $y_1 v_2, y_3 v_1$ and $y_2 v_4$ produces a realization G' of S containing a $K_5 - C_4$ where $K_5 - C_4$ with five vertices v_1, v_2, v_3, v_4, y_1 .

This finishes the inductive step, and thus Theorem 2 is established.

We make the following conjecture:

Conjecture.

$$\sigma(K_m - C_4, n) = (2m - 6)n - (m - 3)(m - 2) + 2$$

for $n \geq m \geq 4$.

This conjecture is true for $m = 4$, by Theorem 5 of [4], and for $m = 5$, by the above Theorem.

Acknowledgment

The author thanks the referees for many helpful comments.

References

- [1] B. Bollobás, *Extremal Graph Theory*, Academic Press, London, 1978.
- [2] P. Erdős, On sequences of integers no one of which divides the product of two others and some related problems, *Izv. Naustno-Issl. Mat. i Meh. Tomsk* 2(1938), 74-82.
- [3] P.Erdős, M.S. Jacobson and J. Lehel, Graphs realizing the same degree sequences and their respective clique numbers, in *Graph Theory, Combinatorics and Application*, Vol. 1(Y. Alavi et al., eds.), John Wiley and Sons, Inc., New York, 1991, 439-449.
- [4] R.J. Gould, M.S. Jacobson and J. Lehel, Potentially G -graphic degree sequences, in *Combinatorics, Graph Theory and Algorithms*, Vol. 2 (Y. Alavi et al., eds.), New Issues Press, Kalamazoo, MI, 1999, 451-460.
- [5] Lai Chunhui, A note on potentially $K_4 - e$ graphical sequences, *Australasian J. of Combinatorics* 24(2001), 123-127.

- [6] Li Jiong-Sheng and Song Zi-Xia, An extremal problem on the potentially P_k -graphic sequences, *Discrete Math*, (212)2000, 223-231.
- [7] Li Jiong-Sheng and Song Zi-Xia, The smallest degree sum that yields potentially P_k -graphical sequences, *J. Graph Theory*,29(1998), 63-72.
- [8] Li Jiong-sheng and Song Zi-Xia, On the potentially P_k -graphic sequences, *Discrete Math.* 195(1999), 255-262.
- [9] Li Jiong-sheng, Song Zi-Xia and Luo Rong, The Erdős-Jacobson-Lehel conjecture on potentially P_k -graphic sequence is true, *Science in China(Series A)*, 41(5)(1998), 510-520.
- [10] P. Turán, On an extremal problem in graph theory, *Mat. Fiz. Lapok* 48(1941), 436-452.