

A Note On Equitable Total Coloring of Middle and Total Graph of Some Graphs

¹G. Jayaraman and ²D. Muthuramakrishnan

^{1,2}Department of Mathematics,

¹Vels Institute of Science Technology and
Advanced Studies, Chennai, India

² National College(Autonomous), Trichy, India
jayaram07maths@gmail.com

Abstract

Among the various coloring of graphs, the concept of equitable total coloring of graph G is the coloring of all its vertices and edges in which the number of elements in any two color classes differ by at most one. The minimum number of colors required is called its equitable total chromatic number. In this paper, we obtained an equitable total chromatic number of middle graph of path, middle graph of cycle, total graph of path and total graph of cycle.

Keywords: Middle graph, Total graph, Equitable total coloring and Equitable total chromatic number.

1 Introduction

All the graphs considered here simple, finite and undirected graph. Let $G = (V(G), E(G))$ be a graph with vertex set $V(G)$ and edge set $E(G)$, respectively. A coloring of a graph G is an assignment of colors to the vertices or edges or both. A vertex-coloring(edge-coloring) is called proper coloring if no two vertices(edges) receive the same color. So many different proper colorings are available in graph theory such as a-coloring, b-coloring, acyclic coloring, star coloring, list coloring, harmonious coloring, total coloring, equitable total coloring. In the present work we focused an equitable

total coloring of graph. The concept of total coloring was independently introduced by Behzad [1] and Vizing [10]. A total coloring of a graph G is an assignment of colors to both the vertices and edges of G , such that no two adjacent or incident vertices and edges of G are receive the same color. They both conjectured that for any graph G the following condition holds: $\Delta(G)+1 \leq \chi''(G) \leq \Delta(G)+2$, where $\Delta(G)$ is the maximum degree of G . We observe that $\Delta(G)+1$ is the suitable lower bound. In general, an equitable total coloring problem is comparatively very difficult than the total coloring problem. In 1994, Fu[4] first introduce the concept of equitable total coloring and the equitable total chromatic number of a graph. A total coloring of a graph G is said to be equitable, if the number of elements(vertices and edges) in any two color classes differ by at most one, for which the required minimum number of colors is called the equitable total chromatic number and it is denoted by $\chi_{et}(G)$. Gong Kun et.al[3] proved some results on the equitable total chromatic number of $W_n \vee K_n$, $F_m \vee K_n$ and $S_m \vee K_n$. In 2012, Ma Gang and Ma Ming[6] proved some results concerning the equitable total chromatic number of $P_m \vee S_n$, $P_m \vee F_n$ and $P_m \vee W_n$. Tong et.al[8] proved that the equitable total chromatic number of $C_m \square C_n$. Girija et.al[2] proved that equitable total chromatic number of Double star graph and fan graph. Gang et.al[7] proved that on the equitable total coloring of multiple join graph. Zhang Zhong-fu[12] proved that on the equitable total coloring of some join graphs. Veninstine vivik et.al[9] proved an algorithmic approach to equitable total chromatic number of wheel graph, Gear graph, Helm graph and sunlet graph.

2 Preliminaries

Definition 2.1. *The middle graph of a graph G , denoted by $M(G)$ is define as follows, the vertex set of $M(G)$ is $V(G) \cup E(G)$. Two vertices x, y in the vertex set of $M(G)$ are adjacent in $M(G)$ in case one of the following condition holds: (i) x, y are in $E(G)$ and x, y is adjacent in G . (ii) x is in $V(G)$, y is in $E(G)$ and x, y are incident in G . The total graph of a graph G , denoted by $T(G)$ is define as, the vertex set of $T(G)$ is $V(G) \cup E(G)$. Two vertices x, y in the vertex set of $T(G)$ are adjacent in $T(G)$ in case one of the following condition holds: (i) x, y are in $V(G)$ and x is adjacent to y in G . (ii) x, y are in $E(G)$ and x, y is adjacent in G (iii) x is in $V(G)$, y is in $E(G)$ and x, y are incident in G . For a simple graph G , let f be a proper k - total coloring of G , $||T_i| - |T_j|| \leq 1$, for $i, j = 1, 2, \dots, k$. The partition $\{T_i\} = \{V_i \cup E_i : 1 \leq i \leq k\}$ is called a k - equitable total coloring and $\chi_{et}(G) = \min \{k / k\text{-equitable total coloring of } G\}$ is called the equitable total chromatic number of G , where for all $x \in T_i = V_i \cup E_i$,*

$f(x) = i,$ for $i = 1, 2, \dots, k.$

Conjecture 2.4([4]) For any simple graph $G(V, E), \chi_{et}(G) \leq \Delta(G) + 2.$

Conjecture 2.5([13]) For any simple graph $G(V, E),$

$$\chi_{et}(G) \geq \chi''(G) \geq \Delta(G) + 1.$$

Conjecture 2.6([11]) For every graph G, G has an equitable total k -coloring for each $k \geq \max\{\chi''(G), \Delta(G) + 2\}$

proposition 2.7([5]) Any four regular multi-graph can be total colored with 6 colors

proposition 2.8([7])For complete graph K_n with order $n,$

$$\chi_{et}(K_n) = \begin{cases} n, & \text{if } n \equiv 1 \pmod{2} \\ n + 1, & \text{if } n \equiv 0 \pmod{2} \end{cases}$$

In this paper, we obtained an equitable total chromatic number of middle graph of path, Middle graph of cycle, Total graph of path and Total graph of cycle.

3 Main Results

Theorem 3.1. For any positive integer $n \geq 3, \chi_{et}(M(P_n)) = 5.$

Proof. Let $V(P_n) = \{v_i : 1 \leq i \leq n\}$ and $E(P_n) = \{e_i : 1 \leq i \leq n - 1\},$ where $\{e_i : 1 \leq i \leq n - 1\}$ be the edges $v_i v_{i+1} (1 \leq i \leq n - 1).$ By the definition of middle graph, each edge $\{e_i : 1 \leq i \leq n - 1\}$ is subdivided by the vertices $\{u_i : 1 \leq i \leq n - 1\}.$ In $M(P_n),$ the vertex set and the edge set is given by $V(M(P_n)) = \{v_i : 1 \leq i \leq n\} \cup \{u_i : 1 \leq i \leq n - 1\}$ and $E(M(P_n)) = \{e_i : 1 \leq i \leq n - 1\} \cup \{e'_i : 1 \leq i \leq n - 1\} \cup \{e''_i : 1 \leq i \leq n - 2\},$ where $e_i (1 \leq i \leq n)$ is an edge $v_i v_{i+1} (1 \leq i \leq n - 1), e'_i (1 \leq i \leq n - 1)$ is an edge $u_i v_{i+1} (1 \leq i \leq n - 1)$ and $e''_i (1 \leq i \leq n - 2)$ is an edge $u_i u_{i+1} (1 \leq i \leq n - 2).$

Define an equitable total coloring $f,$ such that $f : S \rightarrow C,$ where $S = V(M(P_n)) \cup E(M(P_n))$ and $C = \{1, 2, 3, 4, 5\}.$ Now we assign an equitable total coloring to these vertices and edges as follows.

$$f(v_i) = \begin{cases} 1, & \text{if } i \equiv 1 \pmod{3} \\ 2, & \text{if } i \equiv 2 \pmod{3} \\ 3, & \text{if } i \equiv 0 \pmod{3} \end{cases} \text{ for } 1 \leq i \leq n$$

$$f(u_i) = \begin{cases} 3, & \text{if } i \equiv 1 \pmod{3} \\ 1, & \text{if } i \equiv 2 \pmod{3} \\ 2, & \text{if } i \equiv 0 \pmod{3} \end{cases} \quad \text{for } 1 \leq i \leq n-1$$

$$f(e_i) = 4, \quad \text{for } 1 \leq i \leq n-1$$

$$f(e_i') = 5, \quad \text{for } 1 \leq i \leq n-1$$

$$f(e_i'') = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{3} \\ 3, & \text{if } i \equiv 2 \pmod{3} \\ 1, & \text{if } i \equiv 0 \pmod{3} \end{cases} \quad \text{for } 1 \leq i \leq n-2$$

Based on the above procedure of coloring, the graph $M(P_n)$ is equitable total colored with 5 colors, such that the color classes are $T(M(P_n)) = \{T_1, T_2, T_3, T_4, T_5\}$. We observe that these color classes T_1, T_2, T_3, T_4, T_5 are independent sets of $M(P_n)$ and it satisfies the inequality $||T_i| - |T_j|| \leq 1$, for $i \neq j$. Hence $\chi_{et}(M(P_n)) \leq 5$. Further, since $\Delta = 4$, we have $\chi_{et}(M(P_n)) \geq \chi''(M(P_n)) \geq \Delta + 1 \geq 4 + 1 \geq 5$. Therefore $\chi_{et}(M(P_n)) = 5$. \square

Theorem 3.2. For any positive integer $n \geq 4$, $\chi_{et}(M(C_n)) = 5$.

Proof. Let $V(C_n) = \{v_i : 1 \leq i \leq n\}$ and $E(C_n) = \{e_i : 1 \leq i \leq n\}$, where $\{e_i : 1 \leq i \leq n-1\}$ be the edges $v_i v_{i+1}, i+1$ taken modulo $n(1 \leq i \leq n)$. By the definition of middle graph, each edge $\{e_i : 1 \leq i \leq n\}$ is subdivided by the vertices $\{u_i : 1 \leq i \leq n\}$. In $M(C_n)$, the vertex set and the edge set is given by $V(M(C_n)) = \{v_i : 1 \leq i \leq n\} \cup \{u_i : 1 \leq i \leq n\}$ and $E(M(C_n)) = \{e_i : 1 \leq i \leq n\} \cup \{e_i' : 1 \leq i \leq n\} \cup \{e_i'' : 1 \leq i \leq n\}$, where $e_i(1 \leq i \leq n)$ be an edge $v_i u_i(1 \leq i \leq n)$, $e_i'(1 \leq i \leq n-1)$ be an edge $u_i v_{i+1}, i+1$ taken modulo $n(1 \leq i \leq n)$ and $e_i''(1 \leq i \leq n)$ be an edge $u_i u_{i+1}, i+1$ taken modulo $n(1 \leq i \leq n)$.

We define an equitable total coloring f , such that $f : S \rightarrow C$, where $S = V(M(C_n)) \cup E(M(C_n))$ and $C = \{1, 2, 3, 4, 5\}$. The equitable total coloring pattern is as follows. we consider the following two cases

Case(i): when n is even

$$f(v_i) = \begin{cases} 4, & \text{if } i \equiv 1 \pmod{2} \\ 3, & \text{if } i \equiv 0 \pmod{2} \end{cases} \quad \text{for } 1 \leq i \leq n$$

$$f(u_i) = \begin{cases} 1, & \text{if } i \equiv 1 \pmod{2} \\ 2, & \text{if } i \equiv 0 \pmod{2} \end{cases} \quad \text{for } 1 \leq i \leq n$$

$$f(e_i) = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{2} \\ 1, & \text{if } i \equiv 0 \pmod{2} \end{cases} \quad \text{for } 1 \leq i \leq n$$

$$f(e_i') = 5, \quad \text{for } 1 \leq i \leq n$$

$$f(e_i'') = \begin{cases} 3, & \text{if } i \equiv 1 \pmod{2} \\ 4, & \text{if } i \equiv 0 \pmod{2} \end{cases} \quad \text{for } 1 \leq i \leq n$$

Case(ii): when n is odd




Figure 1: Equitable Total Coloring of middle graph of Cycle $M(C_6)$

$$f(v_1) = 2$$

$$f(v_i) = \begin{cases} 3, & \text{if } i \equiv 0 \pmod{2} \\ 4, & \text{if } i \equiv 1 \pmod{2} \end{cases} \quad \text{for } 2 \leq i \leq n$$

$$f(u_i) = \begin{cases} 1, & \text{if } i \equiv 1 \pmod{2} \\ 2, & \text{if } i \equiv 0 \pmod{2} \end{cases} \quad \text{for } 1 \leq i \leq n$$

$$f(e_i) = \begin{cases} 1, & \text{if } i \equiv 0 \pmod{2} \\ 2, & \text{if } i \equiv 1 \pmod{2} \end{cases} \quad \text{for } 2 \leq i \leq n-2$$

$$f(e_1) = 4, \quad f(e_{n-1}) = f(e_n) = 1, \quad f(e_i') = 5, \quad \text{for } 1 \leq i \leq n$$

$$f(e_i'') = \begin{cases} 3, & \text{if } i \equiv 0 \pmod{2} \\ 4, & \text{if } i \equiv 1 \pmod{2} \end{cases} \quad \text{for } 1 \leq i \leq n$$

Using the above pattern of the coloring, we see that the graph $M(C_n)$ is equitable total colored with 5 colors. The color classes of $M(C_n)$ are grouped as $T(M(C_n)) = \{T_1, T_2, T_3, T_4, T_5\}$. We observe that these colors classes T_1, T_2, T_3, T_4, T_5 are independent sets of $M(C_n)$ and its satisfies the

condition $||T_i| - |T_j|| \leq 1$, for $i \neq j$. For example consider the middle graph of cycle with 6 vertices (see Figure 1), for which $|T_1| = |T_2| = |T_3| = |T_4| = |T_5| = 7$ and it holds the inequality $||T_i| - |T_j|| \leq 1$, for $i \neq j$ and it is equitable total colored with 5 colors. Hence $\chi_{et}(M(C_n)) \leq 5$. Further, since $\Delta = 4$, we have $\chi_{et}(M(C_n)) \geq \chi''(M(C_n)) \geq \Delta + 1 \geq 4 + 1 \geq 5$. Therefore $\chi_{et}(M(C_n)) = 5$. \square

Theorem 3.3. For any positive integer $n \geq 3$, $\chi_{et}(T(P_n)) = 5$.

Proof. Let $V(P_n) = \{v_i : 1 \leq i \leq n\}$ and $E(P_n) = \{e_i : 1 \leq i \leq n-1\}$, where $\{e_i : 1 \leq i \leq n-1\}$ be the edges $v_i v_{i+1}$ ($1 \leq i \leq n-1$). By the definition of total graph, each edge $\{e_i : 1 \leq i \leq n-1\}$ is subdivided by the vertices $\{u_i : 1 \leq i \leq n-1\}$. In $T(P_n)$, the vertex set and the edge set is given by $V(T(P_n)) = \{v_i : 1 \leq i \leq n\} \cup \{u_i : 1 \leq i \leq n-1\}$ and $E(T(P_n)) = \{e_i : 1 \leq i \leq n-1\} \cup \{e'_i : 1 \leq i \leq n-1\} \cup \{e''_i : 1 \leq i \leq n-1\} \cup \{e'''_i : 1 \leq i \leq n-2\}$, where e_i ($1 \leq i \leq n-1$) is an edge $v_i v_{i+1}$ ($1 \leq i \leq n-1$), e'_i ($1 \leq i \leq n-1$) is an edge $u_i v_{i+1}$ ($1 \leq i \leq n-1$), e''_i ($1 \leq i \leq n-1$) is an edge $v_i v_{i+1}$ ($1 \leq i \leq n-1$) and e'''_i ($1 \leq i \leq n-2$) is an edge $u_i u_{i+1}$ ($1 \leq i \leq n-2$).

Define an equitable total coloring f , such that $f : S \rightarrow C$, where $S = V(T(P_n)) \cup E(T(P_n))$ and $C = \{1, 2, 3, 4, 5\}$. The equitable total coloring is obtained by coloring these vertices and edges as follows.

$$f(v_i) = \begin{cases} 1, & \text{if } i \equiv 1 \pmod{5} \\ 3, & \text{if } i \equiv 2 \pmod{5} \\ 5, & \text{if } i \equiv 3 \pmod{5} \\ 2, & \text{if } i \equiv 4 \pmod{5} \\ 4, & \text{if } i \equiv 0 \pmod{5} \end{cases} \text{ for } 1 \leq i \leq n$$

$$f(u_i) = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{5} \\ 4, & \text{if } i \equiv 2 \pmod{5} \\ 1, & \text{if } i \equiv 3 \pmod{5} \\ 3, & \text{if } i \equiv 4 \pmod{5} \\ 5, & \text{if } i \equiv 0 \pmod{5} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$f(e_i) = \begin{cases} 4, & \text{if } i \equiv 1 \pmod{5} \\ 1, & \text{if } i \equiv 2 \pmod{5} \\ 3, & \text{if } i \equiv 3 \pmod{5} \\ 5, & \text{if } i \equiv 4 \pmod{5} \\ 2, & \text{if } i \equiv 0 \pmod{5} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$f(e_i') = \begin{cases} 5, & \text{if } i \equiv 1 \pmod{5} \\ 2, & \text{if } i \equiv 2 \pmod{5} \\ 4, & \text{if } i \equiv 3 \pmod{5} \\ 1, & \text{if } i \equiv 4 \pmod{5} \\ 3, & \text{if } i \equiv 0 \pmod{5} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$f(e_i'') = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{5} \\ 4, & \text{if } i \equiv 2 \pmod{5} \\ 1, & \text{if } i \equiv 3 \pmod{5} \\ 3, & \text{if } i \equiv 4 \pmod{5} \\ 5, & \text{if } i \equiv 0 \pmod{5} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$f(e_i''') = \begin{cases} 3, & \text{if } i \equiv 1 \pmod{5} \\ 5, & \text{if } i \equiv 2 \pmod{5} \\ 2, & \text{if } i \equiv 3 \pmod{5} \\ 4, & \text{if } i \equiv 4 \pmod{5} \\ 1, & \text{if } i \equiv 0 \pmod{5} \end{cases} \text{ for } 1 \leq i \leq n-2$$

Based on the above method of coloring, it is clear that the graph $T(P_n)$




Figure 2: Equitable Total Coloring of total graph of Path $T(P_6)$

is equitable total colored with 5 colors, such that the color classes grouped are $T(T(P_n)) = \{T_1, T_2, T_3, T_4, T_5\}$. We observe that these color classes $\{T_1, T_2, T_3, T_4, T_5\}$ are independent sets of $T(P_n)$ and it holds the inequality $||T_i| - |T_j|| \leq 1$, for $i \neq j$. For example consider the total graph of path with 6 vertices (see Figure 2), for which $|T_1| = |T_3| = |T_4| = |T_5| = 7$ and $|T_2| = 8$ and it satisfies the condition $||T_i| - |T_j|| \leq 1$, for $i \neq j$ and so it is

equitable total colored with 5 colors. Hence $\chi_{et}(T(P_n)) \leq 5$. Further, since $\Delta = 4$, we have $\chi_{et}(T(P_n)) \geq \chi''(T(P_n)) \geq \Delta + 1 \geq 4 + 1 \geq 5$. Therefore $\chi_{et}(T(P_n)) = 5$. \square

Theorem 3.4. For any positive integer n , $\chi_{et}(T(C_n)) = \begin{cases} 5, & n = 3 \\ 6, & n \neq 3 \end{cases}$

Proof. Let $V(C_n) = \{v_i : 1 \leq i \leq n\}$ and $E(C_n) = \{e_i : 1 \leq i \leq n\}$, where $\{e_i : 1 \leq i \leq n\}$ be the edges $v_i v_{i+1}, i + 1$ taken modulo $n(1 \leq i \leq n)$. By the definition of total graph, each edge $\{e_i : 1 \leq i \leq n\}$ is subdivided by the vertices $\{u_i : 1 \leq i \leq n\}$. In $T(C_n)$, the vertex set and the edge set is given by $V(T(C_n)) = \{v_i : 1 \leq i \leq n\} \cup \{u_i : 1 \leq i \leq n\}$ and $E(T(C_n)) = \{e_i : 1 \leq i \leq n\} \cup \{e'_i : 1 \leq i \leq n\} \cup \{e''_i : 1 \leq i \leq n\} \cup \{e'''_i : 1 \leq i \leq n\}$, where $e_i(1 \leq i \leq n)$ be an edge $v_i u_i(1 \leq i \leq n)$, $e'_i(1 \leq i \leq n)$ be an edge $u_i v_{i+1}, i + 1$ taken modulo $n(1 \leq i \leq n)$, $e''_i(1 \leq i \leq n)$ be an edge $v_i v_{i+1}, i + 1$ taken modulo $n(1 \leq i \leq n)$ and $e'''_i(1 \leq i \leq n)$ be an edge $u_i u_{i+1}, i + 1$ taken modulo $n(1 \leq i \leq n)$.

We define a function f , such that $f : S \rightarrow C$, where $S = V(T(C_n)) \cup E(T(C_n))$ and $C = \{1, 2, 3, 4, 5, 6\}$. Now we assign an equitable total coloring to these vertices and edges as follows. we consider following three cases
Case(i): When $n = 3$

$$f(v_i) = \begin{cases} 1, & \text{if } i \equiv 1 \pmod{3} \\ 3, & \text{if } i \equiv 2 \pmod{3} \\ 2, & \text{if } i \equiv 0 \pmod{3} \end{cases} \text{ for } 1 \leq i \leq 3$$

$$f(u_i) = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{3} \\ 3, & \text{if } i \equiv 2 \pmod{3} \\ 1, & \text{if } i \equiv 0 \pmod{3} \end{cases} \text{ for } 1 \leq i \leq 3$$

$$\begin{aligned} f(e_i) &= 4, & \text{for } 1 \leq i \leq 3 \\ f(e'_i) &= 5, & \text{for } 1 \leq i \leq 3 \end{aligned}$$

$$f(e''_i) = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{3} \\ 1, & \text{if } i \equiv 2 \pmod{3} \\ 3, & \text{if } i \equiv 0 \pmod{3} \end{cases} \text{ for } 1 \leq i \leq 3$$

$$f(e'''_i) = \begin{cases} 3, & \text{if } i \equiv 1 \pmod{3} \\ 2, & \text{if } i \equiv 2 \pmod{3} \\ 1, & \text{if } i \equiv 0 \pmod{3} \end{cases} \text{ for } 1 \leq i \leq 3$$

From the above rule of total coloring, it is easy to see that the graph $T(C_3)$ is equitable total colored with 5 colors. The color classes of $T(C_3)$ are

grouped as $T(T(C_3)) = \{T_1, T_2, T_3, T_4, T_5\}$. We observe that these color classes $\{T_1, T_2, T_3, T_4, T_5\}$ are independent sets of $T(C_3)$ and it satisfies the inequality $||T_i| - |T_j|| \leq 1$, for $i \neq j$, for which $|T_1| = |T_2| = |T_3| = 4$ and $|T_4| = |T_5| = 3$ and it satisfies the inequality $||T_i| - |T_j|| \leq 1$, for $i \neq j$ and it is equitable total colored with 5 colors. Hence $\chi_{et}(T(C_3)) \leq 5$. Further, since $\Delta = 4$, we have $\chi_{et}(T(C_3)) \geq \chi''(T(C_3)) \geq \Delta + 1 \geq 4 + 1 \geq 5$. Therefore $\chi_{et}(T(C_3)) = 5$.




Figure 3: Equitable Total Coloring of total graph of Cycle $T(C_6)$

Case(ii): When $n > 3$, $n = 2k$, $k = 2, 3, \dots$

$$f(v_i) = \begin{cases} 1, & \text{if } i \equiv 1 \pmod{2} \\ 3, & \text{if } i \equiv 0 \pmod{2} \end{cases} \text{ for } 1 \leq i \leq n$$

$$f(u_i) = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{2} \\ 4, & \text{if } i \equiv 0 \pmod{2} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$f(e_i) = 5, \quad \text{for } 1 \leq i \leq n \\ f(e'_i) = 6, \quad f(e'_n) = 6 \quad \text{for } 1 \leq i \leq n-1$$

$$f(e''_i) = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{2} \\ 4, & \text{if } i \equiv 0 \pmod{2} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$f(e_n'') = 4$$

$$f(e_i''') = \begin{cases} 3, & \text{if } i \equiv 1 \pmod{2} \\ 1, & \text{if } i \equiv 0 \pmod{2} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$f(e_n''') = 1$$

Case(iii): When $n > 3$, $n = 2k + 1$, $k = 2, 3, \dots$

$$f(v_i) = \begin{cases} 1, & \text{if } i \equiv 1 \pmod{2} \\ 3, & \text{if } i \equiv 0 \pmod{2} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$f(v_n) = 5$$

$$f(u_i) = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{2} \\ 4, & \text{if } i \equiv 0 \pmod{2} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$\begin{aligned} f(u_n) &= 6, \quad f(e_i) = 5, \quad \text{for } 1 \leq i \leq n-2 \\ f(e_{n-1}) &= 1, \quad f(e_n) = 3 \\ f(e_i') &= 6, \quad f(e_{n-1}') = 2, \quad f(e_n') = 4 \quad \text{for } 1 \leq i \leq n-2 \end{aligned}$$

$$f(e_i'') = \begin{cases} 2, & \text{if } i \equiv 1 \pmod{2} \\ 4, & \text{if } i \equiv 0 \pmod{2} \end{cases} \text{ for } 1 \leq i \leq n-1$$

$$f(e_n'') = 6$$

$$f(e_i''') = \begin{cases} 3, & \text{if } i \equiv 1 \pmod{2} \\ 1, & \text{if } i \equiv 0 \pmod{2} \end{cases} \text{ for } 1 \leq i \leq n-2$$

$$f(e_{n-1}''') = 5, \quad f(e_n''') = 6$$

The above pattern of total coloring, the graph $T(C_n)$ is equitable total colored with 6 colors. The color classes of $T(C_n)$ are grouped as $T(T(C_n)) = \{T_1, T_2, T_3, T_4, T_5, T_6\}$. We observe that these color classes $\{T_1, T_2, T_3, T_4, T_5, T_6\}$ are independent sets of $T(C_n)$ and its satisfies the inequality $||T_i| - |T_j|| \leq 1$, for $i \neq j$. For example consider the total graph of path with 6 vertices (see Figure 3), for which $|T_1| = |T_2| = |T_3| = |T_4| = |T_5| = |T_6| = 7$ and it satisfies the inequality $||T_i| - |T_j|| \leq 1$, for $i \neq j$. When $n \neq 3$, in this case every vertex is adjacent to exactly four vertices of same degree and due to the incidence and adjacency of elements, five colors can not be suffice for the total coloring. Thus $\chi_{et}(T(C_n)) \geq \chi''(T(C_n)) \neq 5$. Hence $\chi_{et}(T(C_n)) = 6$. \square

References

- [1] Behzad. M, Graphs and Their chromatic numbers, Ph.D. Thesis, Michigan State University, 1965.
- [2] Girija.G, Veninstine Vivek.J, Equitable Total Coloring of Some Graphs, International J. Math. Combin. Vol.1(2015), 107-112.
- [3] Gong Kun, Zhang Zhongfu, Wang Jian Fang, Equitable total coloring of some join graphs, Journal of mathematical Research and Exposition, 28(4),(2008),823-828. DOI:10.3770/j.issn:1000-341X.2008.04.010
- [4] Hung-lin Fu, Some results on equalized total coloring, Congr. Numer., 102,(1994), 111-119.
- [5] Kostochka. A.V, The total coloring of multigraph with maximal degree 4, Discrete Math, 17, (1989),161-163.
- [6] Ma Gang, Ma Ming, The equitable total chromatic number of the some join-graphs,Open Journal of Applied Sciences,(2012),96-99.
- [7] Ma Gang, Zhang Zhong-fu, On the equitable total coloring of multiple join graph, Journal of Mathematical Research and Exposition, 2007, 27 (2) 351-354.
- [8] Tong chunling, lin Xiaohui, Yang Yuansheng, Lizhihe, Equitable Total Coloring of $C_m \square C_n$, Discrete Applied Mathematics, 157,(2009), 596-601.
- [9] Veninstine Vivek.J, Girija. G, An Algorithmic Approach to Equitable Total Chromatic Number of Graphs, Proyecciones Journal of Mathematics, 36(2)(2017),307-324.
- [10] Vizing. V.G, On an estimate of the chromatic class of a p - graph(in Russian), Metody Diskret. Analiz., 5, (1964),25-30.
- [11] Wang Wei-fan, Equitable Total Coloring of Graphs with Maximum Degree 3, Graphs and Combin, 2002, 18, 677-685.
- [12] Zhang Zhong-fu, wang Wei-fan, Bau Sheng. et al. On the equitable total coloring of some join graphs, J.Info Comput. Sci., 2(4),(2005), 829-834.
- [13] Zhang Zhong-fu, Zhang Jianxun, Wang Jianfang, The total chromatic number of some graph, Sci. Sinica, Ser.A, 31(12),(1988) 1434-1441.