

Algebraic Characterization of the SSC $\Delta_s(\mathcal{G}_{n,r}^1)$

Agha Kashif¹, Zahid Raza², Imran Anwar³.

1. Department of Mathematics, University of Management and Technology, Lahore, Pakistan.

2. University of Sharjah, College of Sciences, Department of Mathematics, United Arab Emirates.

3. Abdus Salam School of Mathematical Sciences, Government College University, Lahore, Pakistan.

kashif.khan@umt.edu.pk, aghakashifkhan@hotmail.com,
zraza@sharjah.ac.ae, iimrananwar@gmail.com

Abstract

In this paper, we characterize the set of spanning trees of $\mathcal{G}_{n,r}^1$ (a simple connected graph consisting of n edges, containing exactly one 1-edge-connected chain of r cycles C_r^1 and $\mathcal{G}_{n,r}^1 \setminus C_r^1$ is a forest). We compute the Hilbert series of the face ring $k[\Delta_s(\mathcal{G}_{n,r}^1)]$ for the spanning simplicial complex $\Delta_s(\mathcal{G}_{n,r}^1)$. Also, we characterize associated primes of the facet ideal $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$. Furthermore, we prove that the face ring $k[\Delta_s(\mathcal{G}_{n,r}^1)]$ is Cohen-Macaulay.

Keywords : simplicial complex, f -vector, face ring, facet ideal, spanning trees, primary decomposition, Hilbert series, Cohen-Macaulay ring.

(2000) Mathematics Subject Classification : Primary 13P10, Secondary 13H10, 13F20, 13C14.

1 Introduction

The study of simplicial complexes arising from a simple graph has been an important topic and attracted good literature. One popular chapter of this literature is the complementary simplicial complex Δ_G of a graph G ; for example, see [13]. The notion of spanning simplicial complex (SSC) $\Delta_s(G)$ associated to a simple connected graph $G(V, E)$ was firstly introduced in

[1]. For *uni-cyclic graphs* $U_{n,m}$, it is proved that $\Delta_s(U_{n,m})$ is *shifted* in [1]. Zhu, Shi and Geng [14] further investigated the algebraic and combinatorial properties of SSC associated to n -cyclic graphs with a common edge. In [9], the authors investigated the algebraic properties of SSC $\Delta_s(G_{n,r})$ associated to r -cyclic graphs $G_{n,r}$ (containing exactly r cycles having no edge in common). Moreover, they proved that the facet ideal $I_{\mathcal{F}}(\Delta_s(G_{n,r}))$ has linear quotients with respect to its generating set and computed the *betti numbers* of $I_{\mathcal{F}}(\Delta_s(G_{n,r}))$ for particular cases. Some other interesting classes of simple finite connected graphs are studied for SSC by Pan, Li and Zhu in [11], Guo and Wu in [6] and Raza, Kashif and Anwar in [12]. In this paper, we investigate the class of spanning simplicial complexes $\Delta_s(\mathcal{G}_{n,r}^1)$ associated to $\mathcal{G}_{n,r}^1$. Where $\mathcal{G}_{n,r}^1$ is a connected graph having n edges, containing exactly one *1-edge-connected chain* of r cycles \mathcal{C}_r^1 and $\mathcal{G}_{n,r}^1 \setminus \mathcal{C}_r^1$ is a *forest*. In other words, $\mathcal{G}_{n,r}^1$ is a graph consisting of r cycles such that every pair of consecutive cycles have exactly one edge common between them. If C_1, C_2, \dots, C_r are the r cycles of the graph $\mathcal{G}_{n,r}^1$ forming \mathcal{C}_r^1 with respective lengths m_1, m_2, \dots, m_r then we fix the label of edge set of $\mathcal{G}_{n,r}^1$ as follows;

$$E = \{e_{11}, \dots, e_{1m_1}, e_{21}, \dots, e_{2m_2-1}, \dots, e_{r1}, \dots, e_{rm_r-1}, e_1, \dots, e_t\} \quad (1)$$

where, $t = n - \sum_{i=1}^r m_i + (r-1)$ and $\{e_{i1}, \dots, e_{iv}\}$ is the edge-set of i th-cycle such that $v = m_1$ for $i = 1$, $v = m_i - 1$ for $i > 1$ and e_{i1} always represents the common edge between i th and $(i+1)$ th-cycle (for $1 \leq i < r$). We give the characterization of $s(\mathcal{G}_{n,r}^1)$ in 3.4. The formulation for f -vectors is presented in 3.5 which further applied to device a formula to compute the *Hilbert series* of the *face ring* $k[\Delta_s(\mathcal{G}_{n,r}^1)]$ (see 3.7). Moreover in 4.1, we characterize all the associated primes of the facet ideal $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$. Finally, we prove that the face ring $k[\Delta_s(\mathcal{G}_{n,r}^1)]$ is Cohen-Macaulay in 5.4.

2 Background and basic notions

In this section, we give some background and preliminaries of the topic and define some notions that will be useful in the sequel.

Definition 2.1. A *spanning tree* of a simple connected finite graph $G(V, E)$ is a subtree of G that contains every vertex of G . We represent the collection of all edge-sets of the spanning trees of G by $s(G)$, in other words;

$$s(G) := \{E(T_i) \subset E, \text{ where } T_i \text{ is a spanning tree of } G\}.$$

For any simple connected graph G , the authors mentioned the *cutting-down method* to obtain all the spanning trees of G in [1]. According to this

method a spanning tree is obtained by removing one edge from each cycle appearing in the graph. However, for the graph $\mathcal{G}_{n,r}^1$ with r cycles having one edge common in every consecutive cycles and the labeling given in (1), one can obtain its spanning trees by removing exactly r edges from the graph with not more than two edges deleted from any cycle. Also, keeping in view that if a common edge between two cycles is removed then only one edge can be removed from the non common edges explicitly from the cycles on the either side of the common edge.

For example by using the above said *cutting-down method* for the graph $\mathcal{G}_{10,2}^1$ given in fig. 1:

$$s(\mathcal{G}_{10,2}^1) = \{ \{e_1, e_2, e_3, e_4, e_{13}, e_{11}, e_{23}, e_{21}\}, \{e_1, e_2, e_3, e_4, e_{13}, e_{11}, e_{23}, e_{22}\}, \\ \{e_1, e_2, e_3, e_4, e_{13}, e_{11}, e_{21}, e_{22}\}, \{e_1, e_2, e_3, e_4, e_{13}, e_{23}, e_{21}, e_{22}\}, \{e_1, e_2, e_3, \\ e_4, e_{12}, e_{11}, e_{23}, e_{21}\}, \{e_1, e_2, e_3, e_4, e_{12}, e_{11}, e_{23}, e_{22}\}, \{e_1, e_2, e_3, e_4, e_{12}, e_{11}, \\ e_{21}, e_{22}\}, \{e_1, e_2, e_3, e_4, e_{12}, e_{23}, e_{21}, e_{22}\}, \{e_1, e_2, e_3, e_4, e_{13}, e_{12}, e_{23}, e_{22}\}, \\ \{e_1, e_2, e_3, e_4, e_{13}, e_{12}, e_{23}, e_{21}\}, \{e_1, e_2, e_3, e_4, e_{13}, e_{12}, e_{21}, e_{22}\}, \{e_1, e_2, e_3, \\ e_4, e_{13}, e_{23}, e_{21}, e_{22}\}, \{e_1, e_2, e_3, e_4, e_{12}, e_{23}, e_{21}, e_{22}\} \}$$




Fig. 1. $\mathcal{G}_{10,2}^1$

Definition 2.2. A *simplicial complex* Δ over a finite set $[n] = \{1, 2, \dots, n\}$ is a collection of subsets of $[n]$, with the property that $\{i\} \in \Delta$ for all $i \in [n]$, and if $F \in \Delta$ then Δ will contain all the subsets of F (including the empty set). An element of Δ is called a *face* of Δ , and the *dimension* of a face F of Δ is defined as $|F| - 1$, where $|F|$ is the number of vertices of F . The maximal faces of Δ under inclusion are called *facets* of Δ . The *dimension* of the simplicial complex Δ is :

$$\dim \Delta = \max\{\dim F \mid F \in \Delta\}.$$

We denote the simplicial complex Δ with facets $\{F_1, \dots, F_q\}$ by

$$\Delta = \langle F_1, \dots, F_q \rangle$$

Definition 2.3. For a simplicial complex Δ over $[n]$ having dimension d , its *f - vector* is a $d + 1$ -tuple, defined as:

$$f(\Delta) = (f_0, f_1, \dots, f_d)$$

where f_i denotes the number of i -dimensional faces in Δ .

Definition 2.4. (Spanning Simplicial Complex)

Let $G(V, E)$ be a simple finite connected graph and $s(G) = \{E_1, E_2, \dots, E_t\}$ be the edge-set of all possible spanning trees of $G(V, E)$, then we defined (in [1]) a simplicial complex $\Delta_s(G)$ on E such that the facets of $\Delta_s(G)$ are precisely the elements of $s(G)$, we call $\Delta_s(G)$ as the *spanning simplicial complex* of $G(V, E)$. In other words;

$$\Delta_s(G) = \langle E_1, E_2, \dots, E_t \rangle.$$

Here we recall a definition from [4].

Definition 2.5. Let Δ be a simplicial complex with vertex set $V = [n]$ and facets F_1, F_2, \dots, F_q . A *vertex cover* for Δ is a subset A of V such that $A \cap F_i \neq \emptyset$ for all $i \in \{1, 2, \dots, q\}$. A *minimal vertex cover* of Δ is a subset A of V such that A is a *vertex cover*, and no proper subset of A is a *vertex cover* for Δ .

For example, the *minimal vertex covers* for the *spanning simplicial complex* $\Delta_s(\mathcal{G}_{10,2}^1)$ given in Fig. 1, are as follows:

$$\{e_1\}, \{e_2\}, \{e_3\}, \{e_4\}, \{e_{13}, e_{12}\}, \{e_{23}, e_{22}\}, \{e_{23}, e_{21}\}, \{e_{22}, e_{21}\}$$

3 Spanning trees of $\mathcal{G}_{n,r}^1$ and Face ring $\Delta_s(\mathcal{G}_{n,r}^1)$

In this section, we discuss the combinatorial properties of $\mathcal{G}_{n,r}^1$. We use $\tau(\mathcal{G}_{n,r}^1)$ to denote the **total number of cycles** contained in $\mathcal{G}_{n,r}^1$. We begin with the elementary result, that tells the total number of cycles contained by $\mathcal{G}_{n,r}^1$.

Proposition 3.1. The total number of cycles in the graph $\mathcal{G}_{n,r}^1$ will be

$$\tau(\mathcal{G}_{n,r}^1) = \frac{r(r+1)}{2}$$

Proof. As the graph $\mathcal{G}_{n,r}^1$ contains one-edge connected chain \mathbb{C}_r^1 of r cycles $\{C_1, C_2, \dots, C_r\}$. By removing the common edges between any number of consecutive cycles, we obtain a cycle by the remaining edges. The cycle obtained in this way by adjoining consecutive cycles $C_i, C_{i+1}, \dots, C_{i+k}$ is denoted by $C_{i,i+1,\dots,i+k}$. Therefore, we get the following cycles

$$C_{1,2}, C_{2,3}, \dots, C_{r-1,r}, C_{1,2,3}, \dots, C_{r-2,r-1,r}, \dots, C_{2,3,\dots,r}, C_{12,3,\dots,r}$$

Hence, the set of all possible cycles contained in the graph $\mathcal{G}_{n,r}^1$ will be

$$\{C_{i,i+1,\dots,i+k} \mid i \in \{1, 2, \dots, r-k\} \text{ and } 0 \leq k \leq r-1\}.$$

Therefore, we get the total number of cycles contained in the graph $\mathcal{G}_{n,r}^1$ as

$$\tau(\mathcal{G}_{n,r}^1) = \sum_{k=0}^{r-1} \sum_{i=1}^{r-k} 1 = \frac{r(r+1)}{2}.$$

□

It is clear from above proposition that the cycle $C_{i,i+1,\dots,i+k}$ is obtained by removing the common edges between the adjacent cycles $C_i, C_{i+1}, \dots, C_{i+k}$. We denote the length of cycle $C_{i,i+1,\dots,i+k}$ by $|C_{i,i+1,\dots,i+k}|$.

Proposition 3.2. Let $\mathcal{G}_{n,r}^1$ be a graph containing the one-edge connected chain \mathcal{C}_r^1 of r cycles $\{C_1, C_2, \dots, C_r\}$, then the length of cycle $C_{i,i+1,\dots,i+k}$ will be

$$|C_{i,i+1,\dots,i+k}| = \sum_{\alpha=0}^k |C_{i+\alpha}| - 2k.$$

Proof. It is clear from above that $C_{i,i+1,\dots,i+k}$ is obtained by deleting the common edges shared by the adjacent cycles $\{C_i, C_{i+1}, \dots, C_{i+k}\}$ in $\mathcal{G}_{n,r}^1$. Therefore, the length of the cycle $C_{i,i+1,\dots,i+k}$ is obtained by adding lengths of all $C_i, C_{i+1}, \dots, C_{i+k}$ and subtracting $2k$ from it, since the common edges are being counted twice. Hence, we have

$$|C_{i,i+1,\dots,i+k}| = \sum_{\alpha=0}^k |C_{i+\alpha}| - 2k.$$

□

We use $|C_{i,i+1,\dots,i+k} \cap C_{j,j+1,\dots,j+l}|$ to denote the number of edges shared by the cycles $C_{i,i+1,\dots,i+k}$ and $C_{j,j+1,\dots,j+l}$. The following proposition characterizes $|C_{i,i+1,\dots,i+k} \cap C_{j,j+1,\dots,j+l}|$ in $\mathcal{G}_{n,r}^1$.

Proposition 3.3. Let $\mathcal{G}_{n,r}^1$ be a graph containing the one-edge connected chain \mathcal{C}_r^1 of r cycles $\{C_1, C_2, \dots, C_r\}$ of lengths m_1, m_2, \dots, m_r , then for $1 \leq k \leq l \leq r$ we have

$$|C_{i,i+1,\dots,i+k} \cap C_{j,j+1,\dots,j+l}| = \begin{cases} 1, & i+k=j-1; \\ |C_{j,j+1,\dots,j+\alpha}| - 2, & i+k=j+\alpha; 0 \leq \alpha \leq k-1; \\ |C_{j,j+1,\dots,j+k}| - 1, & i+k=j+k; \\ |C_{j,j+1,\dots,j+l}|, & i+k=j+l \text{ and } l=k; \\ |C_{i,i+1,\dots,i+k}| - 2, & i+k=j+\alpha; k+1 \leq \alpha \leq l-1; \\ |C_{i,i+1,\dots,i+k}| - 1, & i+k=j+l; \\ |C_{i,i+1,\dots,i+k-\alpha}| - 2, & i+k=j+l+\alpha; 1 \leq \alpha \leq k; \\ 1, & i=j+l+1; \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Here we denote $m_{i,i+1,\dots,i+k} = |C_{i,i+1,\dots,i+k}|$.

Now for $1 \leq k \leq l \leq r$ we discuss the following cases for

$$|C_{i,i+1,\dots,i+k} \cap C_{j,j+1,\dots,j+l}| :$$

Case (i) If $i+k = j-1$, then the right most edges of the cycle $C_{i,i+1,\dots,i+k}$ are from its adjoining cycle C_{i+k} and the left most edges of the cycle $C_{j,j+1,\dots,j+l}$ are from its adjoining cycle C_{j+l} , and since C_{i+k} and C_{j+l} are consecutive so they have only one edge in common.

Case (ii) If $i+k = j+\alpha$; $0 \leq \alpha \leq k-1$, then the left most α adjoining cycles of the cycles of $C_{j,j+1,\dots,j+l}$, i.e., $C_j, C_{j+1}, \dots, C_{j+\alpha}$ coincide with the right most α adjoining cycles of the cycles of $C_{i,i+1,\dots,i+k}$. Therefore, the intersection $C_{i,i+1,\dots,i+k} \cap C_{j,j+1,\dots,j+l}$ will contain all edges of $C_{j,j+1,\dots,j+\alpha}$ except its two edges, one the edge of $C_{j,j+1,\dots,j+\alpha}$ which is the common edge of $C_{j+\alpha}$ and $C_{j+\alpha+1}$ and second the edge of $C_{j,j+1,\dots,j+\alpha}$ which is common edge between C_j and C_{j-1} .

Case (iii) If $i+k = j+k$ and $k < l$, then $i = j$ and therefore the cycle $C_{i,i+1,\dots,i+k}$ lies completely in the cycle $C_{j,j+1,\dots,j+l}$ except its one edge which is the common edge between its adjoining cycle C_{j+k} and the cycle C_{j+k+1} . Therefore the intersection $C_{i,i+1,\dots,i+k} \cap C_{j,j+1,\dots,j+l}$ will contain all edges of $C_{j,j+1,\dots,j+k}$ except one.

Case (iv) If $i+k = j+l+\alpha$; $1 \leq \alpha \leq k$, then the left most $k-\alpha+1$ adjoining cycles of the cycle $C_{i,i+1,\dots,i+k}$, which are $C_i, C_{i+1}, \dots, C_{i+k-\alpha}$ coincide with the right most $k-\alpha+1$ adjoining cycles of the cycle $C_{j,j+1,\dots,j+l}$. Hence the intersection $C_{i,i+1,\dots,i+k} \cap C_{j,j+1,\dots,j+l}$ will contain all the edges of the cycle $C_{i,i+1,\dots,i+k-\alpha}$ except two; one is the common edge of its adjoining cycle C_i and the cycle C_{i-1} and the other is the common edge of its adjoining cycle $C_{i+k-\alpha}$ and the cycle $C_{i+k-\alpha+1}$.

The remaining cases can be proved in the similar way. \square

Lemma 3.4. Characterization of $s(\mathcal{G}_{n,r}^1)$

Let $\mathcal{G}_{n,r}^1$ be the r -cycles graph with edge set E as defined in eq (1), then a subset $E(T_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}) \subset E$, where $j_\alpha \in \{1, 2, \dots, r\}$, $i_\alpha \in \{1, 2, \dots, m_{j_\alpha} - 1\}$; $j_\alpha \geq 2$ and $i_\alpha \in \{1, 2, \dots, m_1\}$; $j_\alpha = 1$, will belong to $s(\mathcal{G}_{n,r}^1)$ if and only if it satisfies any of the following:

1. If $j_\alpha i_\alpha \neq j_\alpha 1$ for all α except for which $j_\alpha i_\alpha = r i_\alpha$, then $E(T_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}) = E \setminus \{e_{1i_1}, e_{2i_2}, \dots, e_{ri_r}\}$

2. If $j_\alpha i_\alpha = j_\alpha 1$ for any α , then $E(T_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}) = E \setminus \{e_{j_1 i_1}, e_{j_2 i_2}, \dots, e_{j_r i_r}\}$ where, $\{e_{j_1 i_1}, e_{j_2 i_2}, \dots, e_{j_{\alpha-1} i_{\alpha-1}}, e_{j_{\alpha+1} i_{\alpha+1}}, \dots, e_{j_r i_r}\}$ will contain exactly one edge from $C_{j_\alpha(j_\alpha+1)} \setminus \{e_{(j_\alpha-1)1}, e_{j_\alpha 1}\}$.
3. If $j_\alpha i_\alpha = j_\alpha 1$ for $\alpha \in \{r_1, r_1 + 1, \dots, r_2\}$, where $1 \leq r_1 < r_2 < r$ then
 - (a) If $e_{j_{r_1} 1}, e_{j_{(r_1+1)} 1}, \dots, e_{j_{r_2} 1}$ are common edges from consecutive cycles then $E(T_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}) = E \setminus \{e_{j_1 i_1}, e_{j_2 i_2}, \dots, e_{j_r i_r}\}$ such that $\{e_{j_1 i_1}, e_{j_2 i_2}, \dots, e_{j_r i_r}\} \setminus \{e_{j_{r_1} 1}, e_{j_{(r_1+1)} 1}, \dots, e_{j_{r_2} 1}\}$ will contain exactly one edge from $C_{j_{r_1} j_{(r_1+1)} \dots j_{r_2}} \setminus \{e_{(j_{r_1}-1)1}, e_{j_{r_2} 1}\}$.
 - (b) If none of $e_{j_{r_1} 1}, e_{j_{(r_1+1)} 1}, \dots, e_{j_{r_2} 1}$ are common edges from consecutive cycles then $E(T_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}) = E \setminus \{e_{j_1 i_1}, e_{j_2 i_2}, \dots, e_{j_r i_r}\}$ such that for each edge $e_{j_{r_t} 1}$ case 2 holds.
 - (c) If some of $e_{j_{r_1} 1}, e_{j_{(r_1+1)} 1}, \dots, e_{j_{r_2} 1}$ are common edges from consecutive cycles then $E(T_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}) = E \setminus \{e_{j_1 i_1}, e_{j_2 i_2}, \dots, e_{j_r i_r}\}$ such that (3.(a)) is satisfied for the common edges of consecutive cycles and (3.(b)) is satisfied for remaining common edges.

In particular, if we denote the above classes of subsets of E by $C_{(1)}, C_{(2)}, C_{(3a)}, C_{(3b)}, C_{(3c)}$ respectively then,

$$s(\mathcal{G}_{n,r}^1) = C_{(1)} \cup C_{(2)} \cup C_{(3a)} \cup C_{(3b)} \cup C_{(3c)}$$

Proof. Since $\mathcal{G}_{n,r}^1$ is a r -cycles graph with cycles C_1, C_2, \dots, C_r and $e_{11}, e_{21}, \dots, e_{(r-1)1}$ as common edges between consecutive cycles and by cutting down process a total of r edges must be removed with not more than one edges from the non common edges of each cycle. Therefore, in order to obtain a spanning tree of $\mathcal{G}_{n,r}^1$ with none of common edges $e_{11}, e_{21}, \dots, e_{(r-1)1}$ to be removed, we need to remove exactly one edge from the non common edges from each cycle. This explains the case (1) of the above lemma.

Now for a spanning tree of $\mathcal{G}_{n,r}^1$ such that exactly one common edge $e_{j_\alpha 1}$ is removed, we need to remove precisely $r - 1$ edges using cutting down process from the remaining edges. However, from the non common edges of the cycle $C_{j_\alpha(j_\alpha+1)}$, we cannot remove more than one edge (since that will result in a disconnected graph). This explains the proof of case of (2) of the lemma.

Next for the case (3.a), we need to obtain a spanning tree of $\mathcal{G}_{n,r}^1$ such that $r_2 - r_1$ common edges must be removed from consecutive cycles. If $C_{j_{r_1}}, C_{j_{r_1+1}}, \dots, C_{j_{r_2}}$ are consecutive cycles then the remaining $r - (r_1 - r_2)$ edges must be removed in such a way that exactly one edge is removed from the non common edges of $C_{j_{r_1} j_{(r_1+1)} \dots j_{r_2}}$ and the remaining $r - (r_1 - r_2)$ cycles of the graph $\mathcal{G}_{n,r}^1$ which concludes the case.

The remaining cases of the lemma can be visualised in similar manner using the above cases . Consequently, if we denote the above disjoint classes of subsets of E by $C_{(1)}, C_{(2)}, C_{(3a)}, C_{(3b)}, C_{(3c)}$ respectively, then, we get the desired result for $s(\mathcal{G}_{n,r}^1)$ as follows:

$$s(\mathcal{G}_{n,r}^1) = C_{(1)} \cup C_{(2)} \cup C_{(3a)} \cup C_{(3b)} \cup C_{(3c)}$$

□

Our next result is the characterization of the f -vector of $\Delta_s(\mathcal{G}_{n,r}^1)$.

Proposition 3.5. Let $\Delta_s(\mathcal{G}_{n,r}^1)$ be a spanning simplicial complex of the graph $\mathcal{G}_{n,r}^1$, then the $\dim(\Delta_s(\mathcal{G}_{n,r}^1)) = n-r-1$ with f -vector $f(\Delta_s(\mathcal{G}_{n,r}^1)) = (f_0, f_1, \dots, f_{n-r-1})$ and

$$f_i = \binom{n}{i+1} + \sum_{k=1}^r (-1)^k \left[\sum_{j=1}^k \sum_{k_j=0}^{r-1} \sum_{i_j=1}^{r-k_j} \left(n - \sum_{j=1}^k m_{i_j, i_j+1, \dots, i_j+k_j} + \sum_{u,v=1}^k |C_{i_u, i_u+1, \dots, i_u+k_u} \cap C_{i_v, i_v+1, \dots, i_v+k_v}| \right) \right]$$

where $0 \leq i \leq n-r-1$

Proof. Let E be the edge set of $\mathcal{G}_{n,r}^1$ and $C_{(1)}, C_{(2)}, C_{(3a)}, C_{(3b)}, C_{(3c)}$ are disjoint classes of spanning trees of $\mathcal{G}_{n,r}^1$ then from lemma 3.2 we have

$$s(\mathcal{G}_{n,r}^1) = C_{(1)} \cup C_{(2)} \cup C_{(3a)} \cup C_{(3b)} \cup C_{(3c)}$$

Therefore, by definition 2.4 we can write

$$\Delta_s(\mathcal{G}_{n,r}^1) = \langle C_{(1)} \cup C_{(2)} \cup C_{(3a)} \cup C_{(3b)} \cup C_{(3c)} \rangle$$

Since each facet $\hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)} = E(T_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)})$ is obtained by deleting exactly r edges from the edge set of $\mathcal{G}_{n,r}^1$, keeping in view lemma 3.2, therefore dimension of each facet is same i.e., $n-r-1$ (since $|\hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}| = n-r$) and hence dimension of $\Delta_s(\mathcal{G}_{n,r}^1)$ will be $n-r-1$. Also it is clear from the definition of $\Delta_s(\mathcal{G}_{n,r}^1)$ that it contains all those subsets of E which do not contain the sets $\{e_{11}, \dots, e_{1m_1}\}$ and $\{e_{(i-1)1}, e_{i1}, \dots, e_{im_i-1}\}$ for all $2 \leq i \leq r$, i.e., those subsets of E which do not contain any cycle in the graph $\mathcal{G}_{n,r}^1$.

Now by lemma 3.1 the total cycles in the graph $\mathcal{G}_{n,r}^1$ are $C_{i, i+1, \dots, i+k}$ $i \in \{1, 2, \dots, r-k\}$ and $0 \leq k \leq r-1$, and their total number is τ . Let F be any subset of E of order $i+1$ such that it does not contain any $C_{i, i+1, \dots, i+k}$ $i \in \{1, 2, \dots, r-k\}$ and $0 \leq k \leq r-1$, in it. The total

number of such F is indeed f_i . We use inclusion exclusion principle to find this number. Therefore,

$f_i =$ Total number of subsets of E of order $i+1$ not containing $C_{i,i+1,\dots,i+k}$;
 $i \in \{1, 2, \dots, r-k\}$ and $0 \leq k \leq r-1$.

By Inclusion Exclusion Principle we have,

$$f_i = \left(\text{Total number of subsets of } E \text{ of order } i+1 \right) - \left(\sum_{j=1}^1 \sum_{k_{s_j}=0}^{r-1} \sum_{i_{s_j}=1}^{r-k_{s_j}} \right. \\ \left. \text{number of subsets of } E \text{ of order } i+1 \text{ containing } C_{i_{s_j}, i_{s_j}+1, \dots, i_{s_j}+k_{s_j}} \right) + \\ \left(\sum_{j=1}^2 \sum_{k_{s_j}=0}^{r-1} \sum_{i_{s_j}=1}^{r-k_{s_j}} \text{number of subsets of } E \text{ of order } i+1 \text{ containing both} \right. \\ \left. C_{i_{s_j}, i_{s_j}+1, \dots, i_{s_j}+k_{s_j}} \right) - \dots + (-1)^\tau \left(\sum_{j=1}^\tau \sum_{k_{s_j}=0}^{r-1} \sum_{i_{s_j}=1}^{r-k_{s_j}} \text{number of subsets of } E \right. \\ \left. \text{of order } i+1 \text{ containing each } C_{i_{s_j}, i_{s_j}+1, \dots, i_{s_j}+k_{s_j}} \right)$$

This implies

$$f_i = \binom{n}{i+1} - \left[\sum_{j=1}^1 \sum_{k_{s_j}=0}^{r-1} \sum_{i_{s_j}=1}^{r-k_{s_j}} \binom{n - m_{i_{s_1}, i_{s_1}+1, \dots, i_{s_1}+k_{s_1}}}{i+1 - m_{i_{s_1}, i_{s_1}+1, \dots, i_{s_1}+k_{s_1}}} \right] + \\ \left[\sum_{j=1}^2 \sum_{k_{s_j}=0}^{r-1} \sum_{i_{s_j}=1}^{r-k_{s_j}} \left(\binom{n - \sum_{j=1}^2 m_{i_{s_j}, i_{s_j}+1, \dots, i_{s_j}+k_{s_j}} + \sum_{u,v=1}^2 |C_{i_{s_u}, i_{s_u}+1, \dots, i_{s_u}+k_{s_u}} \cap C_{i_{s_v}, i_{s_v}+1, \dots, i_{s_v}+k_{s_v}}|}{i+1 - \sum_{j=1}^2 m_{i_{s_j}, i_{s_j}+1, \dots, i_{s_j}+k_{s_j}} + \sum_{u,v=1}^2 |C_{i_{s_u}, i_{s_u}+1, \dots, i_{s_u}+k_{s_u}} \cap C_{i_{s_v}, i_{s_v}+1, \dots, i_{s_v}+k_{s_v}}|} \right) \right] \\ - \dots + (-1)^\tau \\ \left[\sum_{j=1}^\tau \sum_{k_{s_j}=0}^{r-1} \sum_{i_{s_j}=1}^{r-k_{s_j}} \left(\binom{n - \sum_{j=1}^\tau m_{i_{s_j}, i_{s_j}+1, \dots, i_{s_j}+k_{s_j}} + \sum_{u,v=1}^\tau |C_{i_{s_u}, i_{s_u}+1, \dots, i_{s_u}+k_{s_u}} \cap C_{i_{s_v}, i_{s_v}+1, \dots, i_{s_v}+k_{s_v}}|}{i+1 - \sum_{j=1}^\tau m_{i_{s_j}, i_{s_j}+1, \dots, i_{s_j}+k_{s_j}} + \sum_{u,v=1}^\tau |C_{i_{s_u}, i_{s_u}+1, \dots, i_{s_u}+k_{s_u}} \cap C_{i_{s_v}, i_{s_v}+1, \dots, i_{s_v}+k_{s_v}}|} \right) \right]$$

This implies

$$f_i = \binom{n}{i+1} + \sum_{k=1}^\tau (-1)^k \\ \left[\sum_{j=1}^k \sum_{k_{s_j}=0}^{r-1} \sum_{i_{s_j}=1}^{r-k_{s_j}} \left(\binom{n - \sum_{j=1}^k m_{i_{s_j}, i_{s_j}+1, \dots, i_{s_j}+k_{s_j}} + \sum_{u,v=1}^k |C_{i_{s_u}, i_{s_u}+1, \dots, i_{s_u}+k_{s_u}} \cap C_{i_{s_v}, i_{s_v}+1, \dots, i_{s_v}+k_{s_v}}|}{i+1 - \sum_{j=1}^k m_{i_{s_j}, i_{s_j}+1, \dots, i_{s_j}+k_{s_j}} + \sum_{u,v=1}^k |C_{i_{s_u}, i_{s_u}+1, \dots, i_{s_u}+k_{s_u}} \cap C_{i_{s_v}, i_{s_v}+1, \dots, i_{s_v}+k_{s_v}}|} \right) \right]$$

□

Corollary 3.6. Let $\Delta_s(G_{n,2}^1)$ be a spanning simplicial complex of a graph with 2 cycles of lengths m_1, m_2 having one edge common, then the $\dim(\Delta_s(G_{n,2}^1)) = n-3$ with f -vectors $f(\Delta_s(G_{n,2}^1)) = (f_0, f_1, \dots, f_{n-3})$ and

$$f_i = \binom{n}{i+1} - \left[\binom{n-m_1}{i+1-m_1} + \binom{n-m_2}{i+1-m_2} + \binom{n-m_{1,2}}{i+1-m_{1,2}} \right] +$$

$$\left[\binom{n-m_1-m_2+|C_1 \cap C_2|}{i+1-m_1-m_2+|C_1 \cap C_2|} + \binom{n-m_1-m_{1,2}+|C_1 \cap C_{1,2}|}{i+1-m_1-m_{1,2}+|C_1 \cap C_{1,2}|} \right] +$$

$$\binom{n-m_2-m_{1,2}+|C_2 \cap C_{1,2}|}{i+1-m_2-m_{1,2}+|C_2 \cap C_{1,2}|} \Bigg] +$$

$$\left[\binom{n-m_1-m_2-m_{1,2}+|C_1 \cap C_2|+|C_1 \cap C_{1,2}|+|C_2 \cap C_{1,2}|}{i+1-m_1-m_2-m_{1,2}+|C_1 \cap C_2|+|C_1 \cap C_{1,2}|+|C_2 \cap C_{1,2}|} \right]$$

where $0 \leq i \leq n-3$.

For a simplicial complex Δ over $[n]$, one would associate to it the Stanley-Reisner ideal, that is, the monomial ideal $I_{\mathcal{N}}(\Delta)$ in $S = k[x_1, x_2, \dots, x_n]$ generated by monomials corresponding to non-faces of this complex (here we are assigning one variable of the polynomial ring to each vertex of the complex). It is well known that the face ring $k[\Delta] = S/I_{\mathcal{N}}(\Delta)$ is a standard graded algebra. We refer the readers to [8] and [13] for more details about graded algebra A , the Hilbert function $H(A, t)$ and the Hilbert series $H_t(A)$ of a graded algebra.

Our main result of this section is as follows;

Theorem 3.7. Let $\Delta_s(\mathcal{G}_{n,r}^1)$ be the spanning simplicial complex of $\mathcal{G}_{n,r}^1$, then the Hilbert series of the face ring $k[\Delta_s(\mathcal{G}_{n,r}^1)]$ is given by,

$$H(k[\Delta_s(\mathcal{G}_{n,r}^1)], t) = 1 + \sum_{i=0}^d \frac{\binom{n}{i+1} t^{i+1}}{(1-t)^{i+1}} + \sum_{i=0}^d \sum_{k=1}^{\tau} (-1)^k$$

$$\left[\sum_{j=1}^k \sum_{k_j=0}^{r-1} \sum_{i_j=1}^{r-k_j} \binom{n - \sum_{j=1}^k m_{i_j, i_j+1, \dots, i_j+k_j} + \sum_{u,v=1}^k |C_{i_u, i_u+1, \dots, i_u+k_u} \cap C_{i_v, i_v+1, \dots, i_v+k_v}|}{i+1 - \sum_{j=1}^k m_{i_j, i_j+1, \dots, i_j+k_j} + \sum_{u,v=1}^k |C_{i_u, i_u+1, \dots, i_u+k_u} \cap C_{i_v, i_v+1, \dots, i_v+k_v}|} \right]$$

$$\frac{t^{i+1}}{(1-t)^{i+1}}.$$

Proof. From [13], we know that if Δ is a simplicial complex of dimension d and $f(\Delta) = (f_0, f_1, \dots, f_d)$ its f -vector, then the Hilbert series of face ring $k[\Delta]$ is given by

$$H(k[\Delta], t) = 1 + \sum_{i=0}^d \frac{f_i t^{i+1}}{(1-t)^{i+1}}.$$

By substituting the values of f_i 's from Proposition 3.5 in this above expression, we get the desired result. \square

4 Associated primes of the facet ideal $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$

We present the characterization of all associated primes of the facet ideal $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$ of spanning simplicial complex $\Delta_s(\mathcal{G}_{n,r}^1)$ in this section.

Associated to a simplicial complex Δ over $[n]$, one defines the *facet ideal* $I_{\mathcal{F}}(\Delta) \subset S$, which is generated by square-free monomials $x_{i_1} \dots x_{i_s}$, where $\{v_{i_1}, \dots, v_{i_s}\}$ is a facet of Δ .

Lemma 4.1. If $\Delta_s(\mathcal{G}_{n,r}^1)$ be the spanning simplicial complex of the r -cycles graph $\mathcal{G}_{n,r}^1$, then

$$I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1)) = \left(\bigcap_{e_t \notin C_i}^{1 \leq i \leq r} (x_t) \right) \cap \left(\bigcap_{2 \leq i_\alpha \neq i_\beta \leq m_{j_\alpha} - 1}^{2 \leq j_\alpha \leq r-1} (x_{j_\alpha i_\alpha}, x_{j_\alpha i_\beta}) \right) \\ \cap \left(\bigcap_{2 \leq i_\alpha \neq i_\beta \leq m_1} (x_{1i_\alpha}, x_{1i_\beta}) \right) \cap \left(\bigcap_{1 \leq i_\alpha \neq i_\beta \leq m_r - 1} (x_{ri_\alpha}, x_{ri_\beta}) \right)$$

Proof. Consider the spanning simplicial complex $\Delta_s(\mathcal{G}_{n,r}^1)$ and let $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$ be the facet ideal of $\Delta_s(\mathcal{G}_{n,r}^1)$. Since from [4, Proposition 1.8], we know that a minimal prime ideal of the facet ideal $I_{\mathcal{F}}(\Delta)$ has one to one correspondence with the minimal vertex cover of the simplicial complex. Therefore, in order to compute the primary decomposition of the facet ideal $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$; it is sufficient to compute all the minimal vertex cover of $\Delta_s(\mathcal{G}_{n,r}^1)$.

Indeed clear from the definition of $\Delta_s(\mathcal{G}_{n,r}^1)$ and by Lemma 3.4 that $\{e_t\}$ is a minimal vertex cover of $\Delta_s(\mathcal{G}_{n,r}^1)$ such that $e_t \notin C_i \forall i \in \{1, \dots, r\}$. Moreover, $\{e_{j_\alpha i_\alpha}, e_{j_\alpha i_\beta}\}$ is also a minimal vertex cover of $\Delta_s(\mathcal{G}_{n,r}^1)$ with $2 \leq i_\alpha \neq i_\beta \leq m_{j_\alpha} - 1$ for $j_\alpha \in \{2, \dots, r-1\}$, $2 \leq i_\alpha \neq i_\beta \leq m_1$ for $j_\alpha = 1$ and $1 \leq i_\alpha \neq i_\beta \leq m_r - 1$ for $j_\alpha = r$. Indeed for any $\hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)} \in s(\mathcal{G}_{n,r}^1)$ the intersection $\{e_{j_\alpha i_\alpha}, e_{j_\alpha i_\beta}\} \cap \hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}$ is nonempty. \square

5 Cohen-Macaulayness of the face ring of $\Delta_s(\mathcal{G}_{n,r}^1)$

In this section, we include some definitions and results from [2] and use them to show that the face ring of $\Delta_s(\mathcal{G}_{n,r}^1)$ is Cohen-Macaulay.

Definition 5.1. [2]

Let $I \subset S = k[x_1, x_2, \dots, x_n]$ be a monomial ideal, we say that I will have the *quasi-linear quotients*, if there exists a minimal monomial system of generators m_1, m_2, \dots, m_r such that $\text{mindeg}(\hat{I}_{m_i}) = 1$ for all $1 < i \leq r$, where

$$\hat{I}_{m_i} = (m_1, m_2, \dots, m_{i-1}) : (m_i).$$

Theorem 5.2. [2] Let Δ be a pure simplicial complex of dimension d over $[n]$. Then Δ will be a shellable simplicial complex if and only if $I_{\mathcal{F}}(\Delta)$ will have the quasi-linear quotients.

Corollary 5.3. [2] If the facet ideal $I_{\mathcal{F}}(\Delta)$ of a pure simplicial complex Δ over $[n]$ has quasi-linear quotients, then the face ring is Cohen Macaulay.

Theorem 5.4. The face ring of $\Delta_s(\mathcal{G}_{n,r}^1)$ is Cohen-Macaulay.

Proof. By corollary 5.3, it is sufficient to show that $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$ has a quasi-linear quotients in $S = k[x_{11}, x_{12}, \dots, x_{1m_1}, x_{21}, x_{22}, \dots, x_{2(m_2-1)}, \dots, x_{r1}, x_{r2}, \dots, x_{r(m_r-1)}, x_1, x_2, \dots, x_t]$. By lemma 3.4, we have

$$s(\mathcal{G}_{n,r}^1) = C_{(1)} \cup C_{(2)} \cup C_{(3a)} \cup C_{(3b)} \cup C_{(3c)}$$

Therefore,

$$\Delta_s(\mathcal{G}_{n,r}^1) = \langle \hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)} = E \setminus \{e_{j_1 i_1}, e_{j_2 i_2}, \dots, e_{j_r i_r}\} \mid \hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)} \in s(\mathcal{G}_{n,r}^1) \rangle$$

and hence we can write,

$$I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1)) = \left(x_{\hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}} \mid \hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)} \in s(\mathcal{G}_{n,r}^1) \right).$$

Here, $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$ is a pure monomial ideal of degree $n - r$ with $x_{\hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}}$ as the product of all variables in S except $x_{j_1 i_1}, x_{j_2 i_2}, \dots, x_{j_r i_r}$. Now we will show that $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$ has quasi-linear quotients with respect to the following generating system:

$$\begin{aligned} & \{x_{\hat{E}_{(11, 21, \dots, r1)}}\}, \{x_{\hat{E}_{(11, 21, \dots, (r-1)1, j_r i_r)}} \mid i_r \neq 1\}, \{x_{\hat{E}_{(11, 21, \dots, (r-2)1, (r-1)i_{r-1}, j_r i_r)}} \mid \\ & i_{r-1} \neq 1\}, \\ & \{x_{\hat{E}_{(11, 21, \dots, (r-3)1, (r-2)i_{r-2}, j_{r-1}i_{r-1}, j_r i_r)}} \mid i_{r-2} \neq 1\}, \dots, \{x_{\hat{E}_{(11, 2i_2, j_3 i_3, \dots, j_r i_r)}} \mid \\ & i_2 \neq 1\}, \\ & \{x_{\hat{E}_{(1i_1, j_2 i_2, \dots, j_r i_r)}} \mid i_1 \neq 1\} \end{aligned}$$

Let us put

$$\begin{aligned} C_{(11, 21, \dots, (r-1)1, j_r i_r)} &= \{x_{\hat{E}_{(11, 21, \dots, (r-1)1, j_r i_r)}} \mid i_r \neq 1\}, \\ C_{(11, 21, \dots, (r-2)1, (r-1)i_{r-1}, j_r i_r)} &= \{x_{\hat{E}_{(11, 21, \dots, (r-2)1, (r-1)i_{r-1}, j_r i_r)}} \mid i_{r-1} \neq 1\}, \\ & \vdots \\ C_{(1i_1, j_2 i_2, \dots, j_r i_r)} &= \{x_{\hat{E}_{(1i_1, j_2 i_2, \dots, j_r i_r)}} \mid i_1 \neq 1\}. \end{aligned}$$

Also for any $C_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}$, denote $\bar{C}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}$ as the residue collection of all the generators which precedes $C_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}$ in the above order. We will show that

$$(\bar{C}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}) : (x_{\hat{E}_{(j_1 i_1, j_2 i_2, \dots, j_r i_r)}})$$

contains atleast one linear generator.

Now for any generator $x_{\hat{E}_{(11, \dots, (k-1)1, j_k i_k, \dots, j_r i_r)}}$, the above said system of generators guarantee the existence of a generator

$x_{\hat{E}_{(11, \dots, (k-1)1, j_\alpha i_\alpha, j_{k+1} i_{k+1}, \dots, j_r i_r)}}$ in $\bar{C}_{(11, \dots, (k-1)1, j_k i_k, \dots, j_r i_r)}$ such that $j_\alpha i_\alpha \neq j_k i_k$. Therefore, by using the definition of colon ideal it is easy to see that

$$(\bar{C}_{(11, \dots, (k-1)1, j_k i_k, \dots, j_r i_r)}) : (x_{(11, \dots, (k-1)1, j_k i_k, \dots, j_r i_r)})$$

contains a linear generator $x_{j_k i_k}$. Hence $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{n,r}^1))$ has quasi-linear quotients, as required. \square

We conclude this section with an example.

Example 5.5. For the graph $\mathcal{G}_{10,2}^1$ given in Fig. 1., the facet ideal of the spanning simplicial complex is:

$$I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{10,2}^1)) = (x_{11,21}, C_{11,j_2 i_2}, C_{1i_1,j_2 i_2})$$

where $C_{11,j_2 i_2} = x_{11,22}, x_{11,23}, x_{11,12}, x_{11,13}$ and $C_{1i_1,j_2 i_2} = x_{12,21}, x_{12,22}, x_{12,23}, x_{13,21}, x_{13,22}, x_{13,23}$. It is easy to see that, $I_{\mathcal{F}}(\Delta_s(\mathcal{G}_{10,2}^1))$ has quasi-linear quotients with respect to the ordering given to its generators (by applying above theorem). Hence, the face ring of $\Delta_s(\mathcal{G}_{10,2}^1)$ is Cohen Macaulay.

Acknowledgments: The authors are grateful to the reviewer and editor for their valuable suggestions to improve the manuscript.

References

- [1] Anwar I., Raza Z. and Kashif A. Spanning Simplicial Complexes of Uni-Cyclic Graphs. Algebra Colloquium 2015; 22(4) : 707 – 710.
- [2] Anwar I., and Raza Z. Quasi-Linear Quotients and Shellability Of Pure Simplicial Complexes. Communications in Algebra 2015; 43 : 4698 – 4704.
- [3] Bruns W. and Herzog J. Cohen Macaulay Rings. Vol.39, Cambridge studies in advanced mathematics, revised edition, 1998.
- [4] Faridi S. The Facet Ideal of a Simplicial Complex. Manuscripta Mathematica 2002; 109 : 159 – 174.
- [5] Faridi S. Simplicial Tree are Sequentially Cohen-Macaulay. J. Pure and Applied Algebra 2004; 190 : 121 – 136.

- [6] Guo J. and Wu T. On Spanning Complex of a Finite Graph. 2015; http://www.researchgate.net/publication/262070008_Spanning-Simplicial_Complex.
- [7] Harary F. Graph Theory. Reading, MA: Addison-Wesley. 1994.
- [8] Herzog J. and Hibi T. Monomial Algebra. Springer-Verlag New York Inc. 2009.
- [9] Kashif A., Anwar I. and Raza Z. On The Algebraic Study of Spanning Simplicial Complex of r -Cycles Graphs $G_{n,r}$. ARS Combinatoria 2014; 115 : 89 – 99.
- [10] Miller M. and Sturmfels B. Combinatorial Commutative Algebra. Springer-Verlag New York Inc. 2005.
- [11] Pan Y., Li R. and Zhu G. Spanning Simplicial Complexes of n -Cyclic Graphs with a Common Vertex. International Electronic Journal of Algebra. 2015; 17 : 180 – 187.
- [12] Raza Z., Kashif A., and Anwar I. On Algebraic Characterization of SSC of the Jahangir's Graph $\mathcal{J}_{n,m}$. to appear in Open Mathematics 2018.
- [13] Villarreal RH. Monomial Algebras. Dekker, New York. 2001.
- [14] Zhu G., Shi F. and Geng Y. Spanning Simplicial Complexes of n -Cyclic Graphs With a Common Edge. International Electronic Journal of Algebra. 2014; 15 : 132 – 144.